

Mirror Fermat Calabi-Yau Threefolds and Landau-Ginzburg Black Hole Attractors

Stefano Bellucci[♠], Sergio Ferrara^{◇, ✕, ♠}, Alessio Marrani^{♡, ♠} and Armen Yeranyan^{♣, ♠}

♠ *INFN - Laboratori Nazionali di Frascati,
Via Enrico Fermi 40, 00044 Frascati, Italy
bellucci, marrani@lnf.infn.it*

◇ *Physics Department, Theory Unit, CERN,
CH 1211, Geneva 23, Switzerland
sergio.ferrara@cern.ch*

✕ *Department of Physics and Astronomy,
University of California, Los Angeles, CA USA
ferrara@physics.ucla.edu*

♡ *Museo Storico della Fisica e
Centro Studi e Ricerche "Enrico Fermi"
Via Panisperna 89A, 00184 Roma, Italy*

♣ *Department of Physics, Yerevan State University,
Alex Manoogian St., 1, Yerevan, 375025, Armenia
ayeran@ysu.am*

ABSTRACT

We study black hole attractor equations for one-(complex structure)modulus Calabi-Yau spaces which are the mirror dual of Fermat Calabi-Yau threefolds (CY_3 s).

When exploring non-degenerate solutions near the Landau-Ginzburg point of the moduli space of such 4-dimensional compactifications, we always find two species of extremal black hole attractors, depending on the choice of the $Sp(4, \mathbb{Z})$ symplectic charge vector, one $\frac{1}{2}$ -BPS (which is always stable, according to general results of special Kähler geometry) and one non-BPS. The latter turns out to be stable (local minimum of the “effective black hole potential” V_{BH}) for non-vanishing central charge, whereas it is unstable (saddle point of V_{BH}) for the case of vanishing central charge.

This is to be compared to the large volume limit of one-modulus CY_3 -compactifications (of Type II A superstrings), in which the homogeneous symmetric special Kähler geometry based on cubic prepotential admits (beside the $\frac{1}{2}$ -BPS ones) only non-BPS extremal black hole attractors with non-vanishing central charge, which are always stable.

Contents

1	Introduction	3
2	Summary of Special Kähler Geometry	5
2.1	n_V -Moduli	5
2.2	1-Modulus	9
3	Stability of Critical Points of V_{BH}	11
3.1	n_V -Moduli	11
3.2	1-Modulus	14
3.2.1	$\frac{1}{2}$ -BPS	15
3.2.2	Non-BPS, $Z \neq 0$	15
3.2.3	Non-BPS, $Z = 0$	17
4	Holomorphic Geometry and Picard-Fuchs Equations	19
4.1	n_V -Moduli	20
4.2	1-Modulus	24
5	Holomorphic Geometry and Flat Holomorphic Matrix Connection	28
5.1	n_V -Moduli	28
5.2	1-Modulus	30
6	General Analysis of (Mirror) Fermat Calabi-Yau Threefolds near the Landau-Ginzburg Point	32
7	$k = 5$: Mirror <i>Quintic</i> near LG Point	41
7.1	Geometric Setup	41
7.1.1	The Holomorphic Prepotentials F and $\mathcal{F}$	43
7.2	“Criticality Condition” Approach and Attractor Equation	44
7.3	Critical Hessian of V_{BH}	44
7.4	Solutions to Attractor Equation	45
7.4.1	Analysis of non-BPS, $Z \neq 0$ supporting BH charge configurations	48
8	$k = 6$: Mirror <i>Sextic</i> near LG Point	51
8.1	Geometric Setup	51
8.1.1	The Holomorphic Prepotentials F and $\mathcal{F}$	52
8.2	“Criticality Condition” Approach and Attractor Equation	53

8.3	Critical Hessian of V_{BH}	54
8.4	Solutions to Attractor Equations	54
9	$k = 8$: Mirror <i>Octic</i> near LG Point	57
9.1	Geometric Setup	57
9.1.1	The Holomorphic Prepotentials F and $\mathcal{F}$	59
9.2	“Criticality Condition” Approach and Attractor Equation	60
9.3	Critical Hessian of V_{BH}	60
9.4	Solutions to Attractor Equations	61
9.4.1	Analysis of non-BPS, $Z \neq 0$ supporting BH charge configurations	64
10	$k = 10$: Mirror <i>Dectic</i> near LG Point	66
10.1	Geometric Setup	66
10.1.1	The Holomorphic Prepotentials F and $\mathcal{F}$	68
10.2	“Criticality Condition” Approach and Attractor Equation	69
10.3	Critical Hessian of V_{BH}	70
10.4	Solutions to Attractor Equations	71
11	Special Kähler Geometrical Identities and Attractors	72
11.1	$n_V = 1$ Identities near the LG Point	72
11.2	“Special Kähler Geometrical Identities” Approach	
	to LG Attractors in Fermat CY_3 s	75
11.2.1	$\frac{1}{2}$ -BPS	76
11.2.2	Non-BPS, $Z \neq 0$	76
11.2.3	Non-BPS, $Z = 0$	77
12	The (LG Limit of) Yukawa Coupling Function of Fermat CY_3s	77
12.1	Different Approaches to Computation and LG Limit	77
12.2	Check of Stability Conditions	81
12.3	The Exact Holomorphic Yukawa Coupling Function of Fermat CY_3 s	82
13	Conclusions and Outlook	82

1 Introduction

Extremal black hole (BH) attractors [1]-[4] have been recently widely investigated [5]- [35], especially in connection with new classes of solutions to the attractor equations corresponding to non-BPS (Bogomol’ny-Prasad-Sommerfeld) horizon geometries, supported by particular configurations of the BH electric and magnetic charges. Such geometries are *non-degenerate*, *i.e.* they have a finite, non-vanishing horizon area, and their Bekenstein-Hawking entropy [36] is obtained by extremizing an “effective BH potential”.

In $\mathcal{N} = 2$, $d = 4$ Maxwell-Einstein supergravity theories (MESGTs), non-degenerate attractor horizon geometries correspond to BH solitonic states belonging to “short massive multiplets” (for the $\frac{1}{2}$ -BPS case, with $0 < |Z|_H = M_{ADM,H}$) and to “long massive multiplets”, either with non-vanishing or vanishing central charge Z not saturating the BPS bound¹ [37]

$$0 \leq |Z|_H < M_{ADM,H}. \quad (1.1)$$

The Arnowitt-Deser-Misner (ADM) mass [38] at the BH horizon is obtained by extremizing a positive-definite “effective BH potential”² $V_{BH}(\phi, \tilde{\Gamma})$, where $\tilde{\Gamma}$ is the $1 \times (2n_V + 2)$ symplectic vector of the BH field-strength fluxes, *i.e.* of the magnetic and electric charges:

$$\tilde{\Gamma} \equiv (p^\Lambda, q_\Lambda), \quad p^\Lambda \equiv \frac{1}{4\pi} \int_{S_\infty^2} \mathcal{F}^\Lambda, \quad q_\Lambda \equiv \frac{1}{4\pi} \int_{S_\infty^2} \mathcal{G}_\Lambda, \quad \Lambda = 0, 1, \dots, n_V, \quad (1.2)$$

where n_V denotes the number of Abelian vector supermultiplets coupled to the supergravity one (containing the graviphoton A^0); moreover, $\mathcal{F}^\Lambda = dA^\Lambda$, and $\mathcal{G}_\Lambda$ is the “dual” field-strength two-form [48, 49].

The BH entropy S_{BH} is given by the Bekenstein-Hawking entropy-area formula [36, 39]

$$S_{BH}(\tilde{\Gamma}) = \frac{A_H(\tilde{\Gamma})}{4} = \pi V_{BH}(\phi, \tilde{\Gamma}) \Big|_{\partial V_{BH}=0} = \pi V_{BH}(\phi_H(\tilde{\Gamma})), \quad (1.3)$$

where A_H is the event horizon area, and the solution $\phi_H(\tilde{\Gamma})$ to the criticality condition

$$\partial_\phi V_{BH}(\phi, \tilde{\Gamma}) = 0 \quad (1.4)$$

is properly named *attractor* if the critical $(2n_V + 2) \times (2n_V + 2)$ real symmetric Hessian matrix

$$\left. \frac{\partial^2 V_{BH}(\phi, \tilde{\Gamma})}{\partial \phi \partial \phi} \right|_{\phi=\phi_H(\tilde{\Gamma})} \quad (1.5)$$

is a strictly positive-definite matrix³.

¹Here and in what follows, the subscript “ H ” will denote values at the BH event horizon.

²Here and below “ ϕ ” denotes the set of real scalars relevant for Attractor Mechanism, *i.e.* the $2n_V$ ones coming from the n_V vector supermultiplets coupled to the supergravity multiplet.

³The opposite is in general not true, *i.e.* there can be attractor points corresponding to critical Hessian matrices with “flat” directions (*i.e.* vanishing eigenvalues). In general, when a critical Hessian matrix exhibits some vanishing eigenvalues, one has to look at higher-order derivatives of V_{BH} evaluated at the considered point, and study their sign. Depending on the values of the supporting BH charges, one can obtain stable or unstable critical points. Examples in literature of investigations beyond the Hessian level can be found in [9, 24, 25].

Although non-supersymmetric BH attractors exist also in $\mathcal{N} > 2$, $d = 4$ and $d = 5$ supergravities [40, 19], the most interesting examples arise in $\mathcal{N} = 2$, $d = 4$ MESGTs, where the scalar fluctuations relevant for the BH Attractor Mechanism parametrize a special Kähler (SK) manifold. Recently, the classification of “attractor solutions” for extremal BHs has been performed in full generality for the whole class of homogeneous symmetric SK geometries [22], and three distinct classes of extremal BH attractors (namely $\frac{1}{2}$ -BPS, non-BPS $Z \neq 0$ and non-BPS $Z = 0$ ones) were found as solutions to Eqs. (1.4). In such a framework, the non-BPS charge orbits have been found to depend on whether the supporting charge vector $\tilde{\Gamma}$ is such that the $\mathcal{N} = 2$ central charge vanishes or not. Moreover, the critical Hessian matrix (1.5) was usually found to exhibit zero modes (*i.e.* “flat” directions), whose attractor nature seemingly further depends on additional conditions on the charge vector $\tilde{\Gamma}$, other than the ones given by the extremality conditions (1.4) (see *e.g.* [9]).

The aim of the present work is to study a particular class of (1-modulus) SK geometries, namely the ones underlying the complex structure moduli space of (mirror) Fermat Calabi-Yau threefolds (CY_3 s) (classified by the *Fermat parameter* $k = 5, 6, 8, 10$, and firstly found in [42]). The fourth order linear Picard-Fuchs (PF) ordinary differential equations determining the holomorphic fundamental period 4×1 vector for such a class of 1-modulus CY_3 s were found some time ago for $k = 5$ in [43, 44] (see in particular Eq. (3.9) of [43]; see also [45]) and for $k = 6, 8, 10$ in [46] (see also [47]).

In the present work we will make distinction between two different (but, as we will explicitly check in the above-mentioned context, completely equivalent) approaches to extremal BH attractors, namely:

- i)* the so-called “*criticality condition*” approach, based on the extremization of a suitably defined, positive-definite “effective BH potential” $V_{BH}(\phi, \tilde{\Gamma})$, whose criticality conditions (1.4) are called “*Attractor Equations*”;
- ii)* the so-called “*SK geometrical identities*” approach (exploited in Sect. 11), based on the evaluation of some fundamental geometrical identities of SK geometry along the *loci* of the scalar manifold determined by some “critical constraints”.

The plan of the paper is as follows.

In Sect. 2 we sketchily recall the fundamentals of the local SK geometry of the scalar manifolds of $\mathcal{N} = 2$, $d = 4$ MESGTs. After the treatment of the general n_V -moduli case in Subsect. 2.1, in Subsect. 2.2 we focus on the 1-modulus case.

Thence, Sect. 3 is devoted to the issue of the stability of the critical points of V_{BH} . After a general discussion in Subsect. 3.1 (including a treatment of the $\frac{1}{2}$ -BPS attractors, specialized for $n_V = 1$ in Subsubsect. 3.2.1), in Subsect. 3.2 we determine (sufficient, but in general not necessary) conditions for non-BPS $Z \neq 0$ (Subsubsect. 3.2.2) and non-BPS $Z = 0$ (Subsubsect. 3.2.3) (classes of) critical points of V_{BH} to be (*at least local*) minima of V_{BH} , and thus to be *attractors* in a strict sense.

Sects. 4 and 5 are devoted to present the holomorphic geometry embedded in SK geometry. Such a geometry is relevant in order to introduce the PF ordinary linear differential Eqs.. Once again, the 1-modulus case is considered

in detail in Subsects 4.2 and 5.2.

Then, in Sect. 6 the formalism of (mirror) Fermat CY_3 s (classified by the Fermat parameter $k = 5, 6, 8, 10$), in particular near the LG point $\psi = 0$ of their (complex structure deformation) moduli space, is introduced. The general analysis of Sect. 6 is consequently specialized to the study of non-degenerate extremal BH LG attractors in the complex structure moduli space of the four mirror Fermat CY_3 s, corresponding to $k = 5$ (Sect. 7), $k = 6$ (Sect. 8), $k = 8$ (Sect. 9), and $k = 10$ (Sect. 10).

The abovementioned “*SK geometrical identities*” approach is exploited in Sect. 11, in order to study the extremal BH LG attractors for the above-mentioned class of CY_3 s. After an introduction to $n_V = 1$ SK geometrical identities near the LG point in Subsect. 11.1, in Subsect. 11.2 their evaluation along the *loci* corresponding to the various species of critical points of V_{BH} is performed. By analyzing the $\frac{1}{2}$ -BPS, non-BPS $Z \neq 0$ and non-BPS $Z = 0$ classes respectively in Subsubsects. 11.2.1, 11.2.2 and 11.2.3, we obtain results identical to the ones we got in Sects. 7-10 by exploiting the abovementioned “*criticality condition*” approach, corresponding to solve near the LG point the 2 real *criticality conditions* of V_{BH} , which in the 1-modulus case are nothing but the real and imaginary part of the so-called $\mathcal{N} = 2$, $d = 4$ supergravity AEs.

Sect. 12 deals with the Yukawa coupling function of 1-modulus SK geometry. Subsect. 12.1 presents three different, equivalent approaches to compute such a crucial quantity, and apply them to study the LG limit of the moduli space of (mirror) Fermat CY_3 s. In such a framework, the 1-modulus stability conditions for non-BPS $Z \neq 0$ and non-BPS $Z = 0$ critical points of V_{BH} (respectively obtained in Subsubsects. 3.2.2 and 3.2.3) are thence explicitly checked in Subsect. 12.2, obtaining results consistent with the ones of Sects. 7-10. Finally, Subsect. 12.3 deals with the global, exact expression of the holomorphic Yukawa coupling function of the 1-modulus SK geometry underlying the moduli space of (mirror) Fermat CY_3 s, and with its limits near the three typologies of regular singular points of the PF differential Eqs. introduced in Sects. 4 and 5.

Concluding remarks, summarizing observations and outlooking comments are the contents of the final Sect. 13.

2 Summary of Special Kähler Geometry

2.1 n_V -Moduli

In $\mathcal{N} = 2$, $d = 4$ MESGT the following formula holds⁴ [3, 4, 48]

$$V_{BH}(z, \bar{z}; q, p) = |Z|^2(z, \bar{z}; q, p) + g^{j\bar{j}}(z, \bar{z}) D_j Z(z, \bar{z}; q, p) \overline{D_{\bar{j}} Z}(z, \bar{z}; q, p). \quad (2.1.1)$$

Consequently, the criticality conditions (1.4) can be easily shown to acquire the form [39]

$$2\overline{Z}D_i Z + g^{j\bar{j}}(D_i D_j Z) \overline{D_{\bar{j}} Z} = 0; \quad (2.1.2)$$

⁴Here and below we switch to the complex parametrization of the set of scalars being considered:

$$\{\phi^a\}_{a=1,\dots,2n_V} \longrightarrow \{z^i, \bar{z}^{\bar{i}}\}_{i,\bar{i}=1,\dots,n_V}.$$

The relation between such two equivalent parametrizations of the SK scalar manifold is given by Eq. (4.2) of [15].

this is what one should rigorously refer to as the $\mathcal{N} = 2$, $d = 4$ supergravity attractor equations (AEs). $g^{j\bar{j}}(z, \bar{z})$ is the contravariant Kähler metric tensor, satisfying the usual orthonormality condition:

$$g^{i\bar{j}}(z, \bar{z}) \partial_i \bar{\partial}_{\bar{k}} K(z, \bar{z}) = \delta_{\bar{k}}^{\bar{j}}, \quad (2.1.3)$$

where $K(z, \bar{z})$ is the real Kähler potential. As previously mentioned, $Z(z, \bar{z}; q, p)$ is the $\mathcal{N} = 2$ central charge function

$$Z(z, \bar{z}; q, p) \equiv e^{\frac{1}{2}K(z, \bar{z})} \tilde{\Gamma} \Omega \Pi(z) = e^{\frac{1}{2}K(z, \bar{z})} [q_\Lambda X^\Lambda(z) - p^\Lambda F_\Lambda(z)] \equiv e^{\frac{1}{2}K(z, \bar{z})} W(z; q, p), \quad (2.1.4)$$

where Ω is the $(2n_V + 2)$ -dim. symplectic metric (subscripts denote dimensions)

$$\Omega \equiv \begin{pmatrix} 0_{n_V+1} & -\mathbb{I}_{n_V+1} \\ \mathbb{I}_{n_V+1} & 0_{n_V+1} \end{pmatrix}, \quad (2.1.5)$$

and $\Pi(z)$ is the $(2n_V + 2) \times 1$ holomorphic period vector in symplectic basis

$$\Pi(z) \equiv \begin{pmatrix} X^\Lambda(z) \\ F_\Lambda(z) \end{pmatrix}, \quad (2.1.6)$$

with $X^\Lambda(z)$ and $F_\Lambda(z)$ being the holomorphic sections of the $U(1)$ line (Hodge) bundle over the SK manifold (clearly, due to holomorphicity they do not belong to the related $U(1)$ ring). Finally, $W(z; q, p)$ is the holomorphic $\mathcal{N} = 2$ central charge function, also named $\mathcal{N} = 2$ superpotential.

Let us here recall that Z has Kähler weights $(p, \bar{p}) = (1, -1)$; thus, its Kähler-covariant derivatives read

$$\begin{aligned} D_i Z &= (\partial_i + \tfrac{1}{2} \partial_i K) Z, \\ \bar{D}_{\bar{i}} Z &= (\bar{\partial}_{\bar{i}} - \tfrac{1}{2} \bar{\partial}_{\bar{i}} K) Z. \end{aligned} \quad (2.1.7)$$

The non-holomorphic basic, defining differential relations of SK geometry are⁵ (see *e.g.* [48]):

$$\left\{ \begin{array}{l} D_i Z = Z_i; \\ D_i Z_j = i C_{ijk} g^{k\bar{k}} \bar{D}_{\bar{k}} \bar{Z} = i C_{ijk} g^{k\bar{k}} \bar{Z}_{\bar{k}}; \\ D_i \bar{D}_{\bar{j}} \bar{Z} = D_i \bar{Z}_{\bar{j}} = g_{i\bar{j}} \bar{Z}; \\ D_i \bar{Z} = 0, \end{array} \right. \quad (2.1.8)$$

where the first relation is nothing but the definition of the “matter charges” Z_i s and the fourth relation expresses the Kähler-covariant holomorphicity of Z . C_{ijk} is the rank-3, completely symmetric, covariantly holomorphic tensor of

⁵Actually, there are different (equivalent) defining approaches to SK geometry. For subtleties and further elucidation concerning such an issue, see *e.g.* [51] and [52].

SK geometry (with Kähler weights $(2, -2)$) (see *e.g.* [48]- [54]):

$$\left\{ \begin{array}{l} C_{ijk} = C_{(ijk)} = \langle D_i D_j V, D_k V \rangle = e^K (\partial_i \mathcal{N}_{\Lambda\Sigma}) D_j X^\Lambda D_k X^\Sigma = \\ = e^K (\partial_i X^\Lambda) (\partial_j X^\Sigma) (\partial_k X^\Xi) \partial_\Xi \partial_\Sigma F_\Lambda(X) \equiv e^K W_{ijk}, \quad \bar{\partial}_{\bar{l}} W_{ijk} = 0; \\ C_{ijk} = D_i D_j D_k \mathcal{S}, \quad \mathcal{S} \equiv -i L^\Lambda L^\Sigma \text{Im}(F_{\Lambda\Sigma}), \quad F_{\Lambda\Sigma} \equiv \frac{\partial F_\Lambda}{\partial X^\Sigma}, F_{\Lambda\Sigma} \equiv F_{(\Lambda\Sigma)}; \\ C_{ijk} = -i g_{i\bar{l}} \bar{f}_\Lambda^{\bar{l}} D_j D_k L^\Lambda, \quad \bar{f}_\Lambda^{\bar{l}} (\overline{DL_s}^\Lambda) \equiv \delta_{\bar{s}}^{\bar{l}}; \end{array} \right. \quad (2.1.9)$$

$$\bar{D}_{\bar{i}} C_{jkl} = 0 \text{ (covariant holomorphicity);}$$

$$R_{i\bar{j}k\bar{l}} = -g_{i\bar{j}} g_{k\bar{l}} - g_{i\bar{l}} g_{k\bar{j}} + C_{ikp} \bar{C}_{\bar{j}\bar{l}\bar{p}} g^{p\bar{p}} \text{ (usually named SKG constraints);}$$

$$D_{[i} C_{j]kl} = 0,$$

where round (square) brackets denote (anti)symmetrization with respect to enclosed indices, and $R_{i\bar{j}k\bar{l}}$ is the Riemann-Christoffel tensor⁶ of Kähler geometry:

$$R_{i\bar{j}k\bar{l}} \equiv g^{m\bar{n}} \left(\bar{\partial}_{\bar{l}} \bar{\partial}_{\bar{j}} \partial_m K \right) \partial_i \bar{\partial}_{\bar{n}} \partial_k K - \bar{\partial}_{\bar{l}} \partial_i \bar{\partial}_{\bar{j}} \partial_k K = g_{k\bar{n}} \partial_i \bar{\Gamma}_{\bar{l}\bar{j}}^{\bar{n}} = g_{n\bar{l}} \bar{\partial}_{\bar{j}} \Gamma_{ki}^n, \quad \overline{R_{i\bar{j}k\bar{l}}} = R_{j\bar{i}l\bar{k}}; \quad (2.1.10)$$

$$\Gamma_{ij}^l \equiv -g^{l\bar{l}} \partial_i g_{j\bar{l}} = -g^{l\bar{l}} \partial_i \bar{\partial}_{\bar{l}} \partial_j K = \Gamma_{(ij)}^l, \quad (2.1.11)$$

Γ_{ij}^l standing for the Christoffel symbols of the second kind of the Kähler metric $g_{i\bar{j}}$. For later convenience, it is here worth writing the expression for the holomorphic covariant derivative of C_{ijk} :

$$D_i C_{jkl} = D_{(i} C_{j)kl} = \partial_i C_{jkl} + \Gamma_{ij}^m C_{mkl} + \Gamma_{ik}^m C_{mjl} + \Gamma_{il}^m C_{mjk}, \quad (2.1.12)$$

where we used the last of Eqs. (2.1.9), which is a consequence, through the SKG constraints and the covariant holomorphicity of C_{ijk} , of the Bianchi identities for the Riemann tensor $R_{i\bar{j}k\bar{l}}$ (see *e.g.* [53]).

By using the first two of relations (2.1.8), the $\mathcal{N} = 2$ AEs (2.1.2) can be recast as follows [39]:

$$2\bar{Z}Z_i + iC_{ijk}g^{j\bar{j}}g^{k\bar{k}}\bar{Z}_{\bar{j}}\bar{Z}_{\bar{k}} = 0. \quad (2.1.13)$$

It is now worth recalling some fundamental identities defining the geometric structure of SK manifolds [55, 8, 13, 15, 16, 24]

$$\tilde{\Gamma}^T - i\Omega\mathcal{M}(\mathcal{N})\tilde{\Gamma}^T = e^{K/2} \left[-2iZ\bar{\Pi} - 2ig^{j\bar{j}} \left(\bar{D}_{\bar{j}}\bar{Z} \right) D_j \Pi \right], \quad (2.1.14)$$

where $\mathcal{M}(\mathcal{N})$ denotes the $(2n_V + 2) \times (2n_V + 2)$ real symmetric matrix [48, 3, 4]

$$\mathcal{M}(\mathcal{N}) \equiv \begin{pmatrix} \text{Im}(\mathcal{N}) + \text{Re}(\mathcal{N}) (\text{Im}(\mathcal{N}))^{-1} \text{Re}(\mathcal{N}) & -\text{Re}(\mathcal{N}) (\text{Im}(\mathcal{N}))^{-1} \\ -(\text{Im}(\mathcal{N}))^{-1} \text{Re}(\mathcal{N}) & (\text{Im}(\mathcal{N}))^{-1} \end{pmatrix}, \quad (2.1.15)$$

⁶Notice that the third of Eqs. (2.1.9) correctly defines the Riemann tensor $R_{i\bar{j}k\bar{l}}$, and it is actual the opposite of the one which may be found in a large part of existing literature (see *e.g.* [50]). Such a formulation of the so-called *SKG constraints* is well defined, because, as discussed *e.g.* in [33], it yields negative values of the constant scalar curvature of $(n_V + 1\text{-dim.})$ homogeneous symmetric compact SK manifolds.

where $\mathcal{N}_{\Lambda\Sigma}$ is a $(n_V + 1) \times (n_V + 1)$ complex symmetric, moduli-dependent, Kähler gauge-invariant matrix, the so-called *kinetic matrix* of $\mathcal{N} = 2$, $d = 4$ MESGT (see *e.g.* the report [48]). Moreover, it should be here reminded that

$$\begin{aligned} D_i \Pi &= (\partial_i + \partial_i K) \Pi, \\ \overline{D}_{\bar{i}} \Pi &= \overline{\partial}_{\bar{i}} \Pi = 0, \end{aligned} \tag{2.1.16}$$

since Π is holomorphic with Kähler weights $(2, 0)$.

The $2n_V + 2$ complex identities (2.1.14) (whose real and imaginary parts are related by a suitable “rotation” [16]) express nothing but a *change of basis* of the BH charge configurations, between the Kähler-invariant $1 \times (2n_V + 2)$ symplectic (magnetic/electric) basis vector $\tilde{\Gamma}$ defined by Eq. (1.2) and the complex, moduli-dependent $1 \times (n_V + 1)$ *matter charges* vector (with Kähler weights $(1, -1)$)

$$\mathcal{Z}(z, \bar{z}) \equiv (Z(z, \bar{z}), Z_i(z, \bar{z}))_{i=1, \dots, n_V}. \tag{2.1.17}$$

It should be recalled that the BH charges are conserved due to the overall $(U(1))^{n_V+1}$ gauge-invariance of the system under consideration, and $\tilde{\Gamma}$ and $\mathcal{Z}(z, \bar{z})$ are two *equivalent* basis for them. Their very equivalence relations are given by the SKG identities (2.1.14) themselves. By its very definition (1.2), $\tilde{\Gamma}$ is *moduli-independent* (at least in a stationary, spherically symmetric and asymptotically flat extremal BH background, as it is the case being treated here), whereas Z is *moduli-dependent*, since it refers to the eigenstates of the $\mathcal{N} = 2$, $d = 4$ supergravity multiplet and of the n_V Maxwell vector supermultiplets.

Moreover, it is important to stress that identities (2.1.14) entail 2 redundant degrees of freedom, encoded in the homogeneity (of degree 1) of (2.1.14) under complex rescalings of $\tilde{\Gamma}$. Indeed, by recalling the definition (2.1.4) it can be readily checked that the right-hand side of (2.1.14) acquires an overall factor λ under the rescaling

$$\tilde{\Gamma} \longrightarrow \lambda \tilde{\Gamma}, \quad \lambda \in \mathbb{C}. \tag{2.1.18}$$

We will reconsider such a point in Sect. 11, when treating the 1-modulus case more in detail.

It should also be noticed that the $\mathcal{N} = 2$ “effective BH potential” given by Eq. (2.1.1) can also be rewritten as [3, 4, 48]

$$V_{BH}(z, \bar{z}; q, p) = -\frac{1}{2} \tilde{\Gamma} \mathcal{M}(\mathcal{N}) \tilde{\Gamma}^T, \tag{2.1.19}$$

and therefore it can be identified with the first, positive-definite real invariant of SK geometry (see *e.g.* [24, 48]). It is interesting to remark that the result (2.1.19) can be elegantly obtained from the SK geometry identities (2.1.14) by making use of the following relations [19]:

$$\mathcal{M}(\mathcal{N}) \Pi = i \Omega \Pi, \quad \mathcal{M}(\mathcal{N}) \overline{D}_{\bar{j}} \Pi = i \Omega \overline{D}_{\bar{j}} \Pi, \quad \forall \bar{j}. \tag{2.1.20}$$

By introducing the $(2n_V + 2) \times (n_V + 1)$ rectangular matrix ($I = 1, \dots, n_V, n_V + 1$)

$$\mathcal{V} \equiv (\overline{D}_{\bar{i}} \overline{\Pi}, \Pi) = \begin{pmatrix} \overline{D}_{\bar{i}} \overline{X}^\Lambda, X^\Lambda \\ \overline{D}_{\bar{i}} \overline{F}_\Lambda, F_\Lambda \end{pmatrix} \equiv e^{-K/2} \begin{pmatrix} \overline{D}_{\bar{i}} \overline{L}^\Lambda, L^\Lambda \\ \overline{D}_{\bar{i}} \overline{M}_\Lambda, M_\Lambda \end{pmatrix} = e^{-K/2} \begin{pmatrix} f_I^\Lambda \\ h_{I\Lambda} \end{pmatrix}, \quad (2.1.21)$$

$$f_I^\Lambda \equiv (\overline{D}_{\bar{i}} \overline{L}^\Lambda, L^\Lambda), \quad h_{I\Lambda} \equiv (\overline{D}_{\bar{i}} \overline{M}_\Lambda, M_\Lambda), \quad (2.1.22)$$

the relations (2.1.20) can be recast as⁷

$$\mathcal{M}(\mathcal{N}) \mathcal{V} = i\Omega \mathcal{V} \Leftrightarrow \frac{1}{2} (\mathcal{M}(\mathcal{N}) + i\Omega) \mathcal{V} = i\Omega \mathcal{V}. \quad (2.1.23)$$

It is also here worth recalling that the fundamental *Ansätze* [48] of SKG (solving the *SKG constraints* given by the third of Eqs. (2.1.9))

$$M_\Lambda = \mathcal{N}_{\Lambda\Sigma} L^\Sigma, \quad D_i M_\Lambda = \overline{\mathcal{N}}_{\Lambda\Sigma} D_i L^\Sigma. \quad (2.1.24)$$

univoquely determine $\mathcal{N}_{\Lambda\Sigma}(z, \bar{z})$ as

$$\mathcal{N}_{\Lambda\Sigma}(z, \bar{z}) = h_{I\Lambda}(z, \bar{z}) \circ (f^{-1})_\Sigma^I(z, \bar{z}), \quad (2.1.25)$$

where $\circ$ denotes the usual matrix product, $(f^{-1})_\Sigma^I f_I^\Lambda = \delta_\Sigma^\Lambda$, $(f^{-1})_\Lambda^I f_J^\Lambda = \delta_J^I$, and f_I^Λ and $h_{I\Lambda}$ are $(n_V + 1) \times (n_V + 1)$ complex matrices defined by Eqs. (2.1.22).

2.2 1-Modulus

In the 1-modulus case ($n_V = 1$) major simplifications occur in the formalism of SKG.

The potential V_{BH} (2.1.1) and the AEs (given by Eqs. (2.1.2) and (2.1.13)) respectively reduce to⁸ ($z^1 \equiv \psi$, $D_1 \equiv D_\psi \equiv D$, $g_{\psi\bar{\psi}} \equiv g$, $C_{111} \equiv C_{\psi\psi\psi} \equiv C$)

$$\begin{aligned} V_{BH}(\psi, \bar{\psi}; q, p) &\equiv |Z|^2(\psi, \bar{\psi}; q, p) + g^{-1}(\psi, \bar{\psi}) |DZ|^2(\psi, \bar{\psi}; q, p) = \\ &= e^{K(\psi, \bar{\psi})} \left[|W|^2(\psi, \bar{\psi}; q, p) + g^{-1}(\psi, \bar{\psi}) |DW|^2(\psi, \bar{\psi}; q, p) \right]; \end{aligned} \quad (2.2.1)$$

$$2\overline{W}DW + g^{-1}(D^2W)\overline{D\overline{W}} = 2\overline{W}DW + iCg^{-2}(\overline{D\overline{W}})^2 = 0, \quad (2.2.2)$$

where the SK geometry has been assumed to be *regular* (*i.e.* with $|K_k(\psi, \bar{\psi})| < \infty$ globally holding), and the (holomorphic) Kähler weight 2 of the holomorphic superpotential W has been exploited:

$$DW = \partial W + (\partial K)W = e^{-K/2} DZ = e^{-K/2} D(e^{K/2} W). \quad (2.2.3)$$

⁷Up to an exchange of order of L^Λ, M_Λ and $\overline{D}_{\bar{i}} \overline{L}^\Lambda, \overline{D}_{\bar{i}} \overline{M}_\Lambda$ in the definitions (2.1.22), Eq. (2.1.23) matches the $\mathcal{N} \geq 2$ generalization of SKG (and its application to the case $\mathcal{N} = 2$) performed in Sect. 2 of [19].

⁸In the present paper C will be used to denote two different entities in different contexts:

1) the Yukawa coupling function in 1-dim. SK geometry, which is nothing but the tensor C_{ijk} (defined by Eqs. (2.1.9)) for $n_V = 1$ (see also Sect. 12);

2) some (k -parameterized) constants in the general formalism of 1-dim. SK geometry arising from compactifications on (mirror) Fermat CY_3 s, as given by the first of definitions (6.12).

Attention should be paid to correctly identify case-by-case the meaning of C , which however can be effortlessly understood in the various contexts.

By specializing⁹ the definition (2.1.11) for $n_V = 1$:

$$\Gamma_{11}^{-1} \equiv \Gamma_{\psi\psi}^{\psi} \equiv \Gamma = -g^{-1}\partial g = -\partial \ln(g), \quad (2.2.4)$$

one can compute

$$\begin{aligned} D^2W &= \partial DW + (\partial K) DW + \Gamma DW = \\ &= \partial [\partial W + (\partial K) W] + (\partial K) [\partial W + (\partial K) W] - [\partial \ln(g)] DW = \\ &= \left\{ \partial^2 + \partial^2 K + 2\partial K \partial + (\partial K)^2 - [\partial \ln(\partial \bar{\partial} K)] (\partial + \partial K) \right\} W. \end{aligned} \quad (2.2.5)$$

Thus, the three different classes of non-degenerate, critical points of V_{BH} are in order:

$\frac{1}{2}$ -**BPS class**, *i.e.* the geometrical *locus* defined by the constraints

$$Z \neq 0, \quad DZ = 0 \iff W \neq 0, \quad DW = 0, \quad (2.2.6)$$

which directly solve the AEs (2.2.2).

Non-BPS, $Z \neq 0$ class, *i.e.* the geometrical *locus* defined by

$$Z \neq 0, \quad DZ \neq 0 \iff W \neq 0, \quad DW \neq 0, \quad (2.2.7)$$

further constrained to solve AEs (2.2.2), and thus yielding the following relations [15] ($k \in \mathbb{Z}$)

$$\begin{aligned} D^2W &= -2g\bar{W} \frac{DW}{DW} \\ &\iff \\ |D^2W| &= 2g|W|, \quad 2\varphi_{DW} = \varphi_{D^2W} + \varphi_W + (1+2k)\pi; \end{aligned} \quad (2.2.8)$$

$$\begin{aligned} DW &= -\frac{i}{2}Cg^{-2} \frac{(\bar{DW})^2}{W} = -\frac{i}{2}C\bar{W}g^{-2} \overline{D \ln(W)}^2 \\ &\iff \\ |DW| &= 2g^2 \frac{|W|}{|C|}, \quad \varphi_{DW} = \varphi_C + \varphi_W + \left(\frac{3}{2} + 2k\right)\pi, \end{aligned} \quad (2.2.9)$$

where the φ s stand for phases. Notice that Eqs. (2.2.2) and (2.2.7) necessarily yield that

$$C_{non-BPS, Z \neq 0} \neq 0. \quad (2.2.10)$$

For this class, one can also define the *supersymmetry-breaking order parameter* as follows:

$$\mathcal{O}_{non-BPS, Z \neq 0} \equiv \left[\frac{g^{-1}|DZ|^2}{|Z|^2} \right]_{non-BPS, Z \neq 0} = \left[\frac{g^{-1}|DW|^2}{|W|^2} \right]_{non-BPS, Z \neq 0} = \quad (2.2.11)$$

$$= 4 \left[\frac{g^3}{|C|^2} \right]_{non-BPS, Z \neq 0}, \quad (2.2.12)$$

⁹In the present paper Γ will be used to denote different entities in different contexts:

- 1) the Christoffel affine connection of (1-dim.) Kähler geometry, as defined by Eq. (2.1.11) ((2.2.4));
- 2) (one of the forms of) the symplectic charge vector, as given by the first of definitions (6.28);
- 3) the Euler Gamma-function, as defined by Eq. (6.13).

Attention should be paid to carefully identify the meaning of Γ , which however can be understood at a glance in the various frameworks. It should also be recalled that $\tilde{\Gamma}$ denotes the symplectic charge vector, as defined by Eq. (1.2), and $\hat{\Gamma}$ denotes the (holomorphic) Christoffel connection of (1-dim.) holomorphic geometry, as defined by Eqs. (4.1.24)-(4.1.25) (the first of definitions (4.2.2)).

where in the second line we used Eq. (2.2.9). It is worth noticing that for a cubic prepotential (in a suitable special projective coordinate, with Kähler gauge fixed such that $X^0 = 1$) $\mathcal{F}(z) = \varrho z^3$ it holds that $\mathcal{O}_{non-BPS, Z \neq 0} = 3 \forall \varrho \in \mathbb{C}$ [22]; such a result actually holds for cubic prepotentials in generic n_V -moduli SK geometries, such as the ones arising in the large volume limit of CY_3 -compactifications of Type II A superstring theory (see Eq. (111) of [9]).

Non-BPS, $Z = 0$ class, *i.e.* the geometrical *locus* defined by

$$Z = 0, \quad DZ \neq 0 \iff W = 0, \quad DW \neq 0, \quad (2.2.13)$$

further constrained to solve AEs (2.2.2), and thus yielding

$$D^2 W|_{non-BPS, Z=0} = 0 \xLeftrightarrow{DW|_{non-BPS, Z=0} \neq 0} C_{non-BPS, Z=0} = 0. \quad (2.2.14)$$

Along the *loci* of its critical points, the “BH effective potential” V_{BH} respectively reads

$$V_{BH, \frac{1}{2}-BPS} = |Z|_{\frac{1}{2}-BPS}^2; \quad (2.2.15)$$

$$V_{BH, non-BPS, Z \neq 0} = |Z|_{non-BPS, Z \neq 0}^2 \left[1 + 4 \frac{g^3}{|C|^2} \right]_{non-BPS, Z \neq 0}; \quad (2.2.16)$$

$$V_{BH, non-BPS, Z=0} = |DZ|_{non-BPS, Z=0}^2. \quad (2.2.17)$$

In the present paper we will investigate AE (2.2.2) near one of three typologies of regular singular points in the complex structure moduli space of (mirror) Fermat CY_3 s, namely near the so-called Landau-Ginzburg (LG) point $\psi = 0$. In such a framework, the identities (2.1.14) of SK geometry, when considered in the 1-modulus case and in correspondence of the various above-mentioned species of critical points of V_{BH} , can be used to find the BH charge configurations supporting the LG point $\psi = 0$ to be an attractor point of the considered kind. It will be shown that, in spite of the fact that identities (2.1.14) give 4 real Eqs. in the 1-modulus case, only 2 of them are independent, and they are completely equivalent to the 2 real rigorously-named $\mathcal{N} = 2$, $d = 4$ supergravity AEs (2.2.2), which are nothing but the criticality condition $\partial V_{BH} = 0$.

3 Stability of Critical Points of V_{BH}

3.1 n_V -Moduli

In order to decide whether a critical point of V_{BH} is an attractor in strict sense, one has to consider the following condition:

$$H_{\mathbb{R}}^{V_{BH}} \equiv H_{ab}^{V_{BH}} \equiv D_a D_b V_{BH} > 0 \quad \text{at} \quad D_c V_{BH} = \frac{\partial V_{BH}}{\partial \phi^c} = 0 \quad \forall c = 1, \dots, 2n_V, \quad (3.1.1)$$

i.e. the condition of (strict) positive-definiteness of the real $2n_V \times 2n_V$ Hessian matrix $H_{\mathbb{R}}^{V_{BH}} \equiv H_{ab}^{V_{BH}}$ of V_{BH} (which is nothing but the squared mass matrix of the moduli) at the critical points of V_{BH} , expressed in the real parameterization

through the ϕ -coordinates. Since V_{BH} is positive-definite, a stable critical point (namely, an attractor in strict sense) is necessarily a (n at least local) minimum, and therefore it fulfills the condition (3.1.1).

In general, $H_{\mathbb{R}}^{V_{BH}}$ may be block-decomposed in $n_V \times n_V$ real matrices:

$$H_{\mathbb{R}}^{V_{BH}} = \begin{pmatrix} \mathcal{A} & \mathcal{C} \\ \mathcal{C}^T & \mathcal{B} \end{pmatrix}, \quad (3.1.2)$$

with $\mathcal{A}$ and $\mathcal{B}$ being $n_V \times n_V$ real symmetric matrices:

$$\mathcal{A}^T = \mathcal{A}, \mathcal{B}^T = \mathcal{B} \Leftrightarrow \left(H_{\mathbb{R}}^{V_{BH}}\right)^T = H_{\mathbb{R}}^{V_{BH}}. \quad (3.1.3)$$

In the local complex $(z, \bar{z})$ -parameterization, the $2n_V \times 2n_V$ Hessian matrix of V_{BH} reads

$$H_{\mathbb{C}}^{V_{BH}} \equiv H_{\hat{i}\hat{j}}^{V_{BH}} \equiv \begin{pmatrix} D_i D_j V_{BH} & D_i \bar{D}_{\bar{j}} V_{BH} \\ D_j \bar{D}_{\bar{i}} V_{BH} & \bar{D}_{\bar{i}} \bar{D}_{\bar{j}} V_{BH} \end{pmatrix} = \begin{pmatrix} \mathcal{M}_{ij} & \mathcal{N}_{i\bar{j}} \\ \bar{\mathcal{N}}_{\bar{i}j} & \bar{\mathcal{M}}_{\bar{i}\bar{j}} \end{pmatrix}, \quad (3.1.4)$$

where the hatted indices $\hat{i}$ and $\hat{j}$ may be holomorphic or antiholomorphic. $H_{\mathbb{C}}^{V_{BH}}$ is the matrix actually computable in the SKG formalism presented in Sect. 2 (see below, Eqs. (3.1.6) and (3.1.7)).

In general, $\frac{1}{2}$ -BPS critical points are (at least local) minima of V_{BH} , and therefore they are stable; thus, they are *attractors* in strict sense. Indeed, the $2n_V \times 2n_V$ (covariant) Hessian matrix $H_{\mathbb{C}}^{V_{BH}}$ evaluated at such points is strictly positive-definite [39] :

$$\begin{aligned} (D_i D_j V_{BH})_{\frac{1}{2}-BPS} &= (\partial_i \partial_j V_{BH})_{\frac{1}{2}-BPS} = 0, \\ \left(D_i \bar{D}_{\bar{j}} V_{BH}\right)_{\frac{1}{2}-BPS} &= \left(\partial_i \bar{\partial}_{\bar{j}} V_{BH}\right)_{\frac{1}{2}-BPS} = 2 \left(g_{i\bar{j}} V_{BH}\right)_{\frac{1}{2}-BPS} = 2 \left.g_{i\bar{j}}\right|_{\frac{1}{2}-BPS} |Z|_{\frac{1}{2}-BPS}^2 > 0, \end{aligned} \quad (3.1.5)$$

where here and below the notation “ > 0 ” (“ < 0 ”) is understood as strict positive-(negative-)definiteness. The Hermiticity and (strict) positive-definiteness of the (covariant) Hessian matrix $H_{\mathbb{C}}^{V_{BH}}$ at the $\frac{1}{2}$ -BPS critical points are due to the Hermiticity and - assumed - (strict) positive-definiteness (actually holding globally) of the metric $g_{i\bar{j}}$ of the SK scalar manifold being considered.

On the other hand, non-BPS critical points of V_{BH} does not automatically fulfill the condition (3.1.1), and a more detailed analysis [22, 16] is needed.

Using the properties of SKG, one obtains:

$$\begin{aligned} \mathcal{M}_{ij} &\equiv D_i D_j V_{BH} = D_j D_i V_{BH} = \\ &= 3 \bar{Z} D_i D_j Z + g^{k\bar{k}} (D_i D_j D_k Z) \bar{D}_{\bar{k}} \bar{Z} = e^K \left[3 \bar{W} D_i D_j W + g^{k\bar{k}} (D_i D_j D_k W) \bar{D}_{\bar{k}} \bar{W} \right] \\ &= 4 i \bar{Z} C_{ijk} g^{k\bar{k}} (\bar{D}_{\bar{k}} \bar{Z}) + i (D_j C_{ikl}) g^{k\bar{k}} g^{l\bar{l}} (\bar{D}_{\bar{k}} \bar{Z}) (\bar{D}_{\bar{l}} \bar{Z}) = \\ &= e^K \left[4 i \bar{Z} C_{ijk} g^{k\bar{k}} (\bar{D}_{\bar{k}} \bar{W}) + i (D_j C_{ikl}) g^{k\bar{k}} g^{l\bar{l}} (\bar{D}_{\bar{k}} \bar{W}) (\bar{D}_{\bar{l}} \bar{W}) \right]; \end{aligned} \quad (3.1.6)$$

$$\begin{aligned}
\mathcal{N}_{i\bar{j}} &\equiv D_i \bar{D}_{\bar{j}} V_{BH} = \bar{D}_{\bar{j}} D_i V_{BH} = \\
&= 2g_{i\bar{j}} |Z|^2 + 2(D_i Z) (\bar{D}_{\bar{j}} \bar{Z}) + g^{k\bar{k}} \left[(\bar{D}_{\bar{j}} D_i D_k Z) \bar{D}_{\bar{k}} \bar{Z} + (D_i D_k Z) \bar{D}_{\bar{j}} \bar{D}_{\bar{k}} \bar{Z} \right] = \\
&= e^K \left\{ 2g_{i\bar{j}} |W|^2 + 2(D_i W) (\bar{D}_{\bar{j}} \bar{W}) + g^{k\bar{k}} \left[(\bar{D}_{\bar{j}} D_i D_k W) \bar{D}_{\bar{k}} \bar{W} + (D_i D_k W) \bar{D}_{\bar{j}} \bar{D}_{\bar{k}} \bar{W} \right] \right\} = \\
&= 2 \left[g_{i\bar{j}} |Z|^2 + (D_i Z) (\bar{D}_{\bar{j}} \bar{Z}) + g^{l\bar{n}} C_{ikl} \bar{C}_{\bar{j}\bar{m}\bar{n}} g^{k\bar{k}} g^{m\bar{m}} (\bar{D}_{\bar{k}} \bar{Z}) (D_m Z) \right] = \\
&= 2e^K \left[g_{i\bar{j}} |W|^2 + (D_i W) (\bar{D}_{\bar{j}} \bar{W}) + g^{l\bar{n}} C_{ikl} \bar{C}_{\bar{j}\bar{m}\bar{n}} g^{k\bar{k}} g^{m\bar{m}} (\bar{D}_{\bar{k}} \bar{W}) (D_m W) \right],
\end{aligned} \tag{3.1.7}$$

with $D_j C_{ikl}$ given by Eq. (2.1.12). Clearly, evaluating Eqs. (3.1.6) and (3.1.7) constrained by the $\frac{1}{2}$ -BPS conditions $D_i Z = 0, \forall i = 1, \dots, n_V$, one reobtains the results (3.1.5). Here we limit ourselves to point out that further noteworthy elaborations of $\mathcal{M}_{ij}$ and $\mathcal{N}_{i\bar{j}}$ can be performed in homogeneous symmetric SK manifolds, where $D_j C_{ikl} = 0$ globally [22], and that the Kähler-invariant $(2, 2)$ -tensor $g^{l\bar{n}} C_{ikl} \bar{C}_{\bar{j}\bar{m}\bar{n}}$ can be rewritten in terms of the Riemann-Christoffel tensor $R_{i\bar{j}k\bar{m}}$ by using the so-called “SKG constraints” (see the third of Eqs. (2.1.9)) [16]. Moreover, the differential Bianchi identities for $R_{i\bar{j}k\bar{m}}$ imply $\mathcal{M}_{ij}$ to be symmetric (see comment below Eqs. (2.1.9) and (2.1.12)).

Thus, one gets the following global properties:

$$\mathcal{M}^T = \mathcal{M}, \quad \mathcal{N}^\dagger = \mathcal{N} \Leftrightarrow \left(H_{\mathbb{C}}^{V_{BH}} \right)^T = H_{\mathbb{C}}^{V_{BH}}, \tag{3.1.8}$$

implying that

$$\left(H_{\mathbb{C}}^{V_{BH}} \right)^\dagger = H_{\mathbb{C}}^{V_{BH}} \Leftrightarrow \mathcal{M}^\dagger = \mathcal{M}, \quad \mathcal{N}^T = \mathcal{N} \Leftrightarrow \bar{\mathcal{M}} = \mathcal{M}, \quad \bar{\mathcal{N}} = \mathcal{N}. \tag{3.1.9}$$

It should be stressed clearly that the symmetry but non-Hermiticity of $H_{\mathbb{C}}^{V_{BH}}$ actually does not matter, because what one is interested in are the eigenvalues of the real form $H_{\mathbb{R}}^{V_{BH}}$, which is real and symmetric, and therefore admitting $2n_V$ real eigenvalues.

The relation between $H_{\mathbb{R}}^{V_{BH}}$ expressed by Eq. (3.1.2) and $H_{\mathbb{C}}^{V_{BH}}$ given by Eq. (3.1.4) is expressed by the following relations between the $n_V \times n_V$ sub-blocks of $H_{\mathbb{R}}^{V_{BH}}$ and $H_{\mathbb{C}}^{V_{BH}}$ [15, 33]:

$$\begin{cases} \mathcal{M} = \frac{1}{2} (\mathcal{A} - \mathcal{B}) + \frac{i}{2} (\mathcal{C} + \mathcal{C}^T); \\ \mathcal{N} = \frac{1}{2} (\mathcal{A} + \mathcal{B}) + \frac{i}{2} (\mathcal{C}^T - \mathcal{C}), \end{cases} \tag{3.1.10}$$

or its inverse

$$\begin{cases} \mathcal{A} = \text{Re} \mathcal{M} + \text{Re} \mathcal{N}; \\ \mathcal{B} = \text{Re} \mathcal{N} - \text{Re} \mathcal{M}; \\ \mathcal{C} = \text{Im} \mathcal{M} - \text{Im} \mathcal{N}. \end{cases} \tag{3.1.11}$$

The structure of the Hessian matrix gets simplified at the critical points of V_{BH} , because the covariant derivatives may be substituted by the flat ones; the critical Hessian matrices in complex holomorphic/antiholomorphic and real

local parameterizations respectively read

$$H_{\mathbb{C}}^{V_{BH}} \Big|_{\partial V_{BH}=0} \equiv \begin{pmatrix} \partial_i \partial_{\bar{j}} V_{BH} & \partial_i \bar{\partial}_{\bar{j}} V_{BH} \\ \partial_{\bar{j}} \bar{\partial}_i V_{BH} & \bar{\partial}_i \bar{\partial}_{\bar{j}} V_{BH} \end{pmatrix}_{\partial V_{BH}=0} = \begin{pmatrix} \mathcal{M} & \mathcal{N} \\ \bar{\mathcal{N}} & \bar{\mathcal{M}} \end{pmatrix}_{\partial V_{BH}=0}; \quad (3.1.12)$$

$$H_{\mathbb{R}}^{V_{BH}} \Big|_{\partial V_{BH}=0} = \frac{\partial^2 V_{BH}}{\partial \phi^a \partial \phi^b} \Big|_{\partial V_{BH}=0} = \begin{pmatrix} \mathcal{A} & \mathcal{C} \\ \mathcal{C}^T & \mathcal{B} \end{pmatrix}_{\partial V_{BH}=0}. \quad (3.1.13)$$

Thus, the study of the condition (3.1.1) finally amounts to the study of the *eigenvalue problem* of the real symmetric $2n_V \times 2n_V$ critical Hessian matrix $H_{\mathbb{R}}^{V_{BH}} \Big|_{\partial V_{BH}=0}$ given by Eq. (3.1.13), which is the Cayley-transformed of the complex (symmetric, but not necessarily Hermitian) $2n_V \times 2n_V$ critical Hessian $H_{\mathbb{C}}^{V_{BH}} \Big|_{\partial V_{BH}=0}$ given by Eq. (3.1.12).

3.2 1-Modulus

Once again, the situation strongly simplifies in $n_V = 1$ SKG.

Indeed, for $n_V = 1$ the moduli-dependent matrices $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{M}$ and $\mathcal{N}$ introduced above are simply scalar functions. In particular, $\mathcal{N}$ is real, since $\mathcal{C}$ trivially satisfies $\mathcal{C} = \mathcal{C}^T$. The stability condition (3.1.1) can thus be written as

$$H_{\mathbb{R}}^{V_{BH}} \equiv D_a D_b V_{BH} > 0, \quad (a, b = 1, 2) \quad \text{at} \quad D_c V_{BH} = \frac{\partial V_{BH}}{\partial \phi^c} = 0 \quad \forall c = 1, 2, \quad (3.2.1)$$

and Eqs. (3.1.6) and (3.1.7) respectively simplify to

$$\begin{aligned} \mathcal{M} &\equiv D^2 V_{BH} = \\ &= 3\bar{Z} D^2 Z + g^{-1} (D^3 Z) \bar{D}_{\bar{k}} \bar{Z} = e^K [3\bar{W} D^2 W + g^{-1} (D^3 W) \bar{D} \bar{W}] = \\ &= 4i\bar{Z} C g^{-1} \bar{D} \bar{Z} + i(DC) g^{-2} (\bar{D} \bar{Z})^2 = ie^K g^{-1} (\bar{D} \bar{W}) [4\bar{W} C + (DC) g^{-1} \bar{D} \bar{W}]; \end{aligned} \quad (3.2.2)$$

$$\begin{aligned} \mathcal{N} &\equiv D \bar{D} V_{BH} = \bar{D} D V_{BH} = \\ &= 2g |Z|^2 + 2|DZ|^2 + g^{-1} [(\bar{D} D^2 Z) \bar{D} \bar{Z} + |D^2 Z|^2] = \\ &= e^K \left\{ 2g |W|^2 + 2|DW|^2 + g^{-1} [(\bar{D} D^2 W) \bar{D} \bar{W} + |D^2 W|^2] \right\} = \\ &= 2 \left[g |Z|^2 + |DZ|^2 + |C|^2 g^{-3} |DZ|^2 \right] = 2e^K \left[g |W|^2 + |DW|^2 + |C|^2 g^{-3} |DW|^2 \right], \end{aligned} \quad (3.2.3)$$

where DC is given by the case $n_V = 1$ case of Eq. (2.1.12) (also recall definition (2.2.4)):

$$DC = \partial C + [(\partial K) + 3\Gamma] C = \partial C + [(\partial K) - 3\partial \ln(g)] C = \left\{ \partial + \left[\partial \ln \left(\frac{e^K}{(\bar{\partial} \partial K)^3} \right) \right] \right\} C. \quad (3.2.4)$$

It is easy to show that the stability condition (3.2.1) for critical points of V_{BH} in $n_V = 1$ SKG can be equivalently reformulated as the strict bound

$$\mathcal{N}|_{\partial V_{BH}=0} > |\mathcal{M}|_{\partial V_{BH}=0}. \quad (3.2.5)$$

Let us now see how such a bound can be further elaborated for the three possible classes of critical points of V_{BH} .

3.2.1 $\frac{1}{2}$ -BPS

$$\begin{aligned} \mathcal{M}_{\frac{1}{2}-BPS} &\equiv D^2 V_{BH}|_{\frac{1}{2}-BPS} = \\ &= \{e^K [3\bar{W} D^2 W + g^{-1} (D^3 W) \bar{D}\bar{W}]\}_{\frac{1}{2}-BPS} = 0; \end{aligned} \quad (3.2.1.1)$$

$$\begin{aligned} \mathcal{N}_{\frac{1}{2}-BPS} &\equiv D\bar{D} V_{BH}|_{\frac{1}{2}-BPS} = \\ &= \left[e^K \left\{ 2g |W|^2 + g^{-1} |D^2 W|^2 \right\} \right]_{\frac{1}{2}-BPS} = 2 \left[g e^K |W|^2 \right]_{\frac{1}{2}-BPS}. \end{aligned} \quad (3.2.1.2)$$

Eqs. (3.2.1.1) and (3.2.1.2) are nothing but the 1-modulus case of Eq. (3.1.5), and they directly satisfy the bound (3.2.5). Thus, consistently with what stated above, the $\frac{1}{2}$ -BPS class of critical points of V_{BH} actually is a class of attractors in strict sense (*at least* local minima of V_{BH}).

3.2.2 Non-BPS, $Z \neq 0$

$$\begin{aligned} \mathcal{M}_{non-BPS, Z \neq 0} &\equiv D^2 V_{BH}|_{non-BPS, Z \neq 0} = \\ &= \{e^K [3\bar{W} D^2 W + g^{-1} (D^3 W) \bar{D}\bar{W}]\}_{non-BPS, Z \neq 0} = \\ &= \left\{ e^K \left[-6g (DW) \frac{\bar{W}^2}{\bar{D}\bar{W}} + g^{-1} (D^3 W) \bar{D}\bar{W} \right] \right\}_{non-BPS, Z \neq 0} = \\ &= \left[e^K |DW|^2 \left(-6 \frac{g}{D \ln(W)^2} + g^{-1} \frac{D^3 W}{DW} \right) \right]_{non-BPS, Z \neq 0} = \\ &= i \left\{ e^K g^{-1} \left[4\bar{W} C \bar{D}\bar{W} + (DC) g^{-1} (\bar{D}\bar{W})^2 \right] \right\}_{non-BPS, Z \neq 0} = \\ &= -2 \left\{ e^K g^{-1} \bar{W} DW \left[g^{-2} |C|^2 D \ln(W) + g D \ln(C) \right] \right\}_{non-BPS, Z \neq 0} = \\ &= i \left\{ e^K C g^{-3} (\bar{D}\bar{W})^2 \left[g^{-2} |C|^2 D \ln(W) + g D \ln(C) \right] \right\}_{non-BPS, Z \neq 0}; \end{aligned} \quad (3.2.2.1)$$

$$\begin{aligned}
\mathcal{N}_{non-BPS,Z \neq 0} &\equiv D\bar{D}V_{BH}|_{non-BPS,Z \neq 0} = \bar{D}DV_{BH}|_{non-BPS,Z \neq 0} = \\
&= \left\{ e^K \left[2g|W|^2 + 2|DW|^2 + g^{-1}(\bar{D}D^2W)\bar{D}\bar{W} + g^{-1}|D^2W|^2 \right] \right\}_{non-BPS,Z \neq 0} = \\
&= \left\{ e^K |DW|^2 \left[2 + 6g \frac{|W|^2}{|DW|^2} + g^{-1} \frac{\bar{D}D^2W}{\bar{D}\bar{W}} \right] \right\}_{non-BPS,Z \neq 0} = \\
&= \left\{ e^K |DW|^2 \left[2 + \frac{3}{2}g^{-3}|C|^2 + g^{-1} \frac{\bar{D}D^2W}{\bar{D}\bar{W}} \right] \right\}_{non-BPS,Z \neq 0} = \\
&= 2 \left\{ e^K |DW|^2 \left[1 + \frac{5}{4}g^{-3}|C|^2 \right] \right\}_{non-BPS,Z \neq 0},
\end{aligned} \tag{3.2.2.2}$$

where D^2W is given by Eq. (2.2.5) and

$$\begin{aligned}
\bar{D}D^2W &= \bar{\partial}D^2W = \bar{\partial} \left\{ \left[\partial^2 + \partial^2 K + 2\partial K \partial + (\partial K)^2 - (\partial \ln(\partial \bar{\partial} K))(\partial + \partial K) \right] W \right\} = \\
&= g^{-2}|C|^2 DW \neq D\bar{D}DW = gDW;
\end{aligned} \tag{3.2.2.3}$$

$$\begin{aligned}
D^3W &= \partial D^2W + (\partial K) D^2W + 2\Gamma D^2W = \\
&= \partial \left\{ \left[\partial^2 + \partial^2 K + 2\partial K \partial + (\partial K)^2 - (\partial \ln(\partial \bar{\partial} K))(\partial + \partial K) \right] W \right\} + \\
&+ [(\partial K) - 2(\partial \ln(\partial \bar{\partial} K))] \left\{ \left[\partial^2 + \partial^2 K + 2\partial K \partial + (\partial K)^2 - (\partial \ln(\partial \bar{\partial} K))(\partial + \partial K) \right] W \right\}
\end{aligned} \tag{3.2.2.4}$$

Eq. (3.2.2.1) yields that

$$\begin{aligned}
|\mathcal{M}|_{non-BPS,Z \neq 0}^2 &= \left\{ e^{2K} |DW|^4 \left[36 \frac{g^2}{|D \ln(W)|^4} + g^{-2} \frac{|D^3W|^2}{|DW|^2} - 12 \operatorname{Re} \left(\frac{D^3W}{DW} \frac{1}{\overline{D \ln(W)}} \right) \right] \right\}_{non-BPS,Z \neq 0} = \\
&= \left\{ e^{2K} |DW|^2 \left[36g^2 \frac{|W|^4}{|DW|^2} + g^{-2} |D^3W|^2 - 12 \operatorname{Re} \left[\bar{W} (D^3W) \overline{D \ln(W)} \right] \right] \right\}_{non-BPS,Z \neq 0} = \\
&= 4 \left\{ e^{2K} |DW|^4 \left[|C|^4 g^{-6} + \frac{1}{4} g^{-4} |DC|^2 + 2g^{-3} \operatorname{Re} [C (\overline{DC}) D \ln(W)] \right] \right\}_{non-BPS,Z \neq 0}.
\end{aligned} \tag{3.2.2.5}$$

By substituting Eqs. (3.2.2.2) and (3.2.2.5) into the strict inequality (3.2.5), one finally obtains the stability condition

for non-BPS, $Z \neq 0$ critical points of V_{BH} in $n_V = 1$ SKG [15]:

$$\mathcal{N}_{non-BPS, Z \neq 0} > |\mathcal{M}|_{non-BPS, Z \neq 0}; \quad (3.2.2.6)$$

$$\Leftrightarrow$$

$$\left[2|DW| + 6g \frac{|W|^2}{|DW|} + g^{-1} (\bar{D}D^2W) \bar{D}\bar{W} \right]_{non-BPS, Z \neq 0} > \sqrt{\left\{ \begin{array}{l} 36g^2 \frac{|W|^4}{|DW|^2} + g^{-2} |D^3W|^2 + \\ -12Re \left[\bar{W} (D^3W) \bar{D}ln(W) \right] \end{array} \right\}}_{non-BPS, Z \neq 0} \quad (3.2.2.7)$$

$$\Leftrightarrow$$

$$1 + \frac{5}{4} \left(|C|^2 g^{-3} \right)_{non-BPS, Z \neq 0} > \sqrt{\left[|C|^4 g^{-6} + \frac{1}{4} g^{-4} |DC|^2 + 2g^{-3} Re \left[C (\bar{D}C) (\bar{D}ln\bar{Z}) \right] \right]_{non-BPS, Z \neq 0}}. \quad (3.2.2.8)$$

As it is seen from such a condition, in general $(DC)_{non-BPS, Z \neq 0}$ is the fundamental geometrical quantity playing a key role in determining the stability of non-BPS, $Z \neq 0$ critical points of V_{BH} in 1-modulus SK geometry.

3.2.3 Non-BPS, $Z = 0$

$$\begin{aligned} \mathcal{M}_{non-BPS, Z=0} &\equiv D^2V_{BH}|_{non-BPS, Z=0} = \\ &= [e^K g^{-1} (D^3W) \bar{D}\bar{W}]_{non-BPS, Z=0} = [e^K g^{-1} (D^3W) \bar{\partial}\bar{W}]_{non-BPS, Z=0} = \\ &= i [e^K g^{-2} (DC) (\bar{D}\bar{W})^2]_{non-BPS, Z=0} = i [e^K g^{-2} (\partial C) (\bar{\partial}\bar{W})^2]_{non-BPS, Z=0}; \end{aligned} \quad (3.2.3.1)$$

$$\begin{aligned} \mathcal{N}_{non-BPS, Z=0} &\equiv D\bar{D}V_{BH}|_{non-BPS, Z=0} = \\ &= \left\{ e^K \left[2|DW|^2 + g^{-1} (\bar{\partial}D^2W) \bar{D}\bar{W} \right] \right\}_{non-BPS, Z=0} = \left\{ e^K \left[2|\partial W|^2 + g^{-1} (\bar{\partial}D^2W) \bar{\partial}\bar{W} \right] \right\}_{non-BPS, Z=0} = \\ &= 2e^K |DW|_{non-BPS, Z=0}^2 = 2e^K |\partial W|_{non-BPS, Z=0}^2, \end{aligned} \quad (3.2.3.2)$$

where Eqs. (2.2.13) and (2.2.14) have been used. Eq. (3.2.3.1) yields that

$$|\mathcal{M}|_{non-BPS, Z=0} = [e^K g^{-1} |D^3W| |\partial W|]_{non-BPS, Z=0} = [e^K g^{-2} |\partial C| |\partial W|^2]_{non-BPS, Z=0}. \quad (3.2.3.3)$$

By substituting Eqs. (3.2.3.2) and (3.2.3.3) into the strict inequality (3.2.5), one finally obtains the stability condition

for non-BPS, $Z = 0$ critical points of V_{BH} in $n_V = 1$ SKG:

$$\mathcal{N}_{non-BPS, Z=0} > |\mathcal{M}|_{non-BPS, Z=0}; \quad (3.2.3.4)$$

$$\Updownarrow$$

$$2g_{non-BPS, Z=0} + \left(\frac{\bar{\partial} D^2 W}{\partial W} \right)_{non-BPS, Z=0} > \left| \frac{D^3 W}{\partial W} \right|_{non-BPS, Z=0}; \quad (3.2.3.5)$$

$$\Updownarrow$$

$$2g_{non-BPS, Z=0}^2 > |\partial C|_{non-BPS, Z=0}. \quad (3.2.3.6)$$

Even though the stability condition (3.2.3.6) have been obtained by correctly using Eqs. (2.2.13) and (2.2.14), holding at the non-BPS, $Z = 0$ critical points of V_{BH} , in some cases it may happen that, in the limit of approaching the non-BPS, $Z = 0$ critical point of V_{BH} , in DC (given by Eq. (3.2.4)) the “connection term” $[(\partial K) + 3\Gamma]C$ is not necessarily sub-leading with respect to the “differential term” ∂C . Thus, for later convenience, it is worth rewriting the condition (3.2.3.6) as follows:

$$2(\bar{\partial}\partial K)_{non-BPS, Z=0}^2 > |\{\partial + [(\partial K) - 3\partial \ln(g)]\}C|_{non-BPS, Z=0} = \left| \left\{ \partial + \left[\partial \ln \left(\frac{e^K}{(\bar{\partial}\partial K)^3} \right) \right] \right\} C \right|_{non-BPS, Z=0}. \quad (3.2.3.7)$$

Remark

Let us consider the 1-modulus stability conditions (3.2.2.8) and (3.2.3.6)-3.2.3.7). It is immediate to realize that they are both satisfied when the function C is globally covariantly constant:

$$DC = \partial C + [(\partial K) + 3\Gamma]C = 0, \quad (3.2.3.8)$$

i.e. for the so-called homogeneous symmetric ($\dim_{\mathbb{C}} = n_V = 1$) SK geometry [56, 57], univoquely associated to the coset manifold $\frac{SU(1,1)}{U(1)}$. Such a SK manifold can be twofold characterized as:

i) the $n = 0$ element of the irreducible rank-1 infinite sequence $\frac{SU(1,1+n)}{U(1) \otimes SU(1+n)}$ (with $n_V = n + rank = n + 1$), endowed with a quadratic holomorphic prepotential function reading (in a suitable projective special coordinate, with Kähler gauge fixed such that $X^0 = 1$; see [22] and Refs. therein)

$$\mathcal{F}(z) = \frac{i}{4} (z^2 - 1). \quad (3.2.3.9)$$

By recalling the first of Eqs. (2.1.9), such a prepotential yields $C = 0$ globally (and thus Eq. (3.2.3.8)), and therefore, by using the SKG constraints (*i.e.* the third of Eqs. (2.1.9)) it yields also the constant scalar curvature

$$\mathcal{R} \equiv g^{-2}R = -2, \quad (3.2.3.10)$$

where $R \equiv R_{\bar{1}\bar{1}\bar{1}}$ denotes the unique component of the Riemann tensor. As obtained in [22], quadratic (homogeneous symmetric) SK geometries only admit $\frac{1}{2}$ -BPS and non-BPS, $Z = 0$ critical points of V_{BH} . Thus, it can be concluded that the 1-dim. quadratic SK geometry determined by the prepotential (3.2.3.9) admits *all* stable critical points of V_{BH} .

ii) the $n = -2$ element of the reducible rank-3 infinite sequence $\frac{SU(1,1)}{U(1)} \otimes \frac{SO(2,2+n)}{SO(2) \otimes SO(2+n)}$ (with $n_V = n + \text{rank} = n + 3$). endowed with a cubic holomorphic prepotential function reading (in a suitable projective special coordinate, with Kähler gauge fixed such that $X^0 = 1$; see [22] and Refs. therein; see also observation below Eqs. (2.2.11)-(2.2.12))

$$\mathcal{F}(z) = \varrho z^3, \quad \varrho \in \mathbb{C}, \quad (3.2.3.11)$$

constrained by the condition $\text{Im}(z) < 0$. By recalling the first of Eqs. (2.1.9), such a prepotential yields $C = 6\varrho e^K$ (and thus Eq. (3.2.3.8)), and consequently it also yields the constant scalar curvature

$$\mathcal{R} \equiv g^{-2} R = g^{-2} \left(-2g^2 + g^{-1} |C|^2 \right) = -\frac{2}{3}, \quad (3.2.3.12)$$

where the SKG constraints (i.e. the third of Eqs. (2.1.9)) and the global value¹⁰ $|C|^2 g^{-3} = \frac{4}{3}$ have been used. As it can be computed (see *e.g.* [35]), the 1-dim. SK geometry determined by the prepotential (3.2.3.11) admits, beside the (stable) $\frac{1}{2}$ -BPS ones, stable non-BPS $Z \neq 0$ critical points of V_{BH} . Thus, it is another example in which *all* critical points of V_{BH} actually are attractors in a strict sense.

Clearly, the quadratic and cubic homogeneous symmetric 1-modulus SK geometries (respectively determined by holomorphic prepotentials (3.2.3.9) and (3.2.3.11)) are not the only ones (with $n_V = 1$) admitting stable non-BPS critical points of V_{BH} . As we will check in Sects. 7 and 9, for instance also the 1-modulus SK geometries of the moduli space of the (mirror) Fermat CY_3 *quintic* $\mathcal{M}'_5$ and *octic* $\mathcal{M}'_8$ admit, in a suitable neighbourhood of the LG point, stable non-BPS ($Z \neq 0$) critical points of V_{BH} .

4 Holomorphic Geometry and Picard-Fuchs Equations

In this and in the next Section we will present a summary of the holomorphic geometry embedded in the SK geometry of the scalar manifolds of $N = 2$, $d = 4$ MESGTs. The main references for such an issue are [58] and [59], to which we will refer at the relevant points of the treatment.

The PF Equations, satisfied in SK geometry by the holomorphic period vector (in a suitable basis, named *PF basis*) are a consequence of SK geometry and of the underlying symplectic structure of the *flat symplectic bundle* [60] (see also next Section), which encodes the differential relations obeyed by the covariantly holomorphic sections and their covariant derivatives.

¹⁰The global value $|C|^2 g^{-3} = \frac{4}{3}$ for homogeneous symmetric cubic $n_V = 1$ SK geometries actually is nothing but the $n_V = 1$ case of the general global relation holding in a generic homogeneous symmetric cubic n_V -dimensional SK geometry [57]:

$$C_{p(kl} C_{ij)n} g^{n\bar{n}} g^{p\bar{p}} \bar{C}_{\bar{n}\bar{p}\bar{m}} = C_{(p(kl} C_{ij)|n)} g^{n\bar{n}} g^{p\bar{p}} \bar{C}_{\bar{n}\bar{p}\bar{m}} = \frac{4}{3} g_{(l|\bar{m}} C_{|ijk)}.$$

4.1 n_V -Moduli

Let us start by considering the Kähler-covariantly holomorphic, symplectic $1 \times (2n_V + 2)$ vector¹¹

$$V(z, \bar{z}) \equiv (L^\Lambda(z, \bar{z}), M_\Lambda(z, \bar{z})) = e^{\frac{1}{2}K(z, \bar{z})} \Pi^T(z), \quad (4.1.1)$$

with $\Pi^T(z)$ defined by Eq. (2.1.6).

Flatness of the symplectic connection entails the following relations [60]:

$$\begin{cases} D_\alpha V = U_\alpha; \\ D_\alpha U_\beta = iC_{\alpha\beta\gamma} g^{\gamma\bar{\gamma}} \bar{D}_{\bar{\gamma}} \bar{V} = iC_{\alpha\beta\gamma} g^{\gamma\bar{\gamma}} \bar{U}_{\bar{\gamma}}; \\ D_\alpha \bar{D}_{\bar{\beta}} \bar{V} = D_\alpha \bar{U}_{\bar{\beta}} = g_{\alpha\bar{\beta}} \bar{V}; \\ D_\alpha \bar{V} = 0. \end{cases} \quad (4.1.2)$$

Notice that, by the definition (4.1.1), the $\mathcal{N} = 2$ central charge function (defined by Eq. (2.1.4)) can be rewritten (in the notation for period vectors used in the present Section) as $Z = \tilde{\Gamma} \Omega V^T$, and the defining relations (2.1.8) of SK geometry can thus be obtained by transposing the relations (4.1.2) and by further left-multiplying them by the $1 \times (2n_V + 2)$ vector $\tilde{\Gamma} \Omega$.

Let us now consider a new $1 \times (2n_V + 2)$ vector of holomorphic sections¹² ($a = 1, \dots, n_V$)

$$V_h(X(z)) \equiv (X^0(z), X^a(z), F_a(X(z)), -F_0(X(z))). \quad (4.1.3)$$

We notice that, while $V(z, \bar{z})$ defined in Eq. (4.1.1) is symplectic with respect to the symplectic metric Ω , this does not hold for $V_h(X(z))$ defined in Eq. (4.1.3), which is instead symplectic with respect to a newly defined anti-diagonal symplectic metric ($Q^T = -Q$, $Q^2 = -\mathbb{I}_{2n_V+2}$):

$$Q \equiv \begin{pmatrix} & & & 1 \\ & & -\mathbb{I}_{n_V} & \\ & \mathbb{I}_{n_V} & & \\ -1 & & & \end{pmatrix}, \quad (4.1.4)$$

where unwritten elements vanish.

In the treatment which follows we will assume the existence of an holomorphic prepotential $F(X(z))$ of $\mathcal{N} = 2$, $d = 4$ vector multiplet couplings such that

$$F_\Lambda(z) \equiv \frac{\partial F(X(z))}{\partial X^\Lambda}, \quad (4.1.5)$$

¹¹In order to make the contact with the relevant literature easier, in this Section, as well in the next one, we will change some notations with respect to the previous treatment.

Firstly, we will consider row (*i.e.* $1 \times (2n_V + 2)$), instead of column (*i.e.* $(2n_V + 2) \times 1$), period vectors.

Secondly, we will use lowercase Greek indices to denote homogeneous coordinates (instead of lowercase Latin indices, as done in the previous Section). Lowercase Latin indices will rather be used to denote indices pertaining to the so-called *holomorphic geometry* we are going to discuss.

¹²The subscript “ h ” stands for “holomorphic”.

which is in turn implied by the assumption that the holomorphic square matrix

$$e_\alpha^a(z) \equiv \frac{\partial \left[\frac{X^a(z)}{X^0(z)} \right]}{\partial z^\alpha} \equiv \frac{\partial t^a(z)}{\partial z^\alpha} = \frac{\partial_\alpha X^a(z)}{X^0(z)} - \frac{X^a(z)}{(X^0(z))^2} \partial_\alpha X^0(z) \quad (4.1.6)$$

is invertible (non-singular), where in the last step we introduced the *homogeneous* (Kähler-invariant) coordinates (see e.g. [48])

$$t^a(z) \equiv \frac{X^a(z)}{X^0(z)}. \quad (4.1.7)$$

The matrix $e_\alpha^a(z)$ is nothing but the Jacobian of the change of basis between the $t^a(z)$ s and the z^α s. *Special (symplectic) coordinates* correspond to the case $e_\alpha^a(z) = \delta_\alpha^a$, implying that $t^a(z) \equiv \frac{X^a(z)}{X^0(z)} = z^a$ (in such a case a -indices and α -indices do coincide). By further fixing the Kähler gauge such that $X^0 = 1$, one finally gets $t^a(z) = X^a(z) = z^a$ and $X^0 = 1$, which is the usual definition of special coordinates (yielding $\partial_\alpha X^\Lambda = \delta_\alpha^\Lambda$).

In special coordinates, the holomorphic period vector $V_h(X(z))$ introduced in Eq. (4.1.3) reads as follows (Kähler gauge $X^0 = 1$ fixed understood throughout, unless otherwise noted; here and below the subscript “*sp.*” stands for “special”):

$$V_{h,sp.}(z) \equiv (1, z^a, \partial_a \mathcal{F}(z), -\mathcal{F}_0(z)) = (1, z^a, \partial_a \mathcal{F}(z), z^a \partial_a \mathcal{F}(z) - 2\mathcal{F}(z)), \quad (4.1.8)$$

where

$$\mathcal{F}(z) \equiv F\left(\frac{X(z)}{X^0(z)}\right) = (X^0(z))^{-2} F(X(z)) \quad (4.1.9)$$

is the Kähler gauge-invariant holomorphic prepotential (here in special coordinates and for $X^0 = 1$). In the second step the homogeneity of degree 2 of the prepotential was used; for general symplectic and special ($X^0 = 1$) coordinates it respectively reads

$$X^\Lambda \partial_\Lambda F(X) = X^0 \partial_0 F(X) + X^a \partial_a F(X) = 2F(X); \quad (4.1.10)$$

$$\mathcal{F}_0(z) + z^a \partial_a \mathcal{F}(z) = 2\mathcal{F}(z).$$

By starting from Eq. (4.1.8) and by differentiating once and twice $V_{h,sp.}(z)$, one respectively achieves

$$\partial_b V_{h,sp.}(z) = (0, \delta_b^a, \partial_a \partial_b \mathcal{F}(z), -\partial_b \mathcal{F}(z) + z^a \partial_a \partial_b \mathcal{F}(z)); \quad (4.1.11)$$

$$\partial_b \partial_c V_{h,sp.}(z) = (0, 0, \partial_a \partial_b \partial_c \mathcal{F}(z), z^a \partial_a \partial_b \partial_c \mathcal{F}(z)), \quad (4.1.12)$$

implying that

$$\partial_b \partial_c V_{h,sp.}(z) = W_{abc,sp.}(z) V_{h,sp.}^a(z); \quad (4.1.13)$$

$$\partial_a V_{h,sp.}^b(z) = \delta_a^b V_{h,sp.}^0,$$

where $W_{abc,sp.}(z) \equiv \partial_a \partial_b \partial_c \mathcal{F}(z)$ is the holomorphic part of $C_{\alpha\beta\gamma}$ in special coordinates and for $X^0 = 1$ (see the second row of the first of relations (2.1.9)) and

$$V_{h,sp.}^a(z) \equiv (0, 0, \delta_d^a, z^a), \quad (4.1.14)$$

$$V_{h,sp.}^0(z) \equiv (0, 0, 0, 1). \quad (4.1.15)$$

By adding the definition $V_{h,sp.,a}(z) \equiv \partial_a V_{h,sp.}(z)$ and the trivial result $\partial_a V_{h,sp.}^0(z) = 0$ to Eqs. (4.1.13), one finally gets the set of differential relations [59]

$$\begin{aligned}\partial_a V_{h,sp.}(z) &= V_{h,sp.,a}(z), \\ \partial_a \partial_b V_{h,sp.}(z) &= \partial_a V_{h,sp.,b}(z) = W_{abc,sp.}(z) V_{h,sp.}^c(z), \\ \partial_a V_{h,sp.}^b(z) &= \delta_a^b V_{h,sp.}^0, \\ \partial_a V_{h,sp.}^0(z) &= 0,\end{aligned}\tag{4.1.16}$$

which are the holomorphic counterparts of SK relations (4.1.2), written in special coordinates and for $X^0 = 1$.

By “*holomorphically covariantizing*” the relations (4.1.16), *i.e.* by writing them in a generic system of homogeneous coordinates, one obtains (notice that here a -indices and α -indices in general do not coincide) [59]

$$\begin{aligned}\widehat{D}_\alpha V_h(z) &= V_{h,\alpha}(z), \\ \widehat{D}_\alpha \widehat{D}_\beta V_h(z) &= \widehat{D}_\alpha V_{h,\beta}(z) = W_{\alpha\beta\gamma}(z) V_h^\gamma(z), \\ \widehat{D}_\alpha V_h^\beta(z) &= \delta_\alpha^\beta V_h^0(z), \\ \widehat{D}_\alpha V_h^0(z) &= 0,\end{aligned}\tag{4.1.17}$$

where $V_h(z)$, $V_{h,\beta}(z)$, $V_h^\beta(z)$ and $V_h^0(z)$ are respectively given by the following formulæ¹³ [59]:

$$\begin{aligned}V_h(z) &= (X^0(z), X^a(z), X^0(z) e_a^\alpha(z) \partial_\alpha \mathcal{F}(z), X^a(z) e_a^\alpha(z) \partial_\alpha \mathcal{F}(z) - 2X^0(z) \mathcal{F}(z)); \\ V_{h,\beta}(z) &= \widehat{D}_\beta V_h(z) = (0, X^0(z) e_\beta^a(z), X^0(z) e_a^\alpha(z) \widehat{D}_\alpha \partial_\beta \mathcal{F}(z), -X^0(z) \partial_\beta \mathcal{F}(z) + X^a(z) e_a^\alpha(z) \widehat{D}_\alpha \partial_\beta \mathcal{F}(z)); \\ V_h^\beta(z) &= (0, 0, (X^0(z))^{-1} e_a^\beta(z), (X^0(z))^{-2} X^a(z) e_a^\beta(z)); \\ V_h^0(z) &= (0, 0, 0, (X^0(z))^{-1}),\end{aligned}\tag{4.1.18}$$

which correspond to the “*holomorphically covariantized*” counterparts of Eqs. (4.1.8), (4.1.11), (4.1.14) and (4.1.15), respectively. $e_a^\alpha(z) \equiv e_a^\alpha(t(z))$ is the inverse of $e_a^\alpha(z)$ (defined by Eq. (4.1.6)):

$$e_a^\alpha(t(z)) \frac{\partial t^a(z)}{\partial z^\beta} \equiv \delta_\beta^\alpha, \quad e_a^\alpha(t(z)) \frac{\partial t^b(z)}{\partial z^\alpha} \equiv \delta_a^b \iff e_a^\alpha(t(z)) \equiv \left. \frac{\partial z^\alpha(t)}{\partial t^a} \right|_{t=t(z)}.\tag{4.1.19}$$

$W_{\alpha\beta\gamma}(z)$ (completely symmetric, with Kähler weights $(4, 0)$, and defined in the second row of the first of relations

¹³The first of relations (4.1.18) corresponds to Eq. (4.1.3) by using the definition of $t^a(z)$ s and the homogeneity of degree 2 of the prepotential F .

(2.1.9)) is the covariant part of $C_{\alpha\beta\gamma}(z, \bar{z})$ in generic homogeneous coordinates. It can be further elaborated as follows:

$$W_{\alpha\beta\gamma}(z) \equiv (\partial_\alpha X^\Lambda(z)) (\partial_\beta X^\Sigma(z)) (\partial_\gamma X^\Xi(z)) \partial_\Xi \partial_\Sigma \partial_\Lambda F(X)|_{X=X(z)} = \quad (4.1.20)$$

$$= (X^0(z))^2 e_\alpha^a(z) e_\beta^b(z) e_\gamma^c(z) \partial_c \partial_b \partial_a \mathcal{F}(t)|_{t=t(z)} = \quad (4.1.21)$$

$$= (X^0(z))^2 \left[\begin{aligned} & \partial_\gamma \partial_\beta \partial_\alpha \mathcal{F}(z) + \\ & - \left(\partial_\gamma e_\beta^b(z) \right) e_b^{\beta'}(t(z)) \partial_{\beta'} \partial_\alpha \mathcal{F}(z) - (\partial_\gamma e_\alpha^a(z)) e_a^{\alpha'}(t(z)) \partial_\beta \partial_{\alpha'} \mathcal{F}(z) + \\ & - (\partial_\beta e_\alpha^a(z)) e_a^{\alpha'}(t(z)) \partial_\gamma \partial_{\alpha'} \mathcal{F}(z) + e_\alpha^a(z) \left(\partial_\gamma \partial_\beta e_a^{\alpha'}(t(z)) \right) \partial_{\alpha'} \mathcal{F}(z) + \\ & + (\partial_{\beta'} e_\alpha^a(z)) \left(\partial_\gamma e_\beta^b(z) \right) e_b^{\beta'}(t(z)) e_a^{\alpha'}(t(z)) \partial_{\alpha'} \mathcal{F}(z) \end{aligned} \right]. \quad (4.1.22)$$

Notice that a new holomorphic covariant derivative $\widehat{D}_\alpha$ has been introduced. In analogy with the usual covariant derivative in Kähler-Hodge manifold, the action of $\widehat{D}_\alpha$ on a vector ϕ_β with Kähler weight p reads [58, 59]

$$\widehat{D}_\alpha \phi_\beta(z, \bar{z}) = \left(\partial_\alpha + \frac{p}{2} \widehat{K}_\alpha(z) \right) \phi_\beta(z, \bar{z}) + \widehat{\Gamma}_{\alpha\beta}{}^\gamma(z) \phi_\gamma(z, \bar{z}), \quad (4.1.23)$$

where $\widehat{\Gamma}_{\alpha\beta}{}^\gamma(z)$ is the holomorphic part of the Christoffel connection $\Gamma_{\alpha\beta}{}^\gamma(z, \bar{z})$ of the SK manifold being considered [58, 59] ($e_\alpha^a(z) e_a^\gamma(z) = \delta_\alpha^\gamma$, $e_\alpha^a(z) e_b^\alpha(z) = \delta_b^a$; also recall definition (2.1.11)):

$$\widehat{\Gamma}_{\alpha\beta}{}^\gamma(z) \equiv -(\partial_\beta e_\alpha^a(z)) e_a^\gamma(z) = \Gamma_{\alpha\beta}{}^\gamma(z, \bar{z}) - T_{\alpha\beta}{}^\gamma(z, \bar{z}) \equiv \quad (4.1.24)$$

$$\equiv -g^{\gamma\bar{\gamma}}(z, \bar{z}) \partial_\alpha \partial_\beta \bar{\partial}_{\bar{\gamma}} K(z, \bar{z}) + e_\alpha^a(z) e_\beta^b(z) \left[\frac{\partial^3 K(t(z), \bar{t}(\bar{z}))}{\partial t^b \partial t^a \partial \bar{t}^{\bar{d}}} \right] g^{c\bar{d}}(z, \bar{z}) e_c^\gamma(z). \quad (4.1.25)$$

It can be checked that $\widehat{\Gamma}_{\alpha\beta}{}^\gamma(z)$ transforms as a connection under holomorphic reparametrizations. $\widehat{\Gamma}_{\alpha\beta}{}^\gamma(z)$ is usually called the *Christoffel holomorphic connection*. By its very definition (4.1.24), it follows that $e_\alpha^a(z)$ can be seen as the *Christoffel holomorphic n_V -bein*.

It is worth pointing out that the $\widehat{\Gamma}_{\alpha\beta}{}^\gamma$ s are the Christoffel symbols of the second kind of an holomorphic Riemann metric

$$\widehat{g}_{\alpha\beta}(z) \equiv e_\alpha^a(z) e_\beta^b(z) \eta_{ab} = \frac{\partial \left[\frac{X^a(z)}{X^0(z)} \right]}{\partial z^\alpha} \frac{\partial \left[\frac{X^b(z)}{X^0(z)} \right]}{\partial z^\beta} \eta_{ab}, \quad (4.1.26)$$

where η_{ab} is constant (invertible) symmetric matrix (note that $\widehat{g}_{\alpha\beta}(z)$ has two holomorphic indices, in contrast to the Kähler metric $g_{\alpha\bar{\beta}}(z, \bar{z}) = \partial_\alpha \bar{\partial}_{\bar{\beta}} K(z, \bar{z})$). $\widehat{g}_{\alpha\beta}(z)$ is the metric tensor of the so-called *holomorphic geometry* “embedded” in the considered SK geometry. Due to Eq. (4.1.6), it can be checked that $\widehat{\Gamma}_{\alpha\beta}{}^\gamma(z)$ is actually a Riemann-flat connection, since it holds that

$$\widehat{R}_{\delta\alpha\beta}{}^\gamma(z) \equiv \partial_\delta \widehat{\Gamma}_{\alpha\beta}{}^\gamma(z) - \partial_\alpha \widehat{\Gamma}_{\delta\beta}{}^\gamma(z) - \widehat{\Gamma}_{\alpha\beta}{}^\zeta(z) \widehat{\Gamma}_{\zeta\delta}{}^\gamma(z) + \widehat{\Gamma}_{\delta\beta}{}^\zeta(z) \widehat{\Gamma}_{\zeta\alpha}{}^\gamma(z) = 0. \quad (4.1.27)$$

Moreover, since $X^0(z)$ has Kähler weights $(2, 0)$, the quantity

$$\hat{K}_\alpha(z) \equiv -\partial_\alpha [\ln(X^0(z))] \quad (4.1.28)$$

transforms as a connection under Kähler gauge transformations:

$$K(z, \bar{z}) \longrightarrow K(z, \bar{z}) + f(z) + \bar{f}(\bar{z}) \implies \hat{K}_\alpha(z) \longrightarrow \hat{K}_\alpha(z) + \partial_\alpha f(z). \quad (4.1.29)$$

$\hat{K}_\alpha(z)$ is usually called the *Kähler holomorphic connection*. By its very definition (4.1.28), it thus follows that $X^0(z)$ can be seen as the *Kähler holomorphic 1-bein*.

In the formalism of *holomorphic geometry* introduced above, it can be easily checked that the quantity $e_a^\alpha(z) \hat{D}_\alpha \partial_\beta \mathcal{F}(z)$, appearing in the vector $V_{h,\beta}(z)$ introduced in the second of relations (4.1.18), can be further elaborated as follows:

$$e_a^\alpha(z) \hat{D}_\alpha \partial_\beta \mathcal{F}(z) = e_a^\alpha(z) \hat{D}_\alpha \hat{D}_\beta \mathcal{F}(z) = \partial_\beta [e_a^\alpha(z) \partial_\alpha \mathcal{F}(z)] = e_a^\alpha(z) \hat{D}_\beta \partial_\alpha \mathcal{F}(z) = e_a^\alpha(z) \hat{D}_\beta \hat{D}_\alpha \mathcal{F}(z). \quad (4.1.30)$$

Finally, it should be observed that special coordinates are *flat* coordinates for the holomorphic geometry, because for special coordinates ($e_a^\alpha(z) = \delta_\alpha^a$) (in the Kähler gauge $X^0 = 1$) Eqs. (4.1.24), (4.1.28) and (4.1.26) respectively reduce to

$$\hat{\Gamma}_{\alpha\beta}{}^\gamma(z) = 0; \quad (4.1.31)$$

$$\hat{K}_\alpha(z) = 0; \quad (4.1.32)$$

$$\hat{g}_{\alpha\beta}(z) = \eta_{\alpha\beta}. \quad (4.1.33)$$

It is worth pointing out that the system (4.1.17) is the holomorphic counterpart of the system (4.1.2), and it is manifestly covariant with respect to the holomorphic geometry defined by $\hat{g}_{\alpha\beta}(z)$ and $\hat{K}_\alpha(z)$. By breaking the “*holomorphic covariance*” and choosing special coordinates (and fixing Kähler gauge such that $X^0 = 1$), the system (4.1.17) reduces to the system (4.1.16). The system of holomorphic differential relations (4.1.17) is usually referred to as the (holomorphic) *Picard-Fuchs (PF) system*.

4.2 1-Modulus

Let us now specialize the treatment to the 1-modulus case. Once again, such a case is peculiarly simple, since for $n_V = 1$ the Christoffel holomorphic n_V -bein is nothing but an holomorphic function ($z^1 \equiv \psi$, $t^1 \equiv t$, $\partial_\psi \equiv \partial$):

$$e_\psi^1(\psi) \equiv e(\psi) \equiv \frac{\partial \left[\frac{X^1(\psi)}{X^0(\psi)} \right]}{\partial \psi} \equiv \frac{\partial t(\psi)}{\partial \psi} = \frac{X^0(\psi) \partial X^1(\psi) - X^1(\psi) \partial X^0(\psi)}{[X^0(\psi)]^2}, \quad (4.2.1)$$

and the connections and metric of holomorphic geometry reduce to

$$\begin{aligned} \hat{\Gamma}_{\psi\psi}{}^\psi(\psi) &\equiv \hat{\Gamma}(\psi) = -\partial [\ln(e(\psi))]; \\ \hat{K}_\psi(\psi) &\equiv \hat{K}(\psi) = -\partial [\ln(X^0(\psi))]; \\ \hat{g}_{\psi\psi}(\psi) &\equiv \hat{g}(\psi) = [e(\psi)]^2 \eta, \quad \eta \in \mathbb{C}_0. \end{aligned} \quad (4.2.2)$$

Consequently, the action of $\widehat{D}_\psi \equiv \widehat{D}$ on a 1-vector (function) $\phi(\psi, \bar{\psi})$ with Kähler weights $(p, \bar{p})$ reads

$$\begin{aligned}\widehat{D}\phi(\psi, \bar{\psi}) &= \left(\partial + \frac{p}{2}\widehat{K}(\psi) + \widehat{\Gamma}(\psi)\right)\phi(\psi, \bar{\psi}) = \\ &= \left\{\partial - \frac{p}{2}\partial[\ln(X^0(\psi))] - \partial[\ln(e(\psi))]\right\}\phi(\psi, \bar{\psi}).\end{aligned}\quad (4.2.3)$$

It can be shown [58, 59] that in the 1-modulus case the PF system (4.1.17) is equivalent to the following complex differential relation:

$$\widehat{D}^2 \left\{ [\mathcal{W}(\psi)]^{-1} \widehat{D}^2 V_h(\psi) \right\} = 0, \quad (4.2.4)$$

where (see Eqs. (4.1.20)-(4.1.22), and the second row of the first of relations (2.1.9))

$$\begin{aligned}\mathcal{W}(\psi) &\equiv W_{\psi\psi\psi}(\psi) = e^{-K(\psi, \bar{\psi})} C(\psi, \bar{\psi}) = \\ &= [\partial X^0(\psi)]^3 \frac{\partial^3 F(X)}{(\partial X^0)^3} \Big|_{X=X(\psi)} + 3[\partial X^0(\psi)]^2 [\partial X^1(\psi)] \frac{\partial^3 F(X)}{(\partial X^0)^2 \partial X^1} \Big|_{X=X(\psi)} + \\ &+ 3[\partial X^1(\psi)]^2 [\partial X^0(\psi)] \frac{\partial^3 F(X)}{(\partial X^1)^2 \partial X^0} \Big|_{X=X(\psi)} + [\partial X^1(\psi)]^3 \frac{\partial^3 F(X)}{(\partial X^1)^3} \Big|_{X=X(\psi)} = \\ &= (X^0(\psi))^2 [e(\psi)]^3 \frac{\partial^3 \mathcal{F}(t)}{(\partial t)^3} \Big|_{t=t(\psi)} = \\ &= (X^0(\psi)) \left[\begin{aligned} &3[e(\psi)]^{-2} [\partial e(\psi)]^2 \partial \mathcal{F}(\psi) - [e(\psi)]^{-1} [\partial^2 e(\psi)] \partial \mathcal{F}(\psi) + \\ &-3[e(\psi)]^{-1} [\partial e(\psi)] \partial^2 \mathcal{F}(\psi) + \partial^3 \mathcal{F}(\psi) \end{aligned} \right],\end{aligned}\quad (4.2.5)$$

and (see Eq. (4.1.3) and the first of relations (4.1.18))

$$\begin{aligned}V_h(\psi) &= (X^0(\psi), X^1(\psi), F_1(X(\psi)), -F_0(X(\psi))) = \\ &= (X^0(\psi), X^1(\psi), X^0(\psi)[e(\psi)]^{-1} \partial \mathcal{F}(\psi), X^1(\psi)[e(\psi)]^{-1} \partial \mathcal{F}(\psi) - 2X^0(\psi) \mathcal{F}(\psi)).\end{aligned}\quad (4.2.6)$$

Eq. (4.2.4) can also be rewritten as a fourth order linear ordinary differential equation for $V_h(\psi)$ (*1-modulus PF Eq.*) [58, 59]:

$$\sum_{n=0}^4 a_n(\psi) \partial^n V_h(\psi) = 0, \quad (4.2.7)$$

where $\partial^n \equiv \frac{\partial^n}{(\partial \psi)^n}$ ($n=0$ corresponds to the identity operator)¹⁴. The functions $a_n(\psi)$ can be obtained by comparing

¹⁴For a general treatment of the n_V -moduli case, see [59].

Eqs. (4.2.4) and (4.2.7) [58] (notation of dependence on ψ omitted throughout):

$$\begin{aligned}
a_4 &\equiv \mathcal{W}^{-1}; \\
a_3 &\equiv 2\partial(\mathcal{W}^{-1}); \\
a_2 &\equiv \mathcal{W}^{-1}(\partial\hat{\Lambda} - \hat{\Lambda}^2 + 2\hat{\Sigma}) + [\partial(\mathcal{W}^{-1})]\hat{\Lambda} + \partial^2(\mathcal{W}^{-1}); \\
a_1 &\equiv \mathcal{W}^{-1}(\partial^2\hat{\Lambda} + 2\partial\hat{\Sigma} - 2\hat{\Lambda}\partial\hat{\Lambda}) + [\partial(\mathcal{W}^{-1})](2\hat{\Sigma} + 2\partial\hat{\Lambda} - \hat{\Lambda}^2) + [\partial^2(\mathcal{W}^{-1})]\hat{\Lambda}; \\
a_0 &\equiv \mathcal{W}^{-1}(\hat{\Sigma}^2 - \hat{\Sigma}\partial\hat{\Lambda} - \hat{\Lambda}\partial\hat{\Sigma} + \partial^2\hat{\Sigma}) + [\partial(\mathcal{W}^{-1})](2\partial\hat{\Sigma} - \hat{\Lambda}\hat{\Sigma}) + [\partial^2(\mathcal{W}^{-1})]\hat{\Sigma},
\end{aligned} \tag{4.2.8}$$

where the following holomorphic functions have been introduced (recall the first two Eqs. of (4.2.2)):

$$\begin{aligned}
\hat{\Lambda} &\equiv 2\hat{K} + \hat{\Gamma} = -2\partial[\ln(X^0)] - \partial[\ln(e)] = -\partial\ln(X^0\partial X^1 - X^1\partial X^0) = -\frac{X^0\partial^2 X^1 - X^1\partial^2 X^0}{X^0\partial X^1 - X^1\partial X^0}; \\
\hat{\Sigma} &\equiv \partial\hat{K} + \hat{K}^2 + \hat{\Gamma}\hat{K} = -\partial^2[\ln(X^0)] + [\partial[\ln(X^0)]]^2 + [\partial[\ln(X^0)]]\partial[\ln(e)] = \\
&= \frac{(\partial X^0)\partial^2 X^1 - (\partial X^1)\partial^2 X^0}{X^0\partial X^1 - X^1\partial X^0} = \hat{\Lambda} + \frac{(X^0 + \partial X^0)\partial^2 X^1 - (X^1 + \partial X^1)\partial^2 X^0}{X^0\partial X^1 - X^1\partial X^0}.
\end{aligned} \tag{4.2.9}$$

The definitions (4.2.8) entail the following differential relations between the functional coefficients of 1-modulus PF Eq. (4.2.7):

$$\begin{aligned}
a_3 &= 2\partial a_4; \\
a_1 &= \partial(a_2 - \tfrac{1}{2}\partial a_3).
\end{aligned} \tag{4.2.10}$$

Thus, the holomorphic functions a_n only depend on the (inverse of the) holomorphic Yukawa coupling function $\mathcal{W}$, on the holomorphic connections $\hat{K}$ and $\hat{\Gamma}$ (or, equivalently, on the holomorphic 1-beins X^0 and e). If $\mathcal{W}$, $\hat{K}$ and $\hat{\Gamma}$ (or $\mathcal{W}$, X^0 and e) are given as input, Eq. (4.2.7) is a fourth order linear ordinary differential equation for $V_h(\psi)$. Also, for given a_n the definitions (4.2.8) turn into non-linear ordinary differential equations for X^0 and e .

In special coordinates (with Kähler gauge fixed such that $X^0 = 1$) one has $X^1(\psi) = t(\psi) = \psi$ and $e(\psi) = 1$, and the $a_n(\psi)$ s simplify drastically¹⁵:

$$\begin{aligned}
a_{4,sp.} &\equiv \mathcal{W}_{sp.}^{-1}; \\
a_{3,sp.} &\equiv 2\partial(\mathcal{W}_{sp.}^{-1}) = -2\mathcal{W}_{sp.}^{-1}\frac{\partial\mathcal{W}_{sp.}}{\mathcal{W}_{sp.}}; \\
a_{2,sp.} &\equiv \partial^2(\mathcal{W}_{sp.}^{-1}) = \mathcal{W}_{sp.}^{-1}\left[2\left(\frac{\partial\mathcal{W}_{sp.}}{\mathcal{W}_{sp.}}\right)^2 - \frac{\partial^2\mathcal{W}_{sp.}}{\mathcal{W}_{sp.}}\right]; \\
a_{1,sp.} &= 0 = a_{0,sp.},
\end{aligned} \tag{4.2.12}$$

¹⁵However, it is here worth pointing out that in the following general treatment of (mirror) Fermat CY_3 s $t(\psi)$ is *not* a special coordinate, *i.e.* $t(\psi) \neq \psi$, and moreover the Kähler gauge is left unfixed (in particular, $X^0(\psi) \neq 1$).

and thus Eq. (4.2.7) can be rewritten as

$$\sum_{n=0}^4 a_{n,sp.}(\psi) \partial^n V_{h,sp.}(\psi) = \left\{ \partial^4 - 2(\partial \ln(\mathcal{W}_{sp.})) \partial^3 + \left[2(\partial \ln(\mathcal{W}_{sp.}))^2 - \frac{\partial^2 \mathcal{W}_{sp.}}{\mathcal{W}_{sp.}} \right] \partial^2 \right\} V_{h,sp.} = 0, \quad (4.2.13)$$

where, by recalling Eq. (4.2.6), $V_{h,sp.}$ reads

$$V_{h,sp.}(\psi) = (1, \psi, \partial \mathcal{F}(\psi), \psi \partial \mathcal{F}(\psi) - 2\mathcal{F}(\psi)), \quad (4.2.14)$$

which is nothing but the case $n_V = 1$ of Eq. (4.1.8). Moreover, since in special coordinates (with $X^0 = 1$) the holomorphic Yukawa coupling function $\mathcal{W}$ defined by Eq. (4.2.5) reads

$$\mathcal{W}_{sp.} = \partial^3 \mathcal{F}, \quad (4.2.15)$$

Eq. (4.2.13) can be also rewritten as

$$\left\{ \partial^4 - 2 \frac{\partial^4 \mathcal{F}}{\partial^3 \mathcal{F}} \partial^3 + \left[2 \left(\frac{\partial^4 \mathcal{F}}{\partial^3 \mathcal{F}} \right)^2 - \frac{\partial^5 \mathcal{F}}{\partial^3 \mathcal{F}} \right] \partial^2 \right\} V_{h,sp.} = 0. \quad (4.2.16)$$

The 1-modulus PF Eq. (4.2.7) can be further elaborated, by recalling some basic facts about fourth order linear ordinary differential equations [41]. It is interesting to notice that not all the $a_n(\psi)$ s are actually relevant. Firstly, one can scale a_4 out from the 1-modulus PF Eq. (4.2.7), and furthermore drop the coefficient of the third derivative by performing the following rescaling redefinition of $V_h(\psi)$ [59]:

$$V_h(\psi) \longrightarrow V_h(\psi) \exp \left[-\frac{1}{4} \int^\psi d\psi' \frac{a_3(\psi')}{a_4(\psi')} \right] = V_h(\psi) \exp \left[-\frac{1}{2} \int^\psi d\psi' \partial \ln(a_4(\psi')) \right] = [a_4(\psi)]^{-1/2} V_h(\psi), \quad (4.2.17)$$

where the first of differential relations (4.2.11) was used. By doing this, the PF Eq. (4.2.7) can be recast in the following form:

$$\sum_{n=0}^4 c_n(\psi) \partial^n V_h(\psi) = 0, \quad (4.2.18)$$

where

$$\begin{aligned} c_4 &= 1; \\ c_3 &= 0; \\ c_2 &= a_4^{-1} \left(\frac{3}{8} a_4^{-1} a_3^2 - \frac{3}{2} \partial a_3 + a_2 \right); \\ c_1 &= -\frac{1}{2} a_4^{-1} \left(\frac{3}{4} a_4^{-2} a_3^3 - 3 a_4^{-1} a_3 \partial a_3 + 3 \partial^2 a_3 + a_4^{-1} a_2 a_3 - 2 \partial a_2 \right); \\ c_0 &= -\frac{1}{16} a_4^{-1} \left[\begin{aligned} & -\frac{45}{16} a_4^{-3} a_3^4 + \frac{27}{2} a_4^{-2} a_3^2 \partial a_3 - 9 a_4^{-1} (\partial a_3)^2 + \\ & -10 a_4^{-1} a_3 \partial^2 a_3 + 4 \partial^3 a_3 - 3 a_4^{-2} a_3^2 a_2 + \\ & + 4 a_4^{-1} a_2 \partial a_3 + 4 a_4^{-1} a_3 \partial a_2 - 16 a_0 \end{aligned} \right]. \end{aligned} \quad (4.2.19)$$

Furthermore, it is possible to remove the dependence of Eq. (4.2.7) on the holomorphic Kähler connection by introducing the Kähler invariant holomorphic vector [58]

$$\tilde{V}_h(\psi) \equiv [X^0(\psi)]^{-1} V_h(\psi). \quad (4.2.20)$$

By doing this, the PF Eq. (4.2.7) can be recast in the following form:

$$\sum_{n=0}^4 d_n(\psi) \partial^n \tilde{V}_h(\psi) = 0, \quad (4.2.21)$$

where

$$\begin{aligned} d_4 &= 1; \\ d_3 &= 4 \frac{\partial X^0}{X^0} + \frac{a_3}{a_4} = -2\Theta; \\ d_2 &= 6 \frac{\partial^2 X^0}{X^0} + 3 \frac{a_3}{a_4} \frac{\partial X^0}{X^0} + \frac{a_2}{a_4} = \Theta^2 - \partial\Theta - \hat{\Gamma}\Theta - \hat{\Gamma}^2 + \partial\hat{\Gamma} = \frac{1}{4}d_3^2 + \frac{1}{2}\partial d_3 + \frac{1}{2}\hat{\Gamma}d_3 - \hat{\Gamma}^2 + \partial\hat{\Gamma}; \\ d_1 &= 4 \frac{\partial^3 X^0}{X^0} + 3 \frac{a_3}{a_4} \frac{\partial^2 X^0}{X^0} + 2 \frac{a_2}{a_4} \frac{\partial X^0}{X^0} + \frac{a_1}{a_4} = (\Theta^2 - \partial\Theta) \hat{\Gamma} + \hat{\Gamma}^2 \Theta - 2(\partial\hat{\Gamma}) \Theta + \partial^2 \hat{\Gamma} - 2\hat{\Gamma}\partial\hat{\Gamma} = \\ &= \frac{1}{4}\hat{\Gamma}d_3^2 - \left(\frac{1}{2}\hat{\Gamma}^2 - \partial\hat{\Gamma}\right) d_3 + \frac{1}{2}\hat{\Gamma}\partial d_3 + \partial^2 \hat{\Gamma} - 2\hat{\Gamma}\partial\hat{\Gamma} = \hat{\Gamma}d_2 - \left(\hat{\Gamma}^2 - \partial\hat{\Gamma}\right) d_3 + \partial^2 \hat{\Gamma} - 3\hat{\Gamma}\partial\hat{\Gamma} + \hat{\Gamma}^3; \\ d_0 &= \frac{1}{X^0} \left(\partial^4 + \frac{a_3}{a_4} \partial^3 + \frac{a_2}{a_4} \partial^2 + \frac{a_1}{a_4} \partial + \frac{a_0}{a_4} \right) X^0 = 0, \end{aligned} \quad (4.2.22)$$

where the Kähler gauge-invariant holomorphic function

$$\Theta \equiv \partial \ln \left[\frac{\mathcal{W}}{(X^0)^2} \right] = \frac{\partial \mathcal{W}}{\mathcal{W}} - 2 \frac{\partial X^0}{X^0} = -\frac{1}{2} \frac{a_3}{a_4} + 2\hat{K} \quad (4.2.23)$$

has been introduced. Eqs. (4.2.22) imply the d_n to be formally independent on $\hat{K}$, and all their dependence on X^0 appears through the function Θ . Since X^0 is the first component of the holomorphic period vector V_h given by Eq. (4.2.6), the vanishing of d_0 directly follows from the PF Eq. (4.2.7) (with $a_4 \neq 0$ and $X^0 \neq 0$).

5 Holomorphic Geometry and Flat Holomorphic Matrix Connection

5.1 n_V -Moduli

As shown in [60], the basic, defining differential relations (4.1.2) of SK geometry can be recast as a vanishing condition for a suitably defined flat symplectic non-holomorphic connection. Analogously, the holomorphic differential Eqs. (4.1.17) can be rewritten as a vanishing condition for a suitably defined flat holomorphic connection, *i.e.* as [59]

$$(\mathbb{I}_{2n_V+2} \partial_\alpha - \mathbf{A}_\alpha(z)) \mathbf{V}_h(z) = 0, \quad (5.1.1)$$

where $\mathbf{V}_h(z)$ is a $(2n_V + 2) \times (2n_V + 2)$ holomorphic matrix ($(2n_V + 2) \times 1$ vector with $1 \times (2n_V + 2)$ vector entries) defined as follows [59]:

$$\mathbf{V}_h(z) \equiv \begin{pmatrix} V_h(z) \\ V_{h,\beta}(z) \\ V_h^\beta(z) \\ V_h^0(z) \end{pmatrix}, \quad (5.1.2)$$

where the entries are defined in Eqs. (4.1.18). On the other hand, $\mathbf{A}_\alpha(z)$ is the following $(2n_V + 2) \times (2n_V + 2)$ holomorphic connection matrix:

$$\mathbf{A}_\alpha(z) \equiv \begin{pmatrix} -\widehat{K}_\alpha(z) & \delta_\alpha^\gamma & 0 & 0 \\ 0 & -\left(\widehat{\Gamma}_\alpha(z) + \widehat{K}_\alpha(z) \mathbb{I}_{n_V}\right)_\beta^\gamma & (W_\alpha)_{\gamma\beta}(z) & 0 \\ 0 & 0 & \left(\widehat{\Gamma}_\alpha(z) + \widehat{K}_\alpha(z) \mathbb{I}_{n_V}\right)_\gamma^\beta & \delta_\alpha^\beta \\ 0 & 0 & 0 & \widehat{K}_\alpha(z) \end{pmatrix}. \quad (5.1.3)$$

It should be noticed that $\mathbf{A}_\alpha(z)$ is Lie-algebra valued in $\mathfrak{sp}(2n_V + 2)$, *i.e.* it satisfies the infinitesimal symplecticity condition [59]

$$\mathbf{A}_\alpha^T(z) Q + Q \mathbf{A}_\alpha(z) = 0, \quad (5.1.4)$$

where Q is the symplectic metric defined in Eq. (4.1.4).

Put another way, it can be stated that the PF Eqs. (4.1.17) are equivalent to the matrix system (5.1.1), with $\mathbf{A}_\alpha(z)$ defined by Eq. (5.1.3). The general solution of such an holomorphic matrix system is given by Eqs. (4.1.18) arranged as a vector as given by Eq. (5.1.2).

As expected, by specializing the holomorphic matrix system (5.1.1) in special coordinates and choosing the Kähler gauge to be such that $X^0 = 1$, one gets the following holomorphic matrix system:

$$(\mathbb{I}_{2n_V+2} \partial_a - \mathbf{A}_{a,sp.}(z)) \mathbf{V}_{h,sp.}(z) = 0, \quad (5.1.5)$$

which is equivalent to the holomorphic system (4.1.16).

$\mathbf{V}_{h,sp.}(z)$ is a $(2n_V + 2) \times (2n_V + 2)$ holomorphic matrix ($(2n_V + 2) \times 1$ vector with $1 \times (2n_V + 2)$ vector entries) defined as follows [59]:

$$\mathbf{V}_{h,sp.}(z) \equiv \begin{pmatrix} V_{h,sp.}(z) \\ V_{h,sp.,b}(z) \\ V_{h,sp.}^b(z) \\ V_{h,sp.}^0(z) \end{pmatrix}, \quad (5.1.6)$$

where the entries are defined in Eqs. (4.1.8), (4.1.11), (4.1.14) and (4.1.15). It is worth mentioning that the matrices $\mathbf{V}_{h,sp.}(z)$ and $\mathbf{V}_h(z)$ have a symplectic structure with respect to the symplectic metric relevant for holomorphic geometry, *i.e.* with respect to Q defined in Eq. (4.1.4):

$$\begin{aligned}\mathbf{V}_{h,sp.}^T(z) Q \mathbf{V}_{h,sp.}(z) &= Q; \\ \mathbf{V}_h^T(z) Q \mathbf{V}_h(z) &= Q.\end{aligned}\tag{5.1.7}$$

$\mathbf{A}_{a,sp.}(z)$ (named $\mathbb{C}_a$ in Eq. (3.6) of the first Ref. of [59]) is the $(2n_V + 2) \times (2n_V + 2)$ holomorphic connection matrix obtained by $\mathbf{A}_\alpha(z)$ (given by Eq. (5.1.3)) by putting $\hat{\Gamma}_\alpha(z) = 0 = \hat{K}_\alpha(z)$ (also recalling that in special coordinates a -indices and α -indices coincide). Clearly, as its “holomorphically covariant” counterpart $\mathbf{A}_\alpha(z)$, also $\mathbf{A}_{a,sp.}(z)$ is Lie-algebra valued in $\mathfrak{sp}(2n_V + 2)$, and therefore it satisfies a corresponding infinitesimal symplecticity condition.

In other words, it can be stated that the the holomorphic system (4.1.16) can be recast in the matrix form (5.1.5), with $\mathbf{A}_{a,sp.}(z)$ defined by Eq. (3.6) of the first Ref. of [59]. The general solution of such an holomorphic matrix system is given by Eqs. (4.1.8), (4.1.11), (4.1.14) and (4.1.15) arranged as a vector as given by Eq. (5.1.6).

5.2 1-Modulus

Once again, by considering the 1-modulus case more in detail, one obtains a major simplification. The 1-modulus PF Eq. (4.2.7) can be rewritten in matrix form as follows:

$$(\mathbb{I}_4 \partial - \mathbf{A}(\psi)) \mathbf{V}_h(\psi) = 0,\tag{5.2.1}$$

where $\mathbf{V}_h(\psi)$ is a 4×4 holomorphic matrix, obtained by putting $n_V = 1$ in Eq. (5.1.2):

$$\begin{aligned}\mathbf{V}_h(\psi) &\equiv \begin{pmatrix} V_h(\psi) \\ V_{h,\psi}(\psi) \\ V_h^\psi \psi \\ V_h^0(\psi) \end{pmatrix} = \\ &= \begin{pmatrix} X^0 & X^1 & X^0 e^{-1} \partial \mathcal{F} & X^1 e^{-1} \partial \mathcal{F} - 2X^0 \mathcal{F} \\ 0 & X^0 e & X^0 e^{-1} \hat{D} \partial \mathcal{F} & -X^0 \partial \mathcal{F} + X^1 e^{-1} \hat{D} \partial \mathcal{F} \\ 0 & 0 & (X^0)^{-1} e^{-1} & (X^0)^{-2} X^1 e^{-1} \\ 0 & 0 & 0 & (X^0)^{-1} \end{pmatrix},\end{aligned}\tag{5.2.2}$$

where the first row is given by Eq. (4.2.6), and (recall Eq. (4.1.9))

$$\mathcal{F}(\psi) \equiv F\left(\frac{X^1(\psi)}{X^0(\psi)}\right) = (X^0(\psi))^{-2} F(X(\psi))\tag{5.2.4}$$

is the Kähler gauge-invariant holomorphic prepotential. Moreover, the second row of $\mathbf{V}_h$ can be further elaborated by recalling Eq. (4.1.30) and the first of definitions (4.2.2):

$$\widehat{D}\partial\mathcal{F} = \widehat{D}^2\mathcal{F} = e\partial[e^{-1}\partial\mathcal{F}] = [\partial^2 - e^{-1}(\partial e)\partial]\mathcal{F} = [\partial^2 + \widehat{\Gamma}\partial]\mathcal{F}. \quad (5.2.5)$$

On the other hand, $\mathbf{A}(\psi) \equiv \mathbf{A}_\psi(\psi)$ is a 4×4 holomorphic connection matrix, which is Lie-algebra valued in $\mathfrak{sp}(4)$ and corresponds to $n_V = 1$ in Eq. (5.1.3):

$$\begin{aligned} \mathbf{A}(\psi) = & \\ = & \begin{pmatrix} \partial \ln(X^0) & 1 & 0 & 0 \\ 0 & \partial \ln(e) + \partial \ln(X^0) & \mathcal{W} & 0 \\ 0 & 0 & -\partial \ln(e) - \partial \ln(X^0) & 1 \\ 0 & 0 & 0 & -\partial \ln(X^0) \end{pmatrix}. \end{aligned} \quad (5.2.6)$$

where use of Eqs. (4.2.2) has been made (see also Eq. (4.2.5)). Eq. (5.2.6) can be further elaborated; indeed, by using Eqs. (4.2.1), (4.2.2), (4.2.9) and (4.2.10), one easily gets that

$$\begin{aligned} \partial \ln(e) + \partial \ln(X^0) &= -(\widehat{\Gamma} + \widehat{K}) = -\widehat{\Lambda} - \frac{\partial X^0}{X^0} = -\widehat{\Lambda} + \widehat{K} = \\ &= -\widehat{\Sigma} + \frac{[(X^0)^2 + X^0 \partial X^0] \partial^2 X^1 - (X^1 + \partial X^1) X^0 \partial^2 X^0 - X^0 \partial X^0 \partial X^1 - X^1 (\partial X^0)^2}{(X^0)^2 \partial X^1 - X^0 X^1 \partial X^0} = \\ &= \partial \ln\left(\frac{X^0 \partial X^1 - X^1 \partial X^0}{X^0}\right) = \partial \ln(\partial X^1 - X^1 \partial \ln X^0) = \frac{(X^0)^2 \partial^2 X^1 - X^0 X^1 \partial^2 X^0 - X^0 \partial X^0 \partial X^1 - X^1 (\partial X^0)^2}{(X^0)^2 \partial X^1 - X^0 X^1 \partial X^0}. \end{aligned} \quad (5.2.7)$$

It can be stated that the 1-modulus PF Eq. (4.2.7) is equivalent to the matrix system (5.2.1), with $\mathbf{A}(\psi)$ defined by Eq. (5.2.6). The general solution of such an holomorphic matrix system (which corresponds to the most general solution of the fourth order linear PF Eq. (4.2.7)) is given by Eqs. (5.2.2)-(5.2.3) (implemented by Eq. (5.2.5)).

Let us now further specialize our treatment to the 1-modulus SK geometries endowing the moduli space of Fermat CY_3 s. As previously mentioned, the fourth order linear PF ordinary differential equation for each of the four threefolds (classified by the index $k = 5, 6, 8, 10$: see next Section) of such a class of CY_3 s has been obtained for $k = 5$ in [43, 44] (see in particular Eq. (3.9) of [43], where $z \equiv \psi^{-5}$ and $\varpi_0 \sim \frac{1}{\psi} V_h$; see also [45]), and for $k = 6, 8, 10$ in [46] (see Eq. (3.1) of such a Ref., with notation $\alpha \equiv \psi$), where a unified, k -parametrized treatment was exploited (see also [47]).

In order to recast the 1-modulus PF Eqs. given by Eq. (3.1) of [46] in the form (4.2.7) with the differential relations (4.2.11) between the $a_n(\psi)$ s holding, one must multiply them by the function $\psi^{-\xi_k}$, with $\xi_k = 0, 3, 6, 8$ for $k = 5, 6, 8, 10$ respectively. By doing this, one achieves the result that for Fermat CY_3 s the fourth order linear PF Eqs.

(4.2.7) can be recast in the following k -parametrized form¹⁶:

$$\sum_{n=0}^4 a_{n,k}(\psi) \partial^n V_h(\psi) = 0, \quad (5.2.8)$$

$$a_{n,k}(\psi) \equiv -\sigma_n \psi^{n+1} + (-1)^n \tau_{n,k} \psi^{n+1-k},$$

with the constants σ_n s and $\tau_{n,k}$ s given by the following Tables:

n	σ_n
0	1
1	15
2	25
3	10
4	1

Table 1: Values of the integer constants σ_n

$k \longrightarrow$	5	6	8	10
$\tau_{0,k}$	0	0	0	0
$\tau_{1,k}$	0	0	15	35
$\tau_{2,k}$	0	2	15	35
$\tau_{3,k}$	0	2	6	10
$\tau_{4,k}$	1	1	1	1

Table 2: Values of the integer constants $\tau_{n,k}$

6 General Analysis of (Mirror) Fermat Calabi-Yau Threefolds near the Landau-Ginzburg Point

In the present Section we will briefly present the formalism of one-modulus (mirror) Fermat Calabi-Yau threefolds (CY_3 s), focussing our analysis in a suitable neighbourhood of the LG point $\psi = 0$. We will mainly follow [46], and cite where appropriate other relevant works. We also derive original formal results, which will be then used in the case-by-case analysis of extremal BH LG attractors performed in next Sections.

Fermat CY_3 s can be defined as the vanishing *locus* of quasi-homogeneous polynomials in 5 complex variables, of the general form:

$$\mathcal{P}_0 = \sum_{i=0}^4 \nu_i (x^i)^{n_i} = 0; \quad (6.1)$$

such a *locus* gives the embedding of the considered CY_3 in a suitably weighted complex projective space $\mathbb{WCP}_{\nu_0, \nu_1, \nu_2, \nu_3, \nu_4}^4$. By imposing the defining conditions of vanishing first Chern class and of absence of singularities, it is possible to show

¹⁶As we will see in Sect. 12, for self-consistency reasons the 1-modulus PF Eqs. (5.2.8) (with Tables 1 and 2) (which are “corrected” by an overall factor $\psi^{-\xi_k}$ with respect to the ones given in Eq. (3.1) of [46]) need also to be further multiplied by a suitable “normalization” constant (see Eq. (12.1.15)).

that only four possible sets of $\{\nu_i, n_i\}_{i=0,1,2,3,4}$ exist, *all* corresponding to CY_3 s with $h(1,1) = \dim(H^{1,1}(CY_3)) = 1$ (*i.e.* only one Kähler modulus). The four existing Fermat CY_3 s can be classified by introducing the *Fermat parameter* k , defined as the smallest common multiple of the n_i s¹⁷; the only allowed values of k turn out to be $k = 5, 6, 8, 10$. Thus, the four existing Fermat CY_3 s $\mathcal{M}_k$ are given by the following geometrical *loci*¹⁸ [42, 43, 44, 46, 47]:

$$\begin{aligned} k = 5 : \text{quintic } \mathcal{M}_5 &= \left\{ x^i \in (\mathbb{W}) \mathbb{CP}_{1,1,1,1,1}^4 : \mathcal{P}_{0,5} = \sum_{i=0}^4 (x^i)^5 = 0 \right\}; \\ k = 6 : \text{sextic } \mathcal{M}_6 &= \left\{ x^i \in \mathbb{WCP}_{2,1,1,1,1}^4 : \mathcal{P}_{0,6} = 2(x^0)^3 + \sum_{i=1}^4 (x^i)^6 = 0 \right\}; \\ k = 8 : \text{octic } \mathcal{M}_8 &= \left\{ x^i \in \mathbb{WCP}_{4,1,1,1,1}^4 : \mathcal{P}_{0,8} = 4(x^0)^2 + \sum_{i=1}^4 (x^i)^8 = 0 \right\}; \\ k = 10 : \text{dectic } \mathcal{M}_{10} &= \left\{ x^i \in \mathbb{WCP}_{5,2,1,1,1}^4 : \mathcal{P}_{0,10} = 5(x^0)^2 + 2(x^1)^5 + \sum_{i=2}^4 (x^i)^{10} = 0 \right\}. \end{aligned} \quad (6.2)$$

By orbifolding the $\mathcal{M}_k$ s and quotienting by the full phase symmetry group G (see [46] and Refs. therein), one obtains a pair of Fermat CY_3 s¹⁹ $(\mathcal{M}_k, \mathcal{M}'_k)$ related by the so-called *mirror symmetry* [61, 62, 63, 43, 44], with $h(1,1)$ and $h(2,1) = \dim(H^{2,1}(CY_3))$ interchanged (and therefore opposite Euler number $\varkappa$). Correspondingly, the defining vanishing geometrical *loci* will be “deformed” as follows:

$$\mathcal{P}_0 \longrightarrow \mathcal{P} \equiv \mathcal{P}_0 - k\psi \prod_{i=0}^4 x^i. \quad (6.3)$$

All the relevant topological data of Fermat CY_3 s $\mathcal{M}_k$ s are given in Table 3.

ψ is the *Kähler deformation modulus* for Fermat CY_3 s $\mathcal{M}_k$ s (all having $h(1,1) = 1$) and the *complex structure deformation modulus* for the corresponding *mirror* Fermat CY_3 s $\mathcal{M}'_k$ s (all having $h(2,1) = 1$). Since in the treatment and computations performed below we will consider ψ as a complex structure modulus, we will be actually working in the *mirror description* of the considered CY_3 s, *i.e.* we will be considering the *mirror* Fermat CY_3 s $\mathcal{M}'_k$ s ($k = 5, 6, 8, 10$).

In such a framework, the relevant quantities for the $d = 4$ low-energy effective Lagrangian of the $d = 10$ superstring theory compactified on $\mathcal{M}_k$ are given (within the complex structure moduli space ($\dim_{\mathbb{C}} = 1$)) by the Kähler metric and Yukawa couplings on $\mathcal{M}'_k$ (related to $\mathcal{M}_k$ by mirror symmetry). All such quantities will be obtained by the solutions of the fourth order linear PF ordinary differential Eqs. (4.2.7).

¹⁷ k can equivalently be defined as the degree of $\mathcal{W}_0$. It turns out that $k = n_i \nu_i$ (no summation on i) $\forall i = 0, 1, 2, 3, 4$. Moreover, it also holds that $k = \sum_{i=0}^4 \nu_i$.

¹⁸Here and below, we give a name to the Fermat CY_3 s corresponding to the various possible values of the Fermat parameter.

The Fermat CY_3 with $k = 5$ has been named *quintic* some time ago (see *e.g.* [42, 45, 43, 44, 46]).

In a similar fashion, by using the corresponding Latin cardinal adjectives, we name *sextic*, *octic*, and *dectic* the Fermat CY_3 s with $k = 6, 8, 10$, respectively.

¹⁹For simplicity’s sake, we denote in the same way the starting Fermat CY_3 and the one obtained by orbifolding and then quotienting by G .

k $\downarrow$	G	$Ord(G)$	$h(1, 1)$	$h(2, 1)$	$\varkappa \equiv 2[h(1, 1) - h(2, 1)]$
5	$(\mathbb{Z}_5)^3$	5^3	1	101	-200
6	$\mathbb{Z}_3 \otimes (\mathbb{Z}_6)^2$	$3 \cdot 6^2$	1	103	-204
8	$(\mathbb{Z}_8)^2 \otimes \mathbb{Z}_2$	$2 \cdot 8^2$	1	149	-296
10	$(\mathbb{Z}_{10})^2$	$1 \cdot 10^2$	1	145	-288

Table 3: **Basic topological data of Fermat CY_3 s $\mathcal{M}_k$ s**

Near the LG point $\psi = 0$, the 4×1 period vector²⁰ in the PF basis $\varpi_k(\psi)$ is obtained by solving the PF Eqs.²¹

$$\sum_{n=0}^4 a_{n,k}(\psi) \partial^n \varpi_k(\psi) = 0. \quad (6.4)$$

²⁰Once again, in order to make the contact with the relevant literature easier, in this Section as well as in the next ones, we will reconsider column (*i.e.* 4×1), rather than row (*i.e.* 1×4), period vectors. Moreover, the holomorphic period vector in the symplectic basis (hitherto named V_h) will be henceforth denoted by Π .

²¹When comparing Eq. (6.4) to Eq. (4.2.7) (and, more in general, considering the treatment given in Sects. 4 and 5), the 4×1 symplectic holomorphic period vector $V_h \equiv \Pi$ and the 4×1 PF holomorphic period vector ϖ turn out to satisfy the same fourth order linear ordinary differential equation.

Consequently, they necessarily have to be related by a *global* (*i.e.* ψ -independent) “rotation” in the moduli space. This is precisely what happens, with such a “rotation” in the moduli space expressed by the 4×4 real matrices M_k s given in Eqs. (6.16)-(6.17) (see Sect. 4 of [46]).

Here we choose the normalization and the gauge of the holomorphic 3-form defined on $\mathcal{M}'_k$ such that²²

$$\varpi_k(\psi) \equiv -\frac{1}{\psi} \frac{(2\pi i)^3}{\text{Ord}(G_k)} \begin{pmatrix} \omega_{2,k}(\psi) \\ \omega_{1,k}(\psi) \\ \omega_{0,k}(\psi) \\ \omega_{k-1,k}(\psi) \end{pmatrix}, \quad (6.5)$$

with

$$\begin{aligned} \omega_{j,k}(\psi) &\equiv \omega_{0,k}(\beta_k^{2j}\psi), \\ \beta_k &\equiv \exp\left(\frac{\pi i}{k}\right) \end{aligned} \quad (6.6)$$

all being solutions of Eq. (6.4) ($j = 0, 1, \dots, k-1$). However, since Eq. (6.4) is a fourth order (linear) differential equation, only 4 linearly independent solutions $\omega_{j,k}(\psi)$ s exist. Thus, $\forall k = 5, 6, 8, 10$, $k-4$ linear relations between the $\omega_{j,k}(\psi)$ s hold. One possible choice is the following one [43, 44, 46]:

$$k = 5 : \quad \sum_{j=0}^4 \omega_{j,5}(\psi) = 0. \quad (6.7)$$

$$k = 6 : \quad \begin{cases} \omega_{0,6}(\psi) + \omega_{2,6}(\psi) + \omega_{4,6}(\psi) = 0; \\ \omega_{1,6}(\psi) + \omega_{3,6}(\psi) + \omega_{5,6}(\psi) = 0. \end{cases} \quad (6.8)$$

$$k = 8 : \quad \omega_{i,8}(\psi) + \omega_{i+4,8}(\psi) = 0, \quad i = 0, 1, 2, 3. \quad (6.9)$$

$$k = 10 : \quad \begin{cases} \omega_{i,10}(\psi) + \omega_{i+5,10}(\psi) = 0, \quad i = 0, 1, 2, 3, 4; \\ \omega_{0,10}(\psi) + \omega_{2,10}(\psi) + \omega_{3,10}(\psi) + \omega_{4,10}(\psi) + \omega_{5,10}(\psi) = 0. \end{cases} \quad (6.10)$$

The defining Eq. (6.5) expresses the usual conventions, in which one takes $\omega_{0,k}(\psi)$, $\omega_{1,k}(\psi)$, $\omega_{2,k}(\psi)$ and $\omega_{k-1,k}(\psi)$ as basis for $\varpi_k(\psi)$. Therefore, due to relations (6.6), the key quantity turns out to be the holomorphic function $\omega_{0,k}$,

²²The normalization of $\varpi_k(\psi)$ adopted in the present work is the same of [43, 44, 46], and it differs from the one adopted in (a part of the) literature on flux compactifications (see *e.g.* Subsect. 3.2 of [64]) by a factor $\frac{1}{\text{Ord}(G_k)}$ (the reason is that we are interested in the mirror manifolds $\mathcal{M}'_k$ s, not in $\mathcal{M}_k$ s).

On the other hand, it is easy to realize that the gauge of the holomorphic 3-form Ξ adopted in [43, 44, 46] is mostly convenient in order to study the *large complex structure modulus limit* $\psi \rightarrow \infty$. Since we will investigate the *LG limit* $\psi \rightarrow 0$, for our purposes it is better to adopt the gauge of [64], which amounts to redefining the holomorphic 3-form (with respect to the holomorphic 3-form of [43, 44, 46]) as follows:

$$\Xi \longrightarrow \frac{1}{\psi} \Xi.$$

Such a redefinition yields the following Kähler gauge transformation:

$$K(\psi, \bar{\psi}) \longrightarrow K(\psi, \bar{\psi}) + f(\psi) + \bar{f}(\bar{\psi}), \quad f(\psi) \equiv 2\ln(\psi).$$

Such a gauge transformation will affect the Kähler-covariant quantities (*i.e.* the ones with non-vanishing Kähler weights), such as $\exp(K)$, $\mathcal{W}$ and C , but clearly not the Kähler-invariant ones, such as the metric g . In the *LG limit*, for the Kähler-invariant quantities we obtain a matching with the results of [43, 44, 46] (see also [47]); such a matching can be checked to hold also for Kähler-covariant quantities, up to the different overall normalization and Kähler gauge discussed above.

whose series expansion (convergent for $|\psi| < 1$, with the *fundamental region* [43, 44, 46] selected by $0 \leq \arg(\psi) < \frac{2\pi}{k}$) reads [46]

$$\omega_{0,k}(\psi) = - \sum_{m=1}^{\infty} C_{k,m-1} \beta_k^{(k-1)m} \psi^m, \quad (6.11)$$

with

$$C_{k,m-1} \equiv \frac{\Gamma(\frac{m}{k})\Gamma(1-\frac{m}{k})k^{m-1}}{\Gamma(m)\prod_{i=0}^4 \Gamma(1-\frac{m}{k}\nu_{i,k})} \gamma_k^m \in \mathbb{R}, \quad (6.12)$$

$$\gamma_k \equiv \prod_{i=0}^4 (\nu_{i,k})^{-\nu_{i,k}/k} \in \mathbb{R}_0,$$

where in $C_{k,m-1}$ Γ denotes the Euler gamma function

$$\Gamma(s) \equiv \int_0^\infty t^{s-1} e^{-t} dt, \quad \text{Re}(s) > 0. \quad (6.13)$$

By using Eqs. (6.11)-(6.12), the series expansion (convergent for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$) of $\varpi_k(\psi)$ can be written as follows:

$$\varpi_k(\psi) = - \frac{(2\pi i)^3}{\text{Ord}(G_k)} \sum_{m=1}^{\infty} (-1)^m C_{k,m-1} \psi^{m-1} \begin{pmatrix} \beta_k^{3m} \\ \beta_k^m \\ \beta_k^{-m} \\ \beta_k^{-3m} \end{pmatrix}. \quad (6.14)$$

The change between the PF basis and the symplectic basis for holomorphic 4×1 period vector is given by:

$$\Pi_k(\psi) = M_k \varpi_k(\psi). \quad (6.15)$$

where the 4×4 constant matrices M_k read [43, 44, 46]

$$M_5 = \begin{pmatrix} -\frac{3}{5} & -\frac{1}{5} & \frac{21}{5} & \frac{8}{5} \\ 0 & 0 & -1 & 0 \\ -1 & 0 & 8 & 3 \\ 0 & 1 & -1 & 0 \end{pmatrix}, \quad M_6 = \begin{pmatrix} -\frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\ 0 & 0 & -1 & 0 \\ -1 & 0 & 3 & 2 \\ 0 & 1 & -1 & 0 \end{pmatrix}, \quad (6.16)$$

$$M_8 = \begin{pmatrix} -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & -1 & 0 \\ -1 & 0 & 3 & 2 \\ 0 & 1 & -1 & 0 \end{pmatrix}, \quad M_{10} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & -1 & 0 \end{pmatrix}. \quad (6.17)$$

In SKG the holomorphic period vector Π is usually normalized such that (see *e.g.* the report [48]; recall Eq. (2.1.5))

$$K(z, \bar{z}) = -\ln [i\Pi^T(z) \Omega \bar{\Pi}(\bar{z})] = -\ln \left\{ i \left[\bar{X}^\Lambda(\bar{z}) F_\Lambda(z) - X^\Lambda(z) \bar{F}_\Lambda(\bar{z}) \right] \right\}. \quad (6.18)$$

Specializing such a formula for the class of (mirror) CY_3 s being treated, one obtains that the Kähler potential is given

k $\downarrow$	χ_k	λ_k	ς_k
5	5	3	3
6	3	2	0
8	2	2	1
10	1	1	-1

Table 4: **Values of the integer constants χ_k , λ_k and ς_k**

by:

$$\begin{aligned}
K_k(\psi, \bar{\psi}) &= -\ln [i \Pi_k^T(\psi) \Omega \bar{\Pi}_k(\bar{\psi})] = -\ln [-i \Pi_k^\dagger(\bar{\psi}) \Omega \Pi_k(\psi)] = \\
&= -\ln \left\{ i \left[\bar{X}_k^\Lambda(\bar{\psi}) F_{k,\Lambda}(\psi) - X_k^\Lambda(\psi) \bar{F}_{k,\Lambda}(\bar{\psi}) \right] \right\} =
\end{aligned} \tag{6.19}$$

$$\begin{aligned}
&= -\ln [i \Pi_k^\dagger(\bar{\psi}) \Sigma \Pi_k(\psi)] = -\ln [-i \Pi_k^T(\psi) \Sigma \bar{\Pi}_k(\bar{\psi})] = \\
&= -\ln [-i \varpi_k^\dagger(\bar{\psi}) m_k \varpi_k(\psi)] = -\ln [i \varpi_k^T(\psi) m_k \bar{\varpi}_k(\bar{\psi})],
\end{aligned} \tag{6.20}$$

where²³

$$\Sigma \equiv \begin{pmatrix} 0_2 & \mathbb{I}_2 \\ -\mathbb{I}_2 & 0_2 \end{pmatrix}; \tag{6.21}$$

$$\Pi_k(\psi) \equiv \begin{pmatrix} X_k^0(\psi) \\ X_k^1(\psi) \\ F_{k,0}(X(\psi)) \\ F_{k,1}(X(\psi)) \end{pmatrix}; \tag{6.22}$$

$$m_k \equiv M_k^\dagger \Sigma M_k = M_k^T \Sigma M_k = \frac{1}{\chi_k} \begin{pmatrix} 0 & -1 & -\lambda_k & -1 \\ 1 & 0 & -\varsigma_k & -\lambda_k \\ \lambda_k & \varsigma_k & 0 & -1 \\ 1 & \lambda_k & 1 & 0 \end{pmatrix}, \tag{6.23}$$

with the values of χ_k , λ_k and ς_k given in Table 4. By recalling the third column from the left of Table 3, one can observe that $Ord(G_k) = \chi_k k^2$. Substituting Eq. (6.14) and definition (6.23) into Eq. (6.20), one obtains the series

²³Note the change of convention with respect to (the case $n_V = 1$ of) the defining Eq. (2.1.5): $\Sigma = -\Omega|_{n_V=1} = \Omega^T|_{n_V=1}$. On the other hand, Eq. (6.22) is nothing but the (k -indexed) case $n_V = 1$ of the definition (2.1.6).

expansion (converging for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$) of the Kähler potential:

$$K_k(\psi, \bar{\psi}) = -\ln \left(\frac{(2\pi)^6}{(\text{Ord}(G_k))^2} \sum_{m,n=1}^{\infty} C_{k,m-1} C_{k,n-1} \psi^{m-1} \bar{\psi}^{n-1} F_{k,mn} \right), \quad (6.24)$$

where the following infinite rank-2 tensor has been introduced²⁴:

$$\begin{aligned} F_{k,mn} &\equiv i(-1)^{m+n+1} \begin{pmatrix} \beta_k^{-3n}, & \beta_k^{-n}, & \beta_k^n, & \beta_k^{3n} \end{pmatrix} m_k \begin{pmatrix} \beta_k^{3m} \\ \beta_k^m \\ \beta_k^{-m} \\ \beta_k^{-3m} \end{pmatrix} = \\ &= \frac{2}{\chi_k} e^{i(m+n)\pi} \left\{ \sin\left(\frac{3n-m}{k}\pi\right) + \sin\left(\frac{3m-n}{k}\pi\right) + \sin\left(\frac{3(n+m)}{k}\pi\right) + \right. \\ &\quad \left. + \varsigma_k \sin\left(\frac{n+m}{k}\pi\right) + \lambda_k \left[\sin\left(\frac{3n+m}{k}\pi\right) + \sin\left(\frac{3m+n}{k}\pi\right) \right] \right\}. \end{aligned} \quad (6.25)$$

From such a definition, $F_{k,mn}$ turns out to have the following relevant properties:

$$\begin{aligned} \overline{F_{k,mn}} &= F_{k,mn}; \\ F_{k,mn} &= F_{k,nm}; \\ F_{k,m+k,n} &= F_{k,m,n+k} = (-1)^{k+1} F_{k,mn}; \\ F_{k,mn} &= 0 \text{ if } n+m = k; \\ F_{k,kk} &= 0. \end{aligned} \quad (6.26)$$

Consequently, at most only $\frac{k(k+1)}{2} - \left[\frac{k}{2}\right] - 1$ (real) non-vanishing independent elements of $F_{k,mn}$ exist (where $\left[\frac{k}{2}\right]$ denotes the integer part of $\frac{k}{2}$), even though, as evident from Eqs. (7.1.2), (8.1.2), (9.1.2) and (10.1.2) below, actually such an upper bound is never reached for the allowed values of the Fermat parameter $k = 5, 6, 8, 10$.

The holomorphic superpotential (also named $\mathcal{N} = 2$ holomorphic central charge function) is given by:

$$W_k(\psi; q, p) = \Gamma \Pi_k(\psi), \quad (6.27)$$

where the 1×4 BH charge vector in the symplectic basis is here defined as²⁵

$$\begin{aligned} \Gamma &\equiv (-p^0, -p^1, q_0, q_1) = \tilde{\Gamma} \Sigma, \\ \tilde{\Gamma} &\equiv (q_0, q_1, p^0, p^1). \end{aligned} \quad (6.28)$$

²⁴In the present paper F will be used to denote two different entities:

1) the holomorphic prepotential (with Kähler weights $(4, 0)$) of SK geometry, as defined by Eq. (4.1.5);

2) the infinite rank-2 tensor, defined by Eq. (6.25), introduced in the study of the LG limit in the moduli space of (mirror) Fermat CY_3 s.

Attention should be paid to carefully identify the meaning of F , which however can be understood at a glance in the various frameworks. It should also be recalled that $\mathcal{F}$ denotes the Kähler-invariant holomorphic prepotential of SK geometry, as defined by Eq. (4.1.9).

²⁵Notice the change in the notation of the symplectic charge vectors with respect to the notation used in Sects. 1 and 2. $\tilde{\Gamma}$ defined in the second row of (6.28) has the electric and magnetic charges interchanged with respect to the (case $n_V = 1$ of the) symplectic vector defined by Eq. (1.2).

Using Eqs. (6.14), (6.15) and (6.23), one can obtain the following series expansion (convergent for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$) of the holomorphic superpotential:

$$W_k(\psi; q, p) = A_k \sum_{m=1}^{\infty} C_{k,m-1} \psi^{m-1} N_{k,m}(q, p), \quad (6.29)$$

where the following quantities have been introduced:

$$A_k \equiv -\frac{1}{\chi_k} \frac{(2\pi i)^3}{\text{Ord}(G_k)}; \quad (6.30)$$

$$N_{k,m}(q, p) \equiv (-1)^m [n_{k,1}(q, p) \beta_k^{3m} + n_{k,2}(q, p) \beta_k^m + n_{k,3}(q, p) \beta_k^{-m} + n_{k,4}(q, p) \beta_k^{-3m}], \quad (6.31)$$

where

$$n_k(q, p) \equiv \chi_k \Gamma M_k \in \mathbb{Z}^4 \quad (6.32)$$

is the 1×4 BH charge vector in the PF basis.

By inverting the definition (6.32), one obtains

$$\Gamma = \frac{1}{\chi_k} n_k(q, p) M_k^{-1} \in \mathbb{Z}^4, \quad (6.33)$$

where Γ is defined in Eq. (6.28). Eqs. (6.32)-(6.33) express the change between the symplectic and PF basis of BH charges.

By recalling Eq. (2.2.1) and using Eqs. (6.24) and (6.29), the general form of the “effective BH potential” function $V_{BH,k}(\psi, \bar{\psi}; q, p)$ for the Calabi-Yau threefolds $\mathcal{M}'_k$ s reads

$$\begin{aligned} V_{BH,k}(\psi, \bar{\psi}; q, p) &= \frac{1}{\chi_k^2 F_{k,11}} \left[\exp \left[\tilde{K}_k(\psi, \bar{\psi}) \right] \right] \left[\left| \widetilde{W}_k \right|^2(\psi, \bar{\psi}; q, p) + (g_{\psi\bar{\psi},k}(\psi, \bar{\psi}))^{-1} \left| D_{\psi} \widetilde{W}_k \right|^2(\psi, \bar{\psi}; q, p) \right] \equiv \\ &\equiv \frac{1}{\chi_k^2 F_{k,11}} \tilde{V}_{BH,k}(\psi, \bar{\psi}; q, p), \end{aligned} \quad (6.34)$$

where

$$\begin{aligned} \tilde{K}_k(\psi, \bar{\psi}) &\equiv K_k(\psi, \bar{\psi}) + \ln \left[\frac{(2\pi)^6}{(\text{Ord}(G_k))^2} C_{k,0}^2 F_{k,11} \right]; \\ \widetilde{W}_k(\psi; q, p) &\equiv \frac{W_k(\psi; q, p)}{A_k C_{k,0}}. \end{aligned} \quad (6.35)$$

Remark

It is worth spending a few words concerning the holomorphic prepotential $F(X(\psi))$. In the treatment of 1-modulus SK geometry underlying the moduli space of Fermat CY_3 -compactifications, we will assume it to exist. This corresponds to the choice of a symplectic gauge such that the holomorphic function (see Eq. (4.2.1))

$$e_k \equiv \frac{\partial \left(\frac{X_k^1}{X_k^0} \right)}{\partial \psi} = \frac{X_k^0 \partial X_k^1 - X_k^1 \partial X_k^0}{[X_k^0]^2} \quad (6.36)$$

does not vanish, at least for ψ belonging to a suitable neighbourhood of the LG point $\psi = 0$ (see Subsect. 4.5 of the second Ref. of [54]). By specializing Eq. (4.1.10) for $n_V = 1$, one achieves:

$$F(X(\psi)) = \frac{1}{2} [F_0(\psi) X^0(\psi) + F_1(\psi) X^1(\psi)] = \frac{1}{2} [\Pi^1(\psi) \Pi^3(\psi) + \Pi^2(\psi) \Pi^4(\psi)], \quad (6.37)$$

where $\Pi^i(\psi)$ denotes the i -th component ($i = 1, 2, 3, 4$) of the 4×1 symplectic holomorphic period vector $\Pi_k(\psi)$ given by Eq. (6.22).

Consequently, by recalling Eqs. (6.14) and (6.15), $F(X(\psi))$ can be explicitly computed in power series expansion (convergent for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$) for the k -parametrized class of Fermat CY_3 s. By introducing the 4×1 vector

$$\xi_{k,m} = \begin{pmatrix} \xi_{k,m}^1 \\ \xi_{k,m}^2 \\ \xi_{k,m}^3 \\ \xi_{k,m}^4 \end{pmatrix} \equiv M_k \begin{pmatrix} \beta_k^{3m} \\ \beta_k^m \\ \beta_k^{-m} \\ \beta_k^{-3m} \end{pmatrix}, \quad (6.38)$$

one obtains that (in the third line round brackets denote symmetrization with respect to enclosed indices)

$$\begin{aligned} F_k(X_k(\psi)) &= \frac{1}{2} [\Pi_k^1(\psi) \Pi_k^3(\psi) + \Pi_k^2(\psi) \Pi_k^4(\psi)] = \\ &= \frac{1}{2} \frac{(2\pi i)^6}{(Ord(G_k))^2} \sum_{m,n=1}^{\infty} (-1)^{m+n} C_{k,m-1} C_{k,n-1} (\xi_{k,m}^1 \xi_{k,n}^3 + \xi_{k,m}^2 \xi_{k,n}^4) \psi^{m+n-2} = \\ &= -\frac{32\pi^6}{(Ord(G_k))^2} \sum_{m,n=1}^{\infty} (-1)^{m+n} C_{k,m-1} C_{k,n-1} \Phi_{k,(m,n)} \psi^{m+n-2}, \end{aligned} \quad (6.39)$$

$$\Phi_{k,m,n} \equiv (\xi_{k,m}^1 \xi_{k,n}^3 + \xi_{k,m}^2 \xi_{k,n}^4). \quad (6.40)$$

Thence, one can symmetrically truncate with respect to the dummy indices m and n , both ranging up to l . The resulting series expansion is complete up to $\mathcal{O}(\psi^{l-1})$ included; we will denote such a symmetric truncation as

$$F_{k,l}(X_k(\psi)) = -\frac{32\pi^6}{(Ord(G_k))^2} \sum_{m=1}^l \sum_{n=1}^l (-1)^{m+n} C_{k,m-1} C_{k,n-1} \psi^{m+n-2} \Phi_{k,(m,n)}. \quad (6.41)$$

On the other hand, by recalling Eq. (4.1.9) and Eq. (6.22), the Kähler gauge-invariant holomorphic prepotential reads

$$\mathcal{F}_k(\psi) \equiv F_k\left(\frac{X_k^1(\psi)}{X_k^0(\psi)}\right) = \frac{F_k(X_k^0(\psi), X_k^1(\psi))}{(X_k^0(\psi))^2} = \frac{1}{2} \frac{\sum_{m,n=1}^{\infty} (-1)^{m+n} C_{k,m-1} C_{k,n-1} \Phi_{k,(m,n)} \psi^{m+n-2}}{\left[\sum_{r=1}^{\infty} (-1)^r C_{k,r-1} \xi_{k,r}^1 \psi^{r-1}\right]^2}. \quad (6.42)$$

The vector $\xi_{k,m}$, as well as the infinite rank-2 tensor $\Phi_{k,m,n}$ and the holomorphic prepotentials $F_{k,l}(X_k(\psi))$ and $\mathcal{F}_k(\psi)$ will be computed for each value of the Fermat parameter $k = 5, 6, 8, 10$ (classifying the mirror Fermat CY_3 s $\mathcal{M}'_k$ s) in Subsubsects. 7.1.1, 8.1.1, 9.1.1 and 10.1.1, respectively.

As previously pointed out, it is here worth remarking that in the general treatment of (mirror) Fermat CY_3 s given above (and explicitied the next Sects. 7-10) $t_k(\psi) \equiv X_k^1(\psi)/X_k^0(\psi)$ is *not* a special coordinate, *i.e.* $t_k(\psi) \neq \psi$, and moreover the Kähler gauge is left unfixed (in particular, $X_k^0(\psi) \neq 1$).

The next Sects. 7-10 will be devoted to consistently solving the 1-modulus AEs (2.2.2) (with covariant derivatives given by Eqs. (2.2.3)-(2.2.5)) near the LG point $\psi = 0$, using all the formal machinery elaborated above in the framework of 1-modulus SK geometry underlying the moduli space of Fermat CY_3 -compactifications. In other words, we will solve the criticality condition for the “effective BH potential” (6.34) near the LG point $\psi = 0$, obtaining the constraints which define the BH charge configurations supporting the LG point to be a critical point of $V_{BH,k}$ given by Eq. (6.34). Furthermore, we will address the issue of the stability, by inspecting the real form of the 2×2 Hessian matrix, and checking the (un)fulfilling of the relevant stability conditions treated in Subsect. 3.2, as well.

We will find that (beside the $\frac{1}{2}$ -BPS solutions, existing and stable in all cases) for $k = 5, 8$ only non-BPS, $Z \neq 0$ solutions exist, and they are attractors (local *minima* of V_{BH}), whereas for $k = 6, 10$ only non-BPS, $Z = 0$ solutions exist, and they are *not* attractors in a strict sense (since they are *saddle points* of V_{BH}).

7 $k = 5$: Mirror *Quintic* near LG Point

7.1 Geometric Setup

In the case of mirror *quintic* $\mathcal{M}'_5$ it is easy to realize that one has to consider the “effective BH potential” (6.34) (at least) up to $\mathcal{O}(\psi^3)$ (or, as always understood below, $\mathcal{O}(\bar{\psi}^3)$). As a consequence, the AEs (2.2.2) and the Hessian matrix will be known up to $\mathcal{O}(\psi)$.

For $k = 5$ the definitions (6.12) yield

$$C_{5,5l-1} = 0, \quad l \in \mathbb{N}; \quad (7.1.1)$$

moreover, since $F_{5,m+5,n} = F_{5,m,n+5} = F_{5,m,n}$ (see the third of properties (6.26)), the only independent elements of the rank-2 tensor F_5 belong to the 5×5 matrix

$$F_{5,m,n} = \begin{pmatrix} \sqrt{5+2\sqrt{5}} & 0 & 0 & 0 & -\sqrt{5+2\sqrt{5}} \\ 0 & -\sqrt{5-2\sqrt{5}} & 0 & 0 & \sqrt{5-2\sqrt{5}} \\ 0 & 0 & \sqrt{5-2\sqrt{5}} & 0 & -\sqrt{5-2\sqrt{5}} \\ 0 & 0 & 0 & -\sqrt{5+2\sqrt{5}} & \sqrt{5+2\sqrt{5}} \\ -\sqrt{5+2\sqrt{5}} & \sqrt{5-2\sqrt{5}} & -\sqrt{5-2\sqrt{5}} & \sqrt{5+2\sqrt{5}} & 0 \end{pmatrix}. \quad (7.1.2)$$

Let us now write down all the relevant quantities up to the needed order (here and below, unless otherwise specified, we omit the Fermat parameter $k = 5$):

$$\tilde{K} \approx (\sqrt{5}-2) \frac{C_1^2}{C_0^2} \left[\psi \bar{\psi} - \left(\frac{C_2^2}{C_1^2} - \frac{(\sqrt{5}-2)}{2} \frac{C_1^2}{C_0^2} \right) (\psi \bar{\psi})^2 + \frac{C_5 C_0}{C_1^2} (\sqrt{5}+2) (\psi^5 + \bar{\psi}^5) \right] + \mathcal{O}(\psi^6); \quad (7.1.3)$$

$$g \approx (\sqrt{5}-2) \frac{C_1^2}{C_0^2} \left[1 - 4 \left(\frac{C_2^2}{C_1^2} - \frac{(\sqrt{5}-2)}{2} \frac{C_1^2}{C_0^2} \right) \psi \bar{\psi} \right] + \mathcal{O}(\psi^4); \quad (7.1.4)$$

$$\widetilde{W} \approx N_1 + \frac{C_1}{C_0} N_2 \psi + \frac{C_2}{C_0} \bar{N}_2 \psi^2 + \frac{C_3}{C_0} \bar{N}_1 \psi^3 + \mathcal{O}(\psi^5). \quad (7.1.5)$$

Now, by using the formulæ of the general analysis exploited in Sect. 6, we can get the “effective BH potential” and the holomorphic superpotential, as well as their (covariant) derivatives, up to $\mathcal{O}(\psi)$ (notice that in all the treatments of Sects. 7-10 we are interested only in ordinary derivatives of $\tilde{V}_{BH}$, since they coincide with the covariant ones at the critical points of $\tilde{V}_{BH}$):

$$\tilde{W} = N_1 + \frac{C_1}{C_0} N_2 \psi; \quad (7.1.6)$$

$$D\tilde{W} = \frac{C_1}{C_0} \left[N_2 + 2\frac{C_2}{C_1} \bar{N}_2 \psi + \frac{C_1}{C_0} (\sqrt{5} - 2) N_1 \bar{\psi} \right]; \quad (7.1.7)$$

$$D^2\tilde{W} = 2\frac{C_2}{C_0} \bar{N}_2 + 6\frac{C_3}{C_0} \bar{N}_1 \psi + 4\frac{C_2^2}{C_0 C_1} N_2 \bar{\psi}, \quad (7.1.8)$$

$$\begin{aligned} \tilde{V}_{BH} = & |N_1|^2 + (\sqrt{5} + 2)|N_2|^2 + 2\frac{C_1}{C_0} \left[N_2 \bar{N}_1 + (\sqrt{5} + 2) \frac{C_2 C_0}{C_1^2} (\bar{N}_2)^2 \right] \psi + \\ & + 2\frac{C_1}{C_0} \left[\bar{N}_2 N_1 + (\sqrt{5} + 2) \frac{C_2 C_0}{C_1^2} (N_2)^2 \right] \bar{\psi}; \end{aligned} \quad (7.1.9)$$

$$\begin{aligned} \partial\tilde{V}_{BH} = & 2\frac{C_1}{C_0} \left[N_2 \bar{N}_1 + (\sqrt{5} + 2) \frac{C_2 C_0}{C_1^2} (\bar{N}_2)^2 \right] + 6\frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] \bar{N}_1 \bar{N}_2 \psi + \\ & + 2\frac{C_1^2}{C_0^2} \left[|N_1|^2 (\sqrt{5} - 2) + |N_2|^2 \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) \right] \bar{\psi}; \end{aligned} \quad (7.1.10)$$

$$\begin{aligned} \partial^2\tilde{V}_{BH} = & 6\frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] \bar{N}_1 \bar{N}_2 + \\ & + 24\frac{C_3}{C_0} (\bar{N}_1)^2 \psi + 4\frac{C_1}{C_0} \left[\frac{C_2}{C_0} \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) (\bar{N}_2)^2 + \right. \\ & \left. + (\sqrt{5} - 2) \frac{C_1^2}{C_0^2} \left(1 + (\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} + 3(\sqrt{5} + 2)^2 \frac{C_0^3 C_2 C_3}{C_1^5} \right) \bar{N}_1 N_2 \right] \bar{\psi}; \end{aligned} \quad (7.1.11)$$

$$\begin{aligned} \partial\bar{\partial}\tilde{V}_{BH} = & 2\frac{C_1^2}{C_0^2} \left[|N_1|^2 (\sqrt{5} - 2) + |N_2|^2 \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) \right] + \\ & + 4\frac{C_1}{C_0} \left[\frac{C_2}{C_0} \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) (\bar{N}_2)^2 + \right. \\ & + (\sqrt{5} - 2) \frac{C_1^2}{C_0^2} \left(1 + (\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} + 3(\sqrt{5} + 2)^2 \frac{C_0^3 C_2 C_3}{C_1^5} \right) \bar{N}_1 N_2 \left. \right] \psi + \\ & + 4\frac{C_1}{C_0} \left[\frac{C_2}{C_0} \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) (N_2)^2 + \right. \\ & \left. + (\sqrt{5} - 2) \frac{C_1^2}{C_0^2} \left(1 + (\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} + 3(\sqrt{5} + 2)^2 \frac{C_0^3 C_2 C_3}{C_1^5} \right) N_1 \bar{N}_2 \right] \bar{\psi}. \end{aligned} \quad (7.1.12)$$

7.1.1 The Holomorphic Prepotentials F and $\mathcal{F}$

In order to compute the holomorphic prepotential $F(X(\psi))$ and its “Kähler-invariant counterpart” $\mathcal{F}(\psi)$ defined by Eq. (4.1.9), we have to recall the general formulæ for (mirror) Fermat CY_3 s introduced in Sect. 6, and specialize them for $k = 5$. By recalling the definitions (6.38), (6.40), (6.41), (6.42), one can compute:

$$\xi_{5,m} = \begin{pmatrix} \frac{1}{5} [3 + 2\cos(\frac{2\pi m}{5})] [5\cos(\frac{\pi m}{5}) - 11i\sin(\frac{\pi m}{5})] \\ -\cos(\frac{\pi m}{5}) + i\sin(\frac{\pi m}{5}) \\ 2 [3 + 2\cos(\frac{2\pi m}{5})] [\cos(\frac{\pi m}{5}) - 2i\sin(\frac{\pi m}{5})] \\ 2i\sin(\frac{\pi m}{5}) \end{pmatrix}; \quad (7.1.1.1)$$

$$\Phi_{5,m,n} = \frac{1}{5} \left\{ \begin{aligned} & \left[\begin{aligned} & -24\cos\left(\frac{\pi(n-3m)}{5}\right) - 24\cos\left(\frac{\pi(3n-m)}{5}\right) + 64\cos\left(\frac{\pi(3n+m)}{5}\right) + 64\cos\left(\frac{\pi(n+3m)}{5}\right) + \\ & -13\cos\left(\frac{\pi(n-m)}{5}\right) - 17\cos\left(\frac{3\pi(n-m)}{5}\right) + 173\cos\left(\frac{\pi(n+m)}{5}\right) + 27\cos\left(\frac{3\pi(n+m)}{5}\right) \end{aligned} \right] + \\ & -i \left[\begin{aligned} & -21\sin\left(\frac{\pi(n-3m)}{5}\right) + 21\sin\left(\frac{\pi(3n-m)}{5}\right) + 63\sin\left(\frac{\pi(3n+m)}{5}\right) + 63\sin\left(\frac{\pi(n+3m)}{5}\right) + \\ & + 173\sin\left(\frac{\pi(n+m)}{5}\right) + 21\sin\left(\frac{3\pi(n+m)}{5}\right) \end{aligned} \right] \end{aligned} \right\} = \Phi_{5,(m,n)}; \quad (7.1.1.2)$$

$$\begin{aligned} F_{k=5,l=5}(X_5(\psi)) &= -\frac{32\pi^6}{5^6} \sum_{m=1}^5 \sum_{n=1}^5 (-1)^{m+n} C_{5,m-1} C_{5,n-1} \psi^{m+n-2} \Phi_{5,m,n} = \\ &= -\frac{32\pi^8}{5^6} \left[\begin{aligned} & -\frac{43+11\sqrt{5}+i\sqrt{11890+5302\sqrt{5}}}{\Gamma^{10}(\frac{4}{5})} + \frac{4(115-5\sqrt{5}+11i\sqrt{10(5+\sqrt{5})})}{\Gamma^5(\frac{3}{5})\Gamma^5(\frac{4}{5})} \psi + \\ & -5 \left[\frac{20(-40+3\sqrt{5}-i\sqrt{5725-2510\sqrt{5}})}{(5+\sqrt{5})\Gamma^{10}(\frac{3}{5})} - \frac{2\sqrt{5}(5+23\sqrt{5}+11i\sqrt{10-2\sqrt{5}})}{\Gamma^5(\frac{2}{5})\Gamma^5(\frac{4}{5})} \right] \psi^2 + \\ & + \frac{625\sqrt{2050+710\sqrt{5}}}{16\pi^5} \psi^3 + \\ & + 125 \left[\frac{5(-40+3\sqrt{5}+i\sqrt{5725-2510\sqrt{5}})}{(5+\sqrt{5})\Gamma^{10}(\frac{2}{5})} - \frac{2\sqrt{5}(5+23\sqrt{5}-11i\sqrt{10-2\sqrt{5}})}{3\Gamma^5(\frac{1}{5})\Gamma^5(\frac{3}{5})} \right] \psi^4 \end{aligned} \right] + \mathcal{O}(\psi^5); \quad (7.1.1.3) \end{aligned}$$

$$\begin{aligned}
\mathcal{F}_5(\psi) &\equiv F_5\left(\frac{X_5^1(\psi)}{X_5^0(\psi)}\right) = \frac{F_5(X_5^0(\psi), X_5^1(\psi))}{(X_5^0(\psi))^2} = \frac{1}{2} \frac{\sum_{m,n=1}^{\infty} (-1)^{m+n} C_{5,m-1} C_{5,n-1} \Phi_{5,m,n} \psi^{m+n-2}}{\left[\sum_{r=1}^{\infty} (-1)^r C_{5,r-1} \xi_{5,r}^1 \psi^{r-1}\right]^2} \approx \\
&\approx \frac{\Phi_{5,1,1}}{2(\xi_{5,1}^1)^2} \left[1 - 2 \frac{C_{5,1} \Phi_{5,2,1}}{C_{5,0} \Phi_{5,1,1}} \psi + \mathcal{O}(\psi^2)\right] \left[1 + 2 \frac{C_{5,1} \xi_{5,2}^1}{C_{5,0} \xi_{5,1}^1} \psi + \mathcal{O}(\psi^2)\right] \approx \\
&\approx \frac{\Phi_{5,1,1}}{2(\xi_{5,1}^1)^2} \left[1 + 2 \left(\frac{\xi_{5,2}^1}{\xi_{5,1}^1} - \frac{\Phi_{5,2,1}}{\Phi_{5,1,1}}\right) \frac{C_{5,1}}{C_{5,0}} \psi\right] + \mathcal{O}(\psi^2),
\end{aligned} \tag{7.1.1.4}$$

where in the formula (7.1.1.4) for the Kähler-invariant holomorphic prepotential $\mathcal{F}_5$ only the leading terms in the *LG limit* $\psi \rightarrow 0$ have been retained. Eq. (7.1.1.3) is the exact expression (complete up to $\mathcal{O}(\psi^4)$ included) of the holomorphic prepotential F_5 in a suitable neighbourhood of the LG point of the moduli space of the (mirror) *quintic* $\mathcal{M}'_5$.

7.2 “Criticality Condition” Approach and Attractor Equation

Let us now find the solutions of the AE $\partial \tilde{V}_{BH}(\psi, \bar{\psi}; q, p) = 0$, and check their stability. Since we are working (very) near the LG point, by using Eq. (7.1.9) we can rewrite the AE for $\mathcal{M}'_5$ as follows:

$$N_2 \bar{N}_1 + (\sqrt{5} + 2) \frac{C_2 C_0}{C_1^2} (\bar{N}_2)^2 \approx 0. \tag{7.2.1}$$

Here we simply put $\psi = 0$ in Eq. (7.1.10). Solving Eq. (7.2.1), we will find one (or more) set(s) of BH charges supporting $\psi \approx 0$ to be a critical point of V_{BH} . As understood throughout all our treatment of Sects. 7-10, *ça va sans dire* that actual BH charges are very close to the found one, and also that the critical value of ψ may not be zero, rather it may belong to a suitable neighbourhood of the LG point.

7.3 Critical Hessian of V_{BH}

The stability of the critical point $\psi \approx 0$ of V_{BH} is governed by the real form of the symmetric Hessian 2×2 matrix of V_{BH} evaluated at the considered extremum; in general, it reads (see Eq. (3.1.2))

$$H_{\text{real form}}^{V_{BH}} \equiv \begin{pmatrix} \mathcal{A} & \mathcal{C} \\ \mathcal{C} & \mathcal{B} \end{pmatrix}, \tag{7.3.1}$$

where $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathbb{R}$ are given in terms of $\partial^2 V_{BH}, \partial \bar{\partial} V_{BH}, \bar{\partial}^2 V_{BH} \in \mathbb{C}$ by (the $n_V = 1$ case of) Eq. (3.1.11). By using such an Eq., and also Eqs. (7.1.11)-(7.1.12) evaluated along the criticality condition (7.2.1), it can be computed that

the components of $H_{\text{real form}}^{\tilde{V}_{BH}}$ constrained by the AE (7.2.1) read as follows:

$$\begin{aligned} \mathcal{A} \approx & 2 \frac{C_1^2}{C_0^2} \left[|N_1|^2 (\sqrt{5} - 2) + |N_2|^2 \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) \right] + \\ & + 3 \frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] (\bar{N}_1 \bar{N}_2 + N_1 N_2); \end{aligned} \quad (7.3.2)$$

$$\begin{aligned} \mathcal{B} \approx & 2 \frac{C_1^2}{C_0^2} \left[|N_1|^2 (\sqrt{5} - 2) + |N_2|^2 \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) \right] + \\ & - 3 \frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] (\bar{N}_1 \bar{N}_2 + N_1 N_2); \end{aligned} \quad (7.3.3)$$

$$\mathcal{C} \approx -3i \frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] (\bar{N}_1 \bar{N}_2 - N_1 N_2). \quad (7.3.4)$$

The resulting real eigenvalues of $H_{\text{real form}}^{\tilde{V}_{BH}}$ constrained by the AE (7.2.1) read:

$$\begin{aligned} \lambda_{\pm} \approx & 2 \frac{C_1^2}{C_0^2} \left[|N_1|^2 (\sqrt{5} - 2) + |N_2|^2 \left(1 + 4(\sqrt{5} + 2) \frac{C_0^2 C_2^2}{C_1^4} \right) \right] + \\ & \pm 6 \frac{C_2}{C_0} \left[1 + (\sqrt{5} + 2) \frac{C_3 C_0}{C_2 C_1} \right] |N_1| |N_2|. \end{aligned} \quad (7.3.5)$$

By recalling Eq. (6.34) and using Eq. (7.1.9) with $\psi \approx 0$ and constrained by the AE (7.2.1), one obtains that the purely charge-dependent LG critical values of the “effective BH potential” for the mirror *quintic* $\mathcal{M}'_5$ are

$$V_{BH, LG-critical, k=5} \approx \frac{1}{25\sqrt{5} + 2\sqrt{5}} \left[|N_1|^2 + (\sqrt{5} + 2) |N_2|^2 \right]; \quad (7.3.6)$$

by recalling formula (1.3), this directly yields the following purely charge-dependent values of the BH entropy at the LG critical points of $V_{BH,5}$ in the moduli space of $\mathcal{M}'_5$:

$$S_{BH, LG-critical, k=5} \approx \frac{\pi}{25\sqrt{5} + 2\sqrt{5}} \left[|N_1|^2 + (\sqrt{5} + 2) |N_2|^2 \right] \approx 0.013\pi \left[|N_1|^2 + (\sqrt{5} + 2) |N_2|^2 \right]. \quad (7.3.7)$$

By recalling Eqs. (6.2) and (6.12), it is helpful to write down here the numerical values of constants relevant to our treatment:

$$C_0 \approx 2.5, \quad C_1 \approx 2.25, \quad C_2 \approx 0.77, \quad C_3 \approx 0.054. \quad (7.3.8)$$

7.4 Solutions to Attractor Equation

Let us now analyze more in depth the species of LG attractor points arising from the AE (7.2.1). As it can be easily seen, the AE (7.2.1) has two *non-degenerate* solutions (*i.e.* with non-vanishing V_{BH} and therefore with non-vanishing BH entropy, see Eq. (1.3)):

I. The first non-degenerate solution to AE (7.2.1) is

$$N_2 \approx 0. \quad (7.4.1)$$

As one can see from Eqs. (7.1.6)-(7.1.7), such a solution corresponds to a $\frac{1}{2}$ -BPS LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} = 0$). From the definition (6.31) with $(k, m) = (5, 2)$, in order to get the solution (7.4.1), one has to fine-tune two PF BH charges out of four in the following way:

$$n_3 \approx \frac{1}{2}(1 + \sqrt{5})(n_2 - n_1), \quad n_4 \approx -\frac{1}{2}(1 + \sqrt{5})n_1 + n_2. \quad (7.4.2)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition $N_1 \neq 0$. The real eigenvalues (7.3.5) for the $\frac{1}{2}$ -BPS critical solution coincide and, as it is well known [39, 15, 16], are strictly positive:

$$\lambda_{+, \frac{1}{2}-BPS} = \lambda_{-, \frac{1}{2}-BPS} \approx 2(\sqrt{5} - 2) \frac{C_1^2}{C_0^2} |N_1|_{N_2 \approx 0}^2 > 0. \quad (7.4.3)$$

Consequently, the $\frac{1}{2}$ -BPS LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (7.4.2) is a stable extremum, since it is a minimum of V_{BH} , and it is therefore an attractor in a strict sense. The classical (Bekenstein-Hawking) BH entropy of such a (class of) $\frac{1}{2}$ -BPS LG attractor(s) takes the value

$$S_{BH, \frac{1}{2}-BPS} = \pi V_{BH, \frac{1}{2}-BPS} \approx 0.013\pi |N_1|_{N_2 \approx 0}^2, \quad (7.4.4)$$

where $|N_1|_{N_2 \approx 0}$ is given by the general formula (6.31) with $(k, m) = (5, 1)$ constrained by Eq. (7.4.2):

$$|N_1|_{N_2 \approx 0} \approx \frac{n_1}{8} \left[4\sqrt{5} - i\sqrt{2(5 + \sqrt{5})^3} \right] - \frac{n_2}{4} \left[5 + \sqrt{5} - i\sqrt{10(5 - \sqrt{5})} \right]. \quad (7.4.5)$$

II. The second non-degenerate solution to AE (7.2.1) is

$$|N_1| \approx \xi |N_2|, \quad \xi \equiv (\sqrt{5} + 2) \frac{C_0 C_2}{C_1^2} \approx 1, 6; \quad (7.4.6)$$

$$\arg(N_1) - 3\arg(N_2) \approx \pi,$$

where N_1 and N_2 are given by the general formula (6.31) with $k = 5$ and $m = 1, 2$, respectively:

$$N_1 = \frac{\sqrt{5} - 1}{4} \left(n_1 + n_4 - \frac{(3 + \sqrt{5})}{2} (n_2 + n_3) \right) + \frac{i}{2} \sqrt{\frac{(5 + \sqrt{5})}{2}} \left(n_4 - n_1 + \frac{(\sqrt{5} - 1)}{2} (n_3 - n_2) \right), \quad (7.4.7)$$

$$N_2 = -\frac{\sqrt{5} + 1}{4} \left(n_1 + n_4 - \frac{(3 - \sqrt{5})}{2} (n_2 + n_3) \right) + \frac{i}{2} \sqrt{\frac{(5 - \sqrt{5})}{2}} \left(n_4 - n_1 - \frac{(\sqrt{5} + 1)}{2} (n_3 - n_2) \right). \quad (7.4.8)$$

The first of fine-tuning conditions (7.4.6) substituted in Eq. (7.3.7) yields

$$S_{BH,non-BPS,Z \neq 0} \approx \frac{\pi}{25\sqrt{5+2\sqrt{5}}} (\xi^2 + \sqrt{5} + 2) |N_2|^2 \approx 0.088\pi |N_2|^2. \quad (7.4.9)$$

As one can see from Eqs. (7.1.6)-(7.1.7), the solution (7.4.6) corresponds to a non-BPS, $Z \neq 0$ LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} \neq 0$). The real eigenvalues (7.3.5) for such a non-BPS, $Z \neq 0$ critical solution read

$$\lambda_{\pm,non-BPS,Z \neq 0} \approx 2 \frac{C_1^2}{C_0^2} |N_2|^2 \left[1 + 5(\sqrt{5} - 2)\xi^2 \pm 3 \left(\xi(\sqrt{5} - 2) + (\sqrt{5} + 2) \frac{C_0^2 C_3}{C_1^3} \right) \xi \right]. \quad (7.4.10)$$

Substituting the numerical values (7.3.8) of the involved constants in Eq. (7.4.10), one reaches the conclusion that both $\lambda_{\pm,non-BPS,Z \neq 0}$ are strictly positive:

$$\lambda_{\pm,non-BPS,Z \neq 0} \approx 2|N_2|^2 [3.277 \pm 1.97] > 0. \quad (7.4.11)$$

Thus, the non-BPS, $Z \neq 0$ LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (7.4.6)-(7.4.8) is a minimum of V_{BH} and consequently an attractor in a strict sense.

Let us now find the fine-tuning conditions for PF BH charges supporting the considered non-BPS, $Z \neq 0$ LG attractor for the mirror *quintic* $\mathcal{M}'_5$. This amounts to solving Eqs. (7.4.6)-(7.4.8) by recalling the definitions (6.31) and (6.32). By doing so, one gets the following three different sets of constraining relations on the PF BH charges:

$$\text{II.1 : } \left\{ \begin{array}{l} \frac{n_{2,\pm}}{n_1} = \frac{a_2 - a_{1,\pm} - 1}{a_{1,\pm} + 2} \approx \begin{cases} \text{"+" : } 0.342 \\ \text{"-" : } -138.3 \end{cases} , \\ \frac{n_{3,\pm}}{n_1} = -\frac{a_2 + a_{1,\pm} - 1}{a_{1,\pm} + 2} \approx \begin{cases} \text{"+" : } -1.352 \\ \text{"-" : } 35 \end{cases} , \\ \frac{n_{4,\pm}}{n_1} = \frac{a_{1,\pm} - 2}{a_{1,\pm} + 2} \approx \begin{cases} \text{"+" : } 0.009 \\ \text{"-" : } 102.3 \end{cases} , \end{array} \right. \quad (7.4.12)$$

$$\left\{ \begin{array}{l} a_{1,\pm}(\xi) \equiv \pm 2 \sqrt{\frac{\sqrt{5}(1+3\xi)}{2-\sqrt{5}+\xi[-\sqrt{5}-4\xi+2(1+\sqrt{5})\xi^2]}} \approx \pm 2.04, \\ a_2(\xi) \equiv \frac{\sqrt{5\sqrt{5}(2+\sqrt{5})(1+\sqrt{2}\xi)}}{\sqrt{\sqrt{5}(\sqrt{5}-2)+\sqrt{2\sqrt{5}(\sqrt{5}+2)}\xi}} \approx 2.92, \\ n_1 \neq 0; \end{array} \right. \quad (7.4.13)$$

$$\text{II.2 : } \left\{ \begin{array}{l} n_2 = n_1 \frac{(\sqrt{5}-1)-\xi(\sqrt{5}+1)}{(\sqrt{5}+1)-\xi(\sqrt{5}-1)} \approx -3.13n_1, \\ n_3 = n_2, \\ n_4 = n_1; \end{array} \right.$$

$$\text{II.3 : } \left\{ \begin{array}{l} \left\{ \begin{array}{l} n_{2,\pm} + n_{3,\pm} - n_{1,\pm} - n_{4,\pm} = a_{\pm}, \\ n_{1,\pm} + n_{2,\pm} + n_{3,\pm} + n_{4,\pm} = b_{\pm}, \\ 2n_{3,\pm} - 2n_{2,\pm} + d = c, \\ n_{4,\pm} - n_{1,\pm} = d, \end{array} \right. \iff \left\{ \begin{array}{l} n_{1,\pm} = \frac{1}{4}(-a_{\pm} + b_{\pm} - 2d), \\ n_{2,\pm} = \frac{1}{4}(a_{\pm} + b_{\pm} - c + d), \\ n_{3,\pm} = \frac{1}{4}(a_{\pm} + b_{\pm} + c - d), \\ n_{4,\pm} = \frac{1}{4}(-a_{\pm} + b_{\pm} + 2d), \end{array} \right. \\ \\ a_{\pm}(\xi; b_{\pm}, c, d) \equiv -\frac{\sqrt{5+2\sqrt{5}c}-\sqrt{5(5-2\sqrt{5})d}+\sqrt{2(5+\sqrt{5})(-c+\sqrt{5}d)\xi}}{\sqrt{5(5-2\sqrt{5})c}-5\sqrt{(5+2\sqrt{5})d}+\sqrt{2(5+\sqrt{5})(\sqrt{5}c-5d)\xi}} b_{\pm}(\xi; c, d) \approx \\ \approx \left(0.26157 + \frac{0.213c}{-0.3325c+d}\right) b_{\pm}(\xi; c, d) \approx \pm 0.49 \sqrt{\frac{c-0.98995d}{c-2.89698d}} \frac{(c-3.0102d)(c-3.9798d)}{c-3.0077d}, \\ \\ b_{\pm}(\xi; c, d) \equiv \pm \frac{1}{2} \sqrt{\frac{[5-2\sqrt{5}+2\xi(5-\sqrt{5}+(5+\sqrt{5})\xi)][2\sqrt{5}+\sqrt{5}(1+\sqrt{5})\xi]}{[-2\sqrt{5}+3\sqrt{5}(1+\sqrt{5})\xi]}} \left[c - \frac{\sqrt{5}d(1+\sqrt{5}+4\xi)}{3-\sqrt{5}+4\xi} \right] \sqrt{\frac{\left[c - \frac{\sqrt{5}d(-2+(\sqrt{5}+1)\xi)}{2+(\sqrt{5}+1)\xi} \right]}{\left[c - \frac{\sqrt{5}d(2+3(\sqrt{5}+1)\xi)}{-2+3(\sqrt{5}+1)\xi} \right]}} \approx \\ \approx \pm 2.481 (c - 3.00769d) \sqrt{\frac{c-0.98995d}{c-2.89698d}}. \end{array} \right. \quad (7.4.14)$$

By using the definitions (6.32) and (6.33) for the case $k = 5$, one obtains (recall the first of definitions (6.28))

$$\begin{aligned} (n_1, n_2, n_3, n_4) &= (3p^0 - 5q_0, p^0 + 5q_1, -21p^0 + 5p^1 + 40q_0 - 5q_1, -8p^0 + 15q_0) \\ &\quad \Updownarrow \\ \Gamma &= (-3n_1 - n_4, \frac{1}{5}(4n_1 - n_2 - n_3 + 4n_4), \frac{1}{5}(8n_1 + 3n_4), \frac{1}{5}(-3n_1 + n_2 - n_4)). \end{aligned} \quad (7.4.15)$$

7.4.1 Analysis of non-BPS, $Z \neq 0$ supporting BH charge configurations

Let us analyze more in depth the obtained supporting BH charge configurations.

II.1. This configuration is characterized by a physically meaningful “ $\pm$ ” split, affecting the charge-dependent expression of the BH entropy. By using Eqs. (7.4.8), (7.4.9) and (7.4.15), Eqs. (7.4.12) yield

$$\begin{aligned} N_{2,II.1,\pm} &= \left[-\frac{\sqrt{5}}{2} (3 + \sqrt{5}) \frac{a_{1,\pm}}{a_{1,\pm}+2} + \frac{i}{2} (1 + \sqrt{5}) \sqrt{\frac{\sqrt{5}(\sqrt{5}-1)}{2}} \frac{a_{2,\pm}}{a_{1,\pm}+2} \right] n_1 \approx \\ &\approx \left[\left\{ \begin{array}{l} \text{“+”} : -2.95603 \\ \text{“-”} : -298.5592 \end{array} \right. + i \left\{ \begin{array}{l} \text{“+”} : 0.322 \\ \text{“-”} : -32.5229 \end{array} \right. \right] (3p^0 - 5q_0); \\ \\ S_{BH,non-BPS,Z \neq 0,II.1,\pm} &\approx 0.088\pi |N_{2,II.1,\pm}|^2 \approx \left\{ \begin{array}{l} \text{“+”} : 0.121\pi (3p^0 - 5q_0)^2. \\ \text{“-”} : 1192.1\pi (3p^0 - 5q_0)^2. \end{array} \right. \end{aligned} \quad (7.4.1.1)$$

In order to give an example of the range of the symplectic (electric/magnetic) charges in the considered BH charge configuration, by recalling that PF BH charges are integers as are the symplectic BH charges (see definitions (6.32)-(6.33)), one can fulfill the fine-tuning conditions (7.4.12) by taking *e.g.*

$$n_1 = 10 \quad \Downarrow \quad n_{2,\pm} \approx \begin{cases} \text{"+"} : 3 \\ \text{"-"} : -1383 \end{cases}, n_{3,\pm} \approx \begin{cases} \text{"+"} : -14 \\ \text{"-"} : 350 \end{cases}, n_{4,\pm} \approx \begin{cases} \text{"+"} : 0 \\ \text{"-"} : 1023 \end{cases} \quad (7.4.1.2)$$

Using Eq. (7.4.15), one finally gets

$$\Gamma_{\pm} \approx \begin{cases} \text{"+"} : (-30, 51, 16, -5); \\ \text{"-"} : (-1053, 1033, 630, -487). \end{cases} \quad (7.4.1.3)$$

II.2. This configuration is the only one not having any kind of split, and having a purely real N_2 . Also, it is the one adopted in [9] (see in particular Sect. 4 and App. C of such a Ref.). By using Eqs. (7.4.8), (7.4.9) and (7.4.15), Eqs. (7.4.13) yield

$$N_{2,II.2} = -\frac{2\sqrt{5}}{1+\sqrt{5}+(1-\sqrt{5})\xi} n_1 \approx -3.554 (3p^0 - 5q_0);$$

$$S_{BH,non-BPS,Z \neq 0,II.2} \approx 0.088\pi (N_{2,II.2})^2 \approx 1.111\pi (3p^0 - 5q_0)^2. \quad (7.4.1.4)$$

Concerning the range of the symplectic (electric/magnetic) charges in the considered BH charge configuration, by repeating the same procedure exploited above for the case II.1, one achieves the same results obtained at the end of Sect. 4 of [9].

II.3. This case, beside being split in “ $\pm$ ” (as the case II.1), is also parameterized by the couple $(c, d) \in \mathbb{Z}^2$ ($\mathbb{R}^2$ in the -semi- classical limit of large charges), further constrained by the request $a_{\pm}(b_{\pm}, c, d)$, $b_{\pm}(c, d) \in \mathbb{Z}$ ($\mathbb{R}$ for large charges), implying that

$$c \neq 3.077d, \quad \text{and} \quad \begin{cases} \begin{cases} c \geq 0.98995d, \\ c > 2.89698d; \end{cases} \\ \text{or} \\ \begin{cases} c \leq 0.98995d, \\ c < 2.89698d. \end{cases} \end{cases} \quad (7.4.1.5)$$

By using Eq. (7.4.8), Eqs. (7.4.14) yield

$$\begin{aligned}
N_{2,II.3,\pm}(c,d) &= -\frac{1}{4} [-\sqrt{5}a_{\pm}(c,d) + b_{\pm}(c,d)] - \frac{i}{4}\sqrt{\frac{5+\sqrt{5}}{2}} [c - \sqrt{5}d] = \\
&= -\frac{1}{4} \left[\frac{\sqrt{5+2\sqrt{5}+\sqrt{5-2\sqrt{5}}}}{\sqrt{5-2\sqrt{5}+\sqrt{5(2+2\sqrt{5})}}\xi} \left[\frac{(c-\sqrt{5}d)}{c-\frac{(\sqrt{5+2\sqrt{5}d+\sqrt{5(2+2\sqrt{5})}\xi)}}{(\sqrt{5-2\sqrt{5}+\sqrt{5(2+2\sqrt{5})}\xi)}}\sqrt{5}d} \right] b_{\pm}(c,d) \right] + \\
&\quad -\frac{i}{4}\sqrt{\frac{5+\sqrt{5}}{2}} (c - \sqrt{5}d) = \\
&= -\frac{1}{4\sqrt{2}} \left\{ \pm \left\{ \sqrt{\frac{(1+\sqrt{5})[5-2\sqrt{5}+2\xi(5-\sqrt{5}+(5+\sqrt{5})\xi)][2\sqrt{5}+\sqrt{5}(1+\sqrt{5})\xi]}{[-2\sqrt{5}+3\sqrt{5}(1+\sqrt{5})\xi][(\sqrt{5}-2)+2\sqrt{2}\sqrt{3-\sqrt{5}\xi+2(1+\sqrt{5})\xi^2}]}} \cdot \right. \right. \\
&\quad \cdot \left. \sqrt{\frac{c-\frac{-2+(\sqrt{5}+1)\xi}{2+(\sqrt{5}+1)\xi}\sqrt{5}d}{c-\frac{2+3(\sqrt{5}+1)\xi}{-2+3(\sqrt{5}+1)\xi}\sqrt{5}d} \frac{c-\frac{1+\sqrt{5}+4\xi}{3-\sqrt{5}+4\xi}\sqrt{5}d}{c-\frac{(\sqrt{5+2\sqrt{5}d+\sqrt{5(2+2\sqrt{5})}\xi)}}{(\sqrt{5-2\sqrt{5}+\sqrt{5(2+2\sqrt{5})}\xi)}}\sqrt{5}d} \right. \right\} + i\sqrt{5+\sqrt{5}} \left. \right\} (c - \sqrt{5}d) \approx \\
&\approx [\mp 0.34631\sqrt{\frac{c-0.98995d}{c-2.89698d}} - 0.47552i] (c - 2.23607d);
\end{aligned} \tag{7.4.1.6}$$

$$\begin{aligned}
|N_{2,II.3,\pm}(c,d)|^2 &= \\
&= \frac{1}{32} \left\{ \left\{ \frac{(1+\sqrt{5})[5-2\sqrt{5}+2\xi(5-\sqrt{5}+(5+\sqrt{5})\xi)][2\sqrt{5}+\sqrt{5}(1+\sqrt{5})\xi]}{[-2\sqrt{5}+3\sqrt{5}(1+\sqrt{5})\xi][(\sqrt{5}-2)+2\sqrt{2}\sqrt{3-\sqrt{5}\xi+2(1+\sqrt{5})\xi^2}]} \cdot \right. \right. \\
&\quad \cdot \left. \frac{c-\frac{-2+(\sqrt{5}+1)\xi}{2+(\sqrt{5}+1)\xi}\sqrt{5}d}{c-\frac{2+3(\sqrt{5}+1)\xi}{-2+3(\sqrt{5}+1)\xi}\sqrt{5}d} \frac{[c-\frac{1+\sqrt{5}+4\xi}{3-\sqrt{5}+4\xi}\sqrt{5}d]^2}{[c-\frac{(\sqrt{5+2\sqrt{5}d+\sqrt{5(2+2\sqrt{5})}\xi)}}{(\sqrt{5-2\sqrt{5}+\sqrt{5(2+2\sqrt{5})}\xi)}}\sqrt{5}d]^2} \right. \right\} + (5+\sqrt{5}) \left. \right\} (c - \sqrt{5}d)^2 \approx \\
&\approx [0.11993(c - 0.98995d) + 0.22613(c - 2.89698d)] \frac{(c-2.23607d)^2}{c-2.89698d} \approx \\
&\approx 0.34606 \frac{(c-2.23607d)^3}{c-2.89698d}.
\end{aligned} \tag{7.4.1.7}$$

By recalling Eqs. (7.4.9) and (7.4.15), one finally gets (notice that in this case the “ $\pm$ ” split is unphysical, because it does not affect the charge-dependent expression of the BH entropy) :

$$\begin{aligned}
S_{BH,non-BPS,Z \neq 0,II.3,\pm}(c,d) &\approx 0.088\pi |N_{2,II.3,\pm}(c,d)|^2 \approx 0.03045\pi \frac{(c-2.23607d)^3}{c-2.89698d} \approx \\
&\approx 0.03045\pi \frac{(-30.40325p^0+10p^1+55.27864q_0-20q_1)^3}{-23.13316p^0+10p^1+42.0603q_0-20q_1} = \\
&= S_{BH,non-BPS,Z \neq 0,II.3,\pm}(p^0, p^1, q_0, q_1).
\end{aligned} \tag{7.4.1.8}$$

Thus, by comparing Eqs. (7.4.1.1), (7.4.1.4) and (7.4.1.8), one can conclude that *all* three distinct sets of fine-tuning conditions for PF BH charges (7.4.12)-(7.4.14) do support a non-BPS, $Z \neq 0$ LG attractor in a strict sense, but they yield different charge-dependent expressions for the classical BH entropy $S_{BH, non-BPS, Z \neq 0}$. As non-BPS, $Z \neq 0$ attractor, the LG point of the moduli space of the Fermat mirror *quintic* $\mathcal{M}'_5$ turns out to originate a *threefold splitting* of the supporting BH charges, and correspondingly of the purely charge-dependent classical (Bekenstein-Hawking) BH entropy. This fact confirms the results of the recent investigation in [35].

Finally, by recalling the definition (2.2.11), one can compute the supersymmetry-breaking order parameter for the non-BPS, $Z \neq 0$ LG attractor in the mirror *quintic* $\mathcal{M}'_5$; by using Eqs. (7.1.4), (7.1.6), (7.1.7) and (7.4.6), one gets

$$\mathcal{O}_{non-BPS, Z \neq 0} \equiv \left[\frac{g^{-1}|DW|^2}{|W|^2} \right]_{non-BPS, Z \neq 0} = \left(g^{-1}|D \ln \widetilde{W}|^2 \right)_{non-BPS, Z \neq 0} \approx \frac{(\sqrt{5}+2)}{\xi^2} \approx 1.65, \quad (7.4.1.9)$$

which is consistent with the result obtained at the end of Sect. 4 of [9].

8 $k = 6$: Mirror *Sextic* near LG Point

8.1 Geometric Setup

For the mirror *sextic* $\mathcal{M}'_6$ the computations (*but not the results!*) go the same way as for the mirror *quintic* $\mathcal{M}'_5$. Thus, also for $k = 6$ it is easy to realize that one has to consider the “effective BH potential” (6.34) (at least) up to $\mathcal{O}(\psi^3)$ in order to get the AE (2.2.2) and the Hessian matrix up to $\mathcal{O}(\psi)$.

For $k = 6$ the definitions (6.12) yield

$$C_{6, 3l-1} = 0, \quad l \in \mathbb{N} \quad (8.1.1)$$

moreover, since $F_{6, m+6, n} = F_{6, m, n+6} = -F_{6, m, n}$ (see the third of properties (6.26)), the only independent elements of the rank-2 tensor F_6 belong to the 6×6 matrix

$$F_{6, m, n} = \begin{pmatrix} 2\sqrt{3} & 0 & -\sqrt{3} & 0 & 0 & 3 \\ 0 & -\frac{2}{\sqrt{3}} & 1 & 0 & 0 & -\frac{1}{\sqrt{3}} \\ -\sqrt{3} & 1 & 0 & -1 & \sqrt{3} & -2 \\ 0 & 0 & -1 & \frac{2}{\sqrt{3}} & 0 & -\frac{1}{\sqrt{3}} \\ 0 & 0 & \sqrt{3} & 0 & -2\sqrt{3} & 3 \\ 3 & -\frac{1}{\sqrt{3}} & -2 & -\frac{1}{\sqrt{3}} & 3 & 0 \end{pmatrix}. \quad (8.1.2)$$

Let us now write down all the relevant quantities up to the needed order (here and below, unless otherwise specified, we omit the Fermat parameter $k = 6$):

$$\widetilde{K} \approx \frac{1}{3} \frac{C_1^2}{C_0^2} \left[\psi \bar{\psi} + \frac{1}{6} \frac{C_1^2}{C_0^2} (\psi \bar{\psi})^2 \right] + \mathcal{O}(\psi^6); \quad (8.1.3)$$

$$g \approx \frac{1}{3} \frac{C_1^2}{C_0^2} \left[1 + \frac{2}{3} \frac{C_1^2}{C_0^2} \psi \bar{\psi} \right] + \mathcal{O}(\psi^4); \quad (8.1.4)$$

$$\widetilde{W} \approx N_1 + \frac{C_1}{C_0} N_2 \psi - \frac{C_3}{C_0} \bar{N}_2 \psi^3 - \frac{C_4}{C_0} \bar{N}_1 \psi^4 + \mathcal{O}(\psi^6). \quad (8.1.5)$$

Now, by using the formulæ of the general analysis exploited in Sect. 6, we can get the “effective BH potential” and the holomorphic superpotential, as well as their (covariant) derivatives, up to $\mathcal{O}(\psi)$:

$$\widetilde{W} = N_1 + \frac{C_1}{C_0} N_2 \psi; \quad (8.1.6)$$

$$D\widetilde{W} = \frac{C_1}{C_0} \left[N_2 + \frac{1}{3} \frac{C_1}{C_0} N_1 \bar{\psi} \right]; \quad (8.1.7)$$

$$D^2\widetilde{W} = -6 \frac{C_3}{C_0^3} \bar{N}_2 \psi - 12 \frac{C_4}{C_0^4} \bar{N}_1 \psi^2; \quad (8.1.8)$$

$$\widetilde{V}_{BH} = |N_1|^2 + 3|N_2|^2 + 2 \frac{C_1}{C_0} N_2 \bar{N}_1 \psi + 2 \frac{C_1}{C_0} \bar{N}_2 N_1 \bar{\psi}; \quad (8.1.9)$$

$$\partial \widetilde{V}_{BH} = 2 \frac{C_1}{C_0} N_2 \bar{N}_1 - 18 \frac{C_3}{C_1^3} (\bar{N}_2)^2 \psi + \frac{2}{3} \frac{C_1^2}{C_0^2} (|N_1|^2 + 3|N_2|^2) \bar{\psi}; \quad (8.1.10)$$

$$\partial^2 \widetilde{V}_{BH} = -18 \frac{C_3}{C_1^3} (\bar{N}_2)^2 - 24 \frac{C_3}{C_0^3} \left(1 + \frac{C_4 C_0}{C_3 C_1} \right) \bar{N}_1 \bar{N}_2 \psi + \frac{4}{3} \frac{C_1^3}{C_0^3} \bar{N}_1 N_2 \bar{\psi}; \quad (8.1.11)$$

$$\partial \bar{\partial} \widetilde{V}_{BH} = \frac{2}{3} \frac{C_1^2}{C_0^2} (|N_1|^2 + 3|N_2|^2) + \frac{4}{3} \frac{C_1^3}{C_0^3} \bar{N}_1 N_2 \psi + \frac{4}{3} \frac{C_3}{C_0^3} N_1 \bar{N}_2 \bar{\psi}. \quad (8.1.12)$$

From the definition (6.31) with $k = 6$ and $m = 1, 2$, one gets respectively:

$$N_1 = -\frac{\sqrt{3}}{2} (n_2 + n_3) + \frac{i}{2} (2n_4 - 2n_1 + n_3 - n_2); \quad (8.1.13)$$

$$N_2 = \frac{1}{2} (n_3 + n_2 - 2n_4 - 2n_1) - \frac{\sqrt{3}i}{2} (n_3 - n_2). \quad (8.1.14)$$

8.1.1 The Holomorphic Prepotentials F and $\mathcal{F}$

As done for $k = 5$ in Subsubsect. 7.1.1, in order to compute the holomorphic prepotential $F(X(\psi))$ and its “Kähler-invariant counterpart” $\mathcal{F}(\psi)$ defined by Eq. (4.1.9), we have to recall the general formulæ for (mirror) Fermat CY_3 s introduced in Sect. 6, and specialize them for $k = 6$. By recalling the definitions (6.38), (6.40), (6.41), (6.42), one can compute:

$$\xi_{6,m} = \begin{pmatrix} -\frac{2}{3}i \left[\sin\left(\frac{\pi m}{6}\right) + \sin\left(\frac{\pi m}{2}\right) \right] \\ -\cos\left(\frac{\pi m}{6}\right) + i \sin\left(\frac{\pi m}{6}\right) \\ 4\cos^2\left(\frac{\pi m}{6}\right) \left[\cos\left(\frac{\pi m}{6}\right) - 3i \sin\left(\frac{\pi m}{6}\right) \right] \\ 2i \sin\left(\frac{\pi m}{6}\right) \end{pmatrix}; \quad (8.1.1.1)$$

$$\begin{aligned}
\Phi_{6,m,n} &= \\
&= \left[\begin{aligned} &-\cos\left(\frac{\pi(n-3m)}{6}\right) - \cos\left(\frac{\pi(3n-m)}{6}\right) - 2\cos\left(\frac{\pi(n-m)}{6}\right) - \cos\left(\frac{\pi(n-m)}{2}\right) + \\ &+ \cos\left(\frac{\pi(n+3m)}{6}\right) + \cos\left(\frac{\pi(3n+m)}{5}\right) + 2\cos\left(\frac{\pi(n+m)}{6}\right) + \cos\left(\frac{\pi(n+m)}{2}\right) \end{aligned} \right] + \\
&\quad - \frac{2}{3}i\cos\left(\frac{\pi m}{6}\right) \left[3\sin\left(\frac{\pi n}{6}\right) + 8\cos^3\left(\frac{\pi n}{6}\right) \sin\left(\frac{\pi m}{3}\right) \right] \neq \Phi_{6,(m,n)};
\end{aligned} \tag{8.1.1.2}$$

$$\begin{aligned}
F_{k=6,l=5}(X_6(\psi)) &= -\frac{2\pi^6}{3^6} \sum_{m=1}^5 \sum_{n=1}^5 (-1)^{m+n} C_{6,m-1} C_{6,n-1} \psi^{m+n-2} \Phi_{6,(m,n)} = \\
&= -\frac{2\pi^6}{3^6} \left[-\frac{2^{4/3}(5+2i\sqrt{3})\pi^2}{\Gamma^2(\frac{2}{3})\Gamma^8(\frac{5}{6})} + \frac{24(i+\sqrt{3})\pi}{\Gamma^4(\frac{2}{3})\Gamma^4(\frac{5}{6})} \psi + \frac{3i2^{2/3}(2\sqrt{3}-9i)}{\Gamma^6(\frac{2}{3})} \psi^2 - \frac{32^{2/3}(6+i\sqrt{3})}{\pi\Gamma^2(\frac{2}{3})\Gamma^2(\frac{5}{6})} \psi^3 + \frac{207\sqrt{3}}{4\pi^3} \psi^4 \right] + \mathcal{O}(\psi^5);
\end{aligned} \tag{8.1.1.3}$$

$$\begin{aligned}
\mathcal{F}_6(\psi) &\equiv F_6\left(\frac{X_6^1(\psi)}{X_6^0(\psi)}\right) = \frac{F_6(X_6^0(\psi), X_6^1(\psi))}{(X_6^0(\psi))^2} = \frac{1}{2} \frac{\sum_{m,n=1}^{\infty} (-1)^{m+n} C_{6,m-1} C_{6,n-1} \Phi_{6,(m,n)} \psi^{m+n-2}}{[\sum_{r=1}^{\infty} (-1)^r C_{6,r-1} \xi_{6,r}^1 \psi^{r-1}]^2} \approx \\
&\approx \frac{\Phi_{6,1,1}}{2(\xi_{6,1}^1)^2} \left[1 - \frac{C_{6,1}(\Phi_{6,2,1} + \Phi_{6,1,2})}{C_{6,0}\Phi_{6,1,1}} \psi + \mathcal{O}(\psi^2) \right] \left[1 + 2 \frac{C_{6,1}\xi_{6,2}^1}{C_{6,0}\xi_{6,1}^1} \psi + \mathcal{O}(\psi^2) \right] \approx \\
&\approx \frac{\Phi_{6,1,1}}{2(\xi_{6,1}^1)^2} \left[1 + 2 \left(\frac{\xi_{6,2}^1}{\xi_{6,1}^1} - \frac{\Phi_{6,2,1} + \Phi_{6,1,2}}{2\Phi_{6,1,1}} \right) \frac{C_{6,1}}{C_{6,0}} \psi \right] + \mathcal{O}(\psi^2),
\end{aligned} \tag{8.1.1.4}$$

where in the formula (8.1.1.4) for the Kähler-invariant holomorphic prepotential $\mathcal{F}_6$ only the leading terms in the *LG limit* $\psi \rightarrow 0$ have been retained. Eq. (8.1.1.3) is the exact expression (complete up to $\mathcal{O}(\psi^4)$ included) of the holomorphic prepotential F_6 in a suitable neighbourhood of the LG point of the moduli space of the (mirror) *sextic* $\mathcal{M}'_6$.

8.2 “Criticality Condition” Approach and Attractor Equation

Let us now find the solutions of the AE $\partial\widetilde{V}_{BH}(\psi, \bar{\psi}; q, p) = 0$, and check their stability. Since we are working near the LG point, by using Eq. (8.1.9) we can rewrite the AE for $\mathcal{M}'_6$ as follows:

$$N_2 \bar{N}_1 \approx 0. \tag{8.2.1}$$

Here we simply put $\psi = 0$. Solving Eq. (8.2.1), we will find one (or more) set(s) of BH charges supporting $\psi \approx 0$ to be a critical point of V_{BH} .

8.3 Critical Hessian of V_{BH}

By using (the $n_V = 1$ case of) Eq. (3.1.11) and Eqs. (8.1.11)-(8.1.12) evaluated along the criticality condition (8.2.1), it can be computed that the components of $H_{\text{real form}}^{\tilde{V}_{BH}}$ (given by Eq. (7.3.1)) constrained by the AE (8.2.1) read as follows:

$$\mathcal{A} = \frac{2}{3} \frac{C_1^2}{C_0^2} (|N_1|^2 + 3|N_2|^2) - 9 \frac{C_3}{C_1} ((\bar{N}_2)^2 + (N_2)^2); \quad (8.3.1)$$

$$\mathcal{B} = \frac{2}{3} \frac{C_1^2}{C_0^2} (|N_1|^2 + 3|N_2|^2) + 9 \frac{C_3}{C_1} ((\bar{N}_2)^2 + (N_2)^2); \quad (8.3.2)$$

$$\mathcal{C} = 9i \frac{C_3}{C_1} ((\bar{N}_2)^2 - (N_2)^2). \quad (8.3.3)$$

The resulting real eigenvalues of $H_{\text{real form}}^{\tilde{V}_{BH}}$ constrained by the AE (8.2.1) read:

$$\lambda_{\pm} \approx 2 \frac{C_1^2}{C_0^2} \left[\frac{1}{3} |N_1|^2 + |N_2|^2 \left(1 \pm 9 \frac{C_0^2 C_3}{C_1^3} \right) \right]. \quad (8.3.4)$$

By recalling Eq. (6.34) and using Eq. (8.1.9) with $\psi \approx 0$ and constrained by the AE (8.2.1), one obtains that the purely charge-dependent LG critical values of the “effective BH potential” for the mirror *sextic* $\mathcal{M}'_6$ are

$$V_{BH, LG-critical, k=6} \approx \frac{1}{18\sqrt{3}} [|N_1|^2 + 3|N_2|^2]; \quad (8.3.5)$$

by recalling formula (1.3), this directly yields the following purely charge-dependent values of the BH entropy at the LG critical points of $V_{BH,6}$ in the moduli space of $\mathcal{M}'_6$:

$$S_{BH, LG-critical, k=6} \approx \frac{\pi}{18\sqrt{3}} [|N_1|^2 + 3|N_2|^2]. \quad (8.3.6)$$

Let us write down here the numerical values of constants relevant to our treatment:

$$C_0 \approx 2.27, \quad C_1 \approx 1.52, \quad C_3 \approx -0.247, \quad C_4 \approx 0.054; \quad 9 \frac{C_0^2 C_3}{C_1^3} \approx -3.25. \quad (8.3.7)$$

8.4 Solutions to Attractor Equations

Let us now analyze more in depth the species of LG attractor points arising from the AE (8.2.1). As it can be easily seen, the AE (8.2.1) has two *non-degenerate* solutions:

I. The first non-degenerate solution to AE (8.2.1) is

$$N_2 \approx 0. \quad (8.4.1)$$

This is nothing but the $k = 5$ solution (7.4.1). As one can see from Eq. (8.1.6)-(8.1.7), also for $k = 6$ such a solution corresponds to a $\frac{1}{2}$ -BPS LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} = 0$). From Eq. (8.1.14), in order to get the solution (8.4.1), we have to fine-tune two PF BH charges out of four in the following way:

$$n_3 \approx n_2, \quad n_4 \approx n_2 - n_1. \quad (8.4.2)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition

$$N_1|_{N_2 \approx 0} \approx -\sqrt{3}n_2 + i(n_2 - 2n_1) \neq 0 \iff n_2 \neq 0, n_1 \neq \frac{n_2}{2}. \quad (8.4.3)$$

As it was for $k = 5$, also the real eigenvalues (8.3.4) for the $\frac{1}{2}$ -BPS critical solution coincide and, as it is well known [39, 15, 16], are strictly positive:

$$\lambda_{+, \frac{1}{2}-BPS} = \lambda_{-, \frac{1}{2}-BPS} \approx \frac{2}{3} \frac{C_1^2}{C_0^2} |N_1|_{N_2 \approx 0}^2 > 0. \quad (8.4.4)$$

Consequently, the $\frac{1}{2}$ -BPS LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (8.4.2) is a stable extremum, since it is a minimum of V_{BH} , and it is therefore an attractor in a strict sense. The classical (Bekenstein-Hawking) BH entropy at such a (class of) $\frac{1}{2}$ -BPS LG attractor(s) takes the value

$$S_{BH, \frac{1}{2}-BPS} = \pi V_{BH, \frac{1}{2}-BPS} \approx 0.032\pi |N_1|_{N_2 \approx 0}^2 = 0.128\pi (n_1^2 + n_2^2 - n_1 n_2). \quad (8.4.5)$$

II. The second non-degenerate solution to AE (8.2.1) is

$$N_1 \approx 0. \quad (8.4.6)$$

As one can see from Eq. (8.1.6)-(8.1.7), such a solution corresponds to a non-BPS, $Z = 0$ LG critical point of V_{BH} ($\widetilde{W} = 0$, $D\widetilde{W} \neq 0$). The real eigenvalues (8.3.4) for such a non-BPS, $Z = 0$ critical solution read

$$\lambda_{\pm, non-BPS, Z=0} \approx 2 \frac{C_1^2}{C_0^2} \left(1 \pm 9 \frac{C_0^2 C_3}{C_1^3} \right) |N_2|_{N_1 \approx 0}^2. \quad (8.4.7)$$

Substituting the numerical values (8.3.7) of the involved constants in Eq. (8.4.7), one reaches the conclusion that one eigenvalue is positive and the other one is negative:

$$\lambda_{\pm, non-BPS, Z=0} \approx 2 (0.45 \mp 1.46) |N_2|_{N_1 \approx 0}^2 \leq 0. \quad (8.4.8)$$

Let us now find the fine-tuning conditions for PF BH charges supporting the considered non-BPS, $Z = 0$ LG attractor for the mirror *sextic* $\mathcal{M}'_6$. This amounts to solving Eq. (8.4.6) by recalling Eq. (8.1.13). By doing so, one gets the following unique set of constraining relations on PF BH charges:

$$n_3 \approx -n_2, \quad n_4 \approx n_2 + n_1. \quad (8.4.9)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition

$$N_2|_{N_1 \approx 0} \approx -(2n_1 + n_2) + \sqrt{3}in_2 \neq 0 \iff n_2 \neq 0, n_1 \neq -\frac{n_2}{2}. \quad (8.4.10)$$

Thus, the non-BPS, $Z = 0$ LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (8.4.9) is a saddle point of V_{BH} and consequently it is *not* an attractor in a strict sense.

The classical (Bekenstein-Hawking) BH entropy at such a (class of) non-BPS, $Z = 0$ LG saddle point(s) takes the value

$$S_{BH, non-BPS, Z=0} = \pi V_{BH, non-BPS, Z=0} \approx 0.096\pi |N_2|_{N_1 \approx 0}^2 = 0.384\pi (n_1^2 + n_2^2 + n_1 n_2). \quad (8.4.11)$$

By using the definitions (6.32) and (6.33) for the case $k = 6$, one obtains (recall the first of definitions (6.28))

$$(n_1, n_2, n_3, n_4) = (p^0 - 3q_0, p^0 + 3q_1, -p^0 + 3p^1 + 9q_0 - 3q_1, -p^0 + 6q_0) \\ \Updownarrow \quad (8.4.12)$$

$$\Gamma = (-2n_1 - n_4, \frac{1}{3}(3n_1 - n_2 - n_3 + 3n_4), \frac{1}{3}(n_1 + n_4), \frac{1}{3}(-2n_1 + n_2 - n_4)).$$

Thus, the fine-tuning conditions (8.4.2) and (8.4.9) can be rewritten in terms of symplectic BH charges respectively as follows:

$$\frac{1}{2} - BPS : \begin{cases} -2p^0 + 3p^1 + 9q_0 - 6q_1 \approx 0, & 3q_0 \approx p^0 + 3q_1, \\ p^0 \neq -3q_1, p^0 \neq 6q_0 + 3q_1; \end{cases} \quad (8.4.13)$$

$$non - BPS, Z = 0 : \begin{cases} p^0 \approx 3q_0 - q_1, & p^1 \approx -3q_0, \\ p^0 \neq -3q_1, p^0 \neq 6q_0 - 3q_1. \end{cases} \quad (8.4.14)$$

Remark

The non-BPS, $Z = 0$ critical solution for the mirror *sextic* $\mathcal{M}'_6$ had been previously investigated in Sect. 7 of [10]. Up to irrelevant changes of notation, Eq. (8.4.14) coincides with Eq. (7.8) of [10]. By considering the second derivatives (8.1.11)-(8.1.12) of the “effective BH potential” constrained by Eq. (8.4.6) and comparing them with Eq. (7.9) of [10], one can state that the crucial difference between the results of [10] and ours lies in the critical value of the second holomorphic derivative of V_{BH} . Indeed, Eq. (7.9) of [10] reads

$$(\partial^2 V_{BH})_{non-BPS, Z=0} = 0. \quad (8.4.15)$$

From our previous computations, consistently taking into account the needed orders in ψ to get the series expansion for V_{BH} up to $\mathcal{O}(\psi^2)$ (or $\mathcal{O}(\psi^3)$), we disagree with the critical value of the second holomorphic derivative of V_{BH}

at the considered non-BPS, $Z = 0$ critical point given by Eq. (8.4.15). According to our results, the statement made in [10] that the considered non-BPS, $Z = 0$ LG critical point of V_{BH} is actually an attractor in a strict sense for all possible supporting symplectic BH charge configurations (8.4.14) does *not* hold. Instead, as correctly stated above, the non-BPS, $Z = 0$ LG critical point $\psi \approx 0$ supported by the BH charge configurations (8.4.9) (PF) and (8.4.14) (symplectic) is a saddle point of V_{BH} and consequently it is *not* an attractor in a strict sense.

Also, by recalling Eq. (8.4.11) and using Eqs. (8.4.10) and (8.4.12)-(8.4.14), our analysis yields that the “effective BH potential” and BH entropy at the considered (class of) non-BPS, $Z = 0$ LG saddle point(s) take the purely charge-dependent values:

$$V_{BH,non-BPS,Z=0} = \frac{1}{2\sqrt{3}}(3q_0^2 + q_1^2), \quad S_{BH,non-BPS,Z=0} = \frac{\pi}{2\sqrt{3}}(3q_0^2 + q_1^2). \quad (8.4.16)$$

As one can see, the value of $V_{BH,non-BPS,Z=0}$ given by Eq. (8.4.16) does *not* coincide with the one given by Eq. (7.10) of [10].

9 $k = 8$: Mirror *Octic* near LG Point

9.1 Geometric Setup

The case of mirror *octic* $\mathcal{M}'_8$ (as well as the one of mirror *dectic* $\mathcal{M}'_{10}$ treated in Sect. 10) needs a different approach with respect to the cases of mirror *quintic* $\mathcal{M}'_5$ and mirror *sextic* $\mathcal{M}'_6$, respectively treated in Sects. 7 and 8.

Indeed, contrary to what happens for $k = 5, 6$ (see Eqs. (7.1.4) and (8.1.4), respectively), for $k = 8, 10$ the series expansion of the Kähler metric $g(\psi, \bar{\psi})$ near the LG point starts with *no* constant term (namely, it is *not* regular at $\psi = 0$). As a consequence, one has to consider the series expansion of the “effective BH potential” V_{BH} up to $\mathcal{O}(\psi^4)$ (rather than up to $\mathcal{O}(\psi^3)$, as it is for $k = 5, 6$), in order to obtain all the relevant quantities up to $\mathcal{O}(\psi^2)$ (rather than up to $\mathcal{O}(\psi)$, as it is for $k = 5, 6$).

For $k = 8$ the definitions (6.12) yield

$$C_{8,2l-1} = 0, \quad l \in \mathbb{N}; \quad (9.1.1)$$

moreover, since $F_{8,m+8,n} = F_{8,m,n+8} = -F_{8,m,n}$ (see the third of properties (6.26)), the only independent elements of the rank-2 tensor F_8 belong to the 8×8 matrix

$$F_{8,mn} = \begin{pmatrix} 2(2+\sqrt{2}) & -\sqrt{10+7\sqrt{2}} & 0 & \sqrt{2+\sqrt{2}} & 0 & -\sqrt{2+\sqrt{2}} & 0 & \sqrt{10+7\sqrt{2}} \\ -\sqrt{10+7\sqrt{2}} & 2 & \sqrt{2-\sqrt{2}} & -\sqrt{2} & \sqrt{10-7\sqrt{2}} & 0 & \sqrt{2+\sqrt{2}} & -3\sqrt{2} \\ 0 & \sqrt{2-\sqrt{2}} & 2(-2+\sqrt{2}) & \sqrt{2-\sqrt{2}} & 0 & -\sqrt{10-7\sqrt{2}} & 0 & \sqrt{10-7\sqrt{2}} \\ \sqrt{2+\sqrt{2}} & -\sqrt{2} & \sqrt{2-\sqrt{2}} & 0 & -\sqrt{2-\sqrt{2}} & \sqrt{2} & -\sqrt{2+\sqrt{2}} & 2 \\ 0 & \sqrt{10-7\sqrt{2}} & 0 & -\sqrt{2-\sqrt{2}} & 4-2\sqrt{2} & -\sqrt{2-\sqrt{2}} & 0 & \sqrt{10-7\sqrt{2}} \\ -\sqrt{2+\sqrt{2}} & 0 & -\sqrt{10-7\sqrt{2}} & \sqrt{2} & -\sqrt{2-\sqrt{2}} & -2 & \sqrt{10+7\sqrt{2}} & -3\sqrt{2} \\ 0 & \sqrt{2+\sqrt{2}} & 0 & -\sqrt{2+\sqrt{2}} & 0 & \sqrt{10+7\sqrt{2}} & -4-2\sqrt{2} & \sqrt{10+7\sqrt{2}} \\ \sqrt{10+7\sqrt{2}} & -3\sqrt{2} & \sqrt{10-7\sqrt{2}} & 2 & \sqrt{10-7\sqrt{2}} & -3\sqrt{2} & \sqrt{10+7\sqrt{2}} & 0 \end{pmatrix}. \quad (9.1.2)$$

Let us now write down all the relevant quantities up to the needed order (here and below, unless otherwise specified, we omit the Fermat parameter $k = 8$):

$$\tilde{K} \approx (3 - 2\sqrt{2}) \frac{C_2^2}{C_0^2} (\psi\bar{\psi})^2 \left[1 - \left(\frac{C_4^2}{C_2^2} - \frac{1}{2}(3 - 2\sqrt{2}) \frac{C_2^2}{C_0^2} \right) (\psi\bar{\psi})^2 \right] + \frac{C_8}{C_0} (\psi^8 + \bar{\psi}^8) + \mathcal{O}(\psi^9); \quad (9.1.3)$$

$$g \approx 4(3 - 2\sqrt{2}) \frac{C_2^2}{C_0^2} \psi\bar{\psi} \left[1 - 4 \left(\frac{C_4^2}{C_2^2} - \frac{1}{2}(3 - 2\sqrt{2}) \frac{C_2^2}{C_0^2} \right) (\psi\bar{\psi})^2 \right] + \mathcal{O}(\psi^7); \quad (9.1.4)$$

$$\widetilde{W} \approx N_1 + \frac{C_2}{C_0} N_3 \psi^2 - \frac{C_4}{C_0} \bar{N}_3 \psi^4 - \frac{C_6}{C_0} \bar{N}_1 \psi^6 + \mathcal{O}(\psi^8). \quad (9.1.5)$$

Now, by using the formulæ of the general analysis exploited in Sect. 6, we can get the “effective BH potential” and the holomorphic superpotential, as well as their (covariant) derivatives, up to $\mathcal{O}(\psi^2)$:

$$\widetilde{W} = N_1 + \frac{C_2}{C_0} N_3 \psi^2; \quad (9.1.6)$$

$$D\widetilde{W} = 2\frac{C_2}{C_0} N_3 \psi - 4\frac{C_4}{C_0} \bar{N}_3 \psi^3 + 2(3 - 2\sqrt{2}) \frac{C_2^2}{C_0^2} N_1 \psi \bar{\psi}^2; \quad (9.1.7)$$

$$D^2\widetilde{W} = -8\frac{C_4}{C_0} \bar{N}_3 \psi^2; \quad (9.1.8)$$

$$\begin{aligned} \tilde{V}_{BH} = & |N_1|^2 + (3 + 2\sqrt{2}) |N_3|^2 + 2\frac{C_2}{C_0} \left(N_3 \bar{N}_1 - (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} (\bar{N}_3)^2 \right) \psi^2 + \\ & + 2\frac{C_2}{C_0} \left(\bar{N}_3 N_1 - (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} (N_3)^2 \right) \bar{\psi}^2; \end{aligned} \quad (9.1.9)$$

$$\begin{aligned} \partial\tilde{V}_{BH} = & 4\psi \left[\frac{C_2}{C_0} \left(N_3 \bar{N}_1 - (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} (\bar{N}_3)^2 \right) - 3\frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) \bar{N}_1 \bar{N}_3 \psi^2 + \right. \\ & \left. + \frac{C_2^2}{C_0^2} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^2} \right) \right) \bar{\psi}^2 \right]; \end{aligned} \quad (9.1.10)$$

$$\begin{aligned} \partial^2\tilde{V}_{BH} = & 4\frac{C_2}{C_0} \left(N_3 \bar{N}_1 - (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} (\bar{N}_3)^2 \right) - 36\frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) \bar{N}_1 \bar{N}_3 \psi^2 + \\ & + 4\frac{C_2^2}{C_0^2} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^2} \right) \right) \bar{\psi}^2; \end{aligned} \quad (9.1.11)$$

$$\partial\bar{\partial}\tilde{V}_{BH} = 8\frac{C_2^2}{C_0^2} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^2} \right) \right) \psi\bar{\psi}. \quad (9.1.12)$$

Let us stress once again that, contrary to the treatment of Sects. 7 and 8, and as evident from Eqs. (9.1.9)-(9.1.12), for the case of mirror *octic* we truncate the series expansion of the “effective BH potential” and of its second derivatives around the LG point up to $\mathcal{O}(\psi^2)$ included, and the series expansion of its first derivative around the LG point up to $\mathcal{O}(\psi^3)$ included. This is due to the absence of an $\mathcal{O}(\psi)$ term in expression of $\tilde{V}_{BH}$ given by Eq. (9.1.9). As mentioned at the start of the present Section, such a fact can be traced back to the non-regularity of g at $\psi = 0$ (see Eq. (9.1.4)).

9.1.1 The Holomorphic Prepotentials F and $\mathcal{F}$

As done for $k = 5, 6$ respectively in Subsubsects. 7.1.1 and 8.1.1, in order to compute the holomorphic prepotential $F(X(\psi))$ and its “Kähler-invariant counterpart” $\mathcal{F}(\psi)$ defined by Eq. (4.1.9), we have to recall the general formulæ for (mirror) Fermat CY_3 s introduced in Sect. 6, and specialize them for $k = 8$. By recalling the definitions (6.38), (6.40), (6.41), (6.42), one can compute:

$$\xi_{8,m} = \begin{pmatrix} -i \left[\sin\left(\frac{\pi m}{8}\right) + \sin\left(\frac{3\pi m}{8}\right) \right] \\ -\cos\left(\frac{\pi m}{8}\right) + i \sin\left(\frac{\pi m}{8}\right) \\ 4\cos^2\left(\frac{\pi m}{8}\right) \left[\cos\left(\frac{\pi m}{8}\right) - 3i \sin\left(\frac{\pi m}{8}\right) \right] \\ 2i \sin\left(\frac{\pi m}{8}\right) \end{pmatrix}; \quad (9.1.1.1)$$

$$\begin{aligned} \Phi_{8,m,n} = \\ = \frac{1}{2} \left\{ \begin{aligned} & \left[-3\cos\left(\frac{\pi(n-3m)}{8}\right) - 3\cos\left(\frac{\pi(3n-m)}{8}\right) - 5\cos\left(\frac{\pi(n-m)}{8}\right) - 3\cos\left(\frac{3\pi(n-m)}{8}\right) + \right. \\ & \left. + 5\cos\left(\frac{\pi(n+m)}{8}\right) + 3\cos\left(\frac{3\pi(n+m)}{8}\right) + 3\cos\left(\frac{\pi(3n+m)}{8}\right) + 3\cos\left(\frac{\pi(n+3m)}{8}\right) \right] + \\ & + i \left[3\sin\left(\frac{\pi(n-3m)}{8}\right) + \sin\left(\frac{\pi(3n-m)}{8}\right) + \sin\left(\frac{\pi(n-m)}{8}\right) + \sin\left(\frac{3\pi(n-m)}{8}\right) + \right. \\ & \left. - 5\sin\left(\frac{\pi(n+m)}{8}\right) - \sin\left(\frac{3\pi(n+m)}{8}\right) - \sin\left(\frac{\pi(3n+m)}{8}\right) - 3\sin\left(\frac{\pi(n+3m)}{8}\right) \right] \end{aligned} \right\} \neq \Phi_{8,(m,n)}; \end{aligned} \quad (9.1.1.2)$$

$$F_{k=8,l=7}(X_8(\psi)) = -\frac{\pi^6}{2^9} \sum_{m=1}^7 \sum_{n=1}^7 (-1)^{m+n} C_{8,m-1} C_{8,n-1} \psi^{m+n-2} \Phi_{8,(m,n)} =$$

$$\begin{aligned} = -\frac{\pi^6}{2^9} \left[\begin{aligned} & -\frac{(4+\sqrt{2}+2i(1+\sqrt{2}))\pi}{\Gamma^2\left(\frac{5}{6}\right)\Gamma^6\left(\frac{7}{8}\right)} + \frac{8(i+\sqrt{2})\sqrt{\pi}}{\Gamma^3\left(\frac{5}{8}\right)\Gamma\left(\frac{5}{6}\right)\Gamma^3\left(\frac{7}{8}\right)} \psi^2 + \\ & + \left[\frac{-16+4\sqrt{2}+8i(1-\sqrt{2})}{\Gamma^6\left(\frac{5}{8}\right)} - \frac{4(2+i)2^{3/4}}{\pi^{3/2}\Gamma^3\left(\frac{3}{4}\right)} \right] \psi^4 + \\ & + \frac{1}{\sqrt{2}\pi^4} \left[\frac{(4-\sqrt{2})\sqrt{\pi}\Gamma\left(\frac{5}{6}\right)}{\sin^3\left(\frac{7\pi}{8}\right)} + \frac{2(4+\sqrt{2})\Gamma^2\left(\frac{7}{6}\right)}{\sin^3\left(\frac{5\pi}{8}\right)} \right] \psi^6 \end{aligned} \right] + \mathcal{O}(\psi^7) = F_{k=8,l=7}(\psi^2); \end{aligned} \quad (9.1.1.3)$$

$$\begin{aligned}
\mathcal{F}_8(\psi) &\equiv F_8\left(\frac{X_8^1(\psi)}{X_8^0(\psi)}\right) = \frac{F_8(X_8^0(\psi), X_8^1(\psi))}{(X_8^0(\psi))^2} = \frac{1}{2} \frac{\sum_{m,n=1}^{\infty} (-1)^{m+n} C_{8,m-1} C_{8,n-1} \Phi_{8,(m,n)} \psi^{m+n-2}}{\left[\sum_{r=1}^{\infty} (-1)^r C_{8,r-1} \xi_{8,r}^1 \psi^{r-1}\right]^2} \approx \\
&\approx \frac{\Phi_{8,1,1}}{2(\xi_{8,1}^1)^2} \left[1 + \frac{C_{8,2}(\Phi_{8,1,3} + \Phi_{8,3,1})}{C_{8,0} \Phi_{8,1,1}} \psi^2 + \mathcal{O}(\psi^3)\right] \left[1 - 2 \frac{C_{8,2} \xi_{8,3}^1}{C_{8,0} \xi_{8,1}^1} \psi^2 + \mathcal{O}(\psi^3)\right] \approx \\
&\approx \frac{\Phi_{8,1,1}}{2(\xi_{8,1}^1)^2} \left[1 - 2 \left(\frac{\xi_{8,3}^1}{\xi_{8,1}^1} - \frac{\Phi_{8,1,3} + \Phi_{8,3,1}}{2\Phi_{8,1,1}}\right) \frac{C_{8,2}}{C_{8,0}} \psi^2\right] + \mathcal{O}(\psi^3),
\end{aligned} \tag{9.1.1.4}$$

where in the formula (9.1.1.4) for the Kähler-invariant holomorphic prepotential $\mathcal{F}_8$ only the leading terms in the *LG limit* $\psi \rightarrow 0$ have been retained. Eq. (9.1.1.3) is the exact expression (complete up to $\mathcal{O}(\psi^6)$ included) of the holomorphic prepotential F_8 in a suitable neighbourhood of the LG point of the moduli space of the (mirror) *octic* $\mathcal{M}'_8$.

9.2 “Criticality Condition” Approach and Attractor Equation

Let us now find the solutions of the AE $\partial \tilde{V}_{BH}(\psi, \bar{\psi}; q, p) = 0$, and check their stability. Since we are working near the LG point, by using Eq. (9.1.9) we can rewrite the AE for $\mathcal{M}'_8$ as follows:

$$\begin{aligned}
&\frac{C_2}{C_0} \left(N_3 \bar{N}_1 - (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} (\bar{N}_3)^2 \right) + \frac{C_2^2}{C_0^2} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^4} \right) \right) \bar{\psi}^2 \approx \\
&\approx 3 \frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) \bar{N}_1 \bar{N}_3 \psi^2.
\end{aligned} \tag{9.2.1}$$

Solving Eq. (9.2.1), we will find one (or more) set(s) of BH charges supporting $\psi \approx 0$ to be a critical point of V_{BH} . Since we are working near the LG point, it is clear that the first term in the left-hand side (l.h.s.) of Eq. (9.2.1) must be small enough. This implies the following fine-tuning condition:

$$\begin{aligned}
N_3 \bar{N}_1 - \vartheta (\bar{N}_3)^2 &\approx 0, \\
\vartheta &\equiv (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2}.
\end{aligned} \tag{9.2.2}$$

9.3 Critical Hessian of V_{BH}

By using (the $n_V = 1$ case of) Eq. (3.1.11) and Eqs. (9.1.11)-(9.1.12) evaluated along the criticality condition (9.2.1)-(9.2.2), it can be computed that the components of $H_{\text{real form}}^{\tilde{V}_{BH}}$ (given by Eq. (7.3.1)) constrained by Eqs. (9.2.1)-(9.2.2)

read as follows:

$$\begin{aligned} \mathcal{A} = & -12 \frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) (\bar{N}_1 \bar{N}_3 \psi^2 + N_1 N_3 \bar{\psi}^2) + \\ & + 8 \frac{C_2^2}{C_0^2} \psi \bar{\psi} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^4} \right) \right); \end{aligned} \quad (9.3.1)$$

$$\begin{aligned} \mathcal{B} = & 12 \frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) (\bar{N}_1 \bar{N}_3 \psi^2 + N_1 N_3 \bar{\psi}^2) + \\ & + 8 \frac{C_2^2}{C_0^2} \psi \bar{\psi} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^4} \right) \right); \end{aligned} \quad (9.3.2)$$

$$\mathcal{C} = -12i \frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) (N_1 N_3 \bar{\psi}^2 - \bar{N}_1 \bar{N}_3 \psi^2). \quad (9.3.3)$$

The resulting real eigenvalues of $H_{\text{real form}}^{\tilde{V}_{BH}}$ constrained by Eqs. (9.2.1)-(9.2.2) read:

$$\lambda_{\pm} \approx 8\psi \bar{\psi} \left[\begin{aligned} & \frac{C_2^2}{C_0^2} \left(|N_1|^2 (3 - 2\sqrt{2}) + |N_3|^2 \left(1 + 4(3 + 2\sqrt{2}) \frac{C_0^2 C_4^2}{C_2^4} \right) \right) + \\ & \pm 3 \frac{C_4}{C_0} \left(1 + (3 + 2\sqrt{2}) \frac{C_6 C_0}{C_4 C_2} \right) |N_1| |N_3| \end{aligned} \right]. \quad (9.3.4)$$

By recalling Eq. (6.34) and using Eq. (9.1.9) with $\psi \approx 0$ and constrained by Eqs. (9.2.1)-(9.2.2), one obtains that the purely charge-dependent LG critical values of the “effective BH potential” for the mirror *octic* $\mathcal{M}'_8$ are

$$V_{BH, LG-critical, k=8} \approx \frac{1}{8(2 + \sqrt{2})} \left[|N_1|^2 + (3 + 2\sqrt{2}) |N_3|^2 \right]; \quad (9.3.5)$$

by recalling formula (1.3), this directly yields the following purely charge-dependent values of the BH entropy at the LG critical points of $V_{BH,8}$ in the moduli space of $\mathcal{M}'_8$:

$$S_{BH, LG-critical, k=8} \approx \frac{\pi}{8(2 + \sqrt{2})} \left[|N_1|^2 + (3 + 2\sqrt{2}) |N_3|^2 \right]. \quad (9.3.6)$$

Let us write down here the numerical values of constants relevant to our treatment:

$$C_0 \approx 1.64, \quad C_2 \approx -0.9, \quad C_4 \approx 0.24, \quad C_6 \approx -0.007, \quad C_8 \approx -0.004; \quad \vartheta \approx 2.83. \quad (9.3.7)$$

9.4 Solutions to Attractor Equations

Let us now analyze more in depth the species of LG attractor points arising from the AE (9.2.2). As it can be easily seen, the AE (9.2.2) has two *non-degenerate* solutions:

I. The first non-degenerate solution to AE (9.2.2) is

$$N_3 \approx 0. \quad (9.4.1)$$

As one can see from Eq. (9.1.6)-(9.1.7), such a solution corresponds to a $\frac{1}{2}$ -BPS LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} = 0$). From the definition (6.31) with $(k, m) = (8, 3)$, in order to get the solution (9.4.1), one has to fine-tune two PF BH charges out of four in the following way:

$$n_3 \approx -n_1 + \sqrt{2}n_2, \quad n_4 \approx -\sqrt{2}n_1 + n_2, \quad (9.4.2)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition $N_1 \neq 0$. The real eigenvalues (9.3.4) for the $\frac{1}{2}$ -BPS critical solution coincide and, as it is well known [39, 15, 16], are strictly positive:

$$\lambda_{+, \frac{1}{2}-BPS} = \lambda_{-, \frac{1}{2}-BPS} \approx 8\psi\bar{\psi} \frac{C_2^2}{C_0^2} (3 - 2\sqrt{2}) |N_1|_{N_3 \approx 0}^2 > 0.$$

Consequently, the $\frac{1}{2}$ -BPS LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (9.4.2) is a stable extremum, since it is a (local) minimum of V_{BH} , and it is therefore an attractor in a strict sense. The classical (Bekenstein-Hawking) BH entropy at such a (class of) $\frac{1}{2}$ -BPS LG attractor(s) takes the value

$$S_{BH, \frac{1}{2}-BPS} = \pi V_{BH, \frac{1}{2}-BPS} \approx 0.0366\pi |N_1|_{N_3 \approx 0}^2, \quad (9.4.3)$$

where $|N_1|_{N_3 \approx 0}$ is given by the general formula (6.31) with $(k, m) = (8, 1)$ constrained by Eq. (9.4.2):

$$|N_1|_{N_3 \approx 0} \approx \sqrt{4 - 2\sqrt{2}[n_1(1 - i(1 + \sqrt{2})) - n_2(1 + \sqrt{2} - i)]}. \quad (9.4.4)$$

II. The second non-degenerate solution to AE (9.2.2) reads (from Eqs. (9.2.2) and (9.3.7): $\vartheta \equiv (3 + 2\sqrt{2}) \frac{C_4 C_0}{C_2^2} \approx 2.83$):

$$|N_1| \approx \vartheta |N_3|, \quad (9.4.5)$$

$$\arg(N_1) \approx 3\arg(N_3),$$

where N_1 and N_3 are given by the general formula (6.31) with $k = 8$ and $m = 1, 3$, respectively:

$$N_1 = -\frac{\sqrt{2 - \sqrt{2}}}{2} \left(n_1 + n_4 + (1 + \sqrt{2})(n_2 + n_3) - i((n_4 - n_1)(1 + \sqrt{2}) + n_3 - n_2) \right); \quad (9.4.6)$$

$$N_3 = \frac{\sqrt{2 + \sqrt{2}}}{2} \left(n_1 + n_4 + (1 - \sqrt{2})(n_2 + n_3) + i((n_4 - n_1)(1 - \sqrt{2}) + n_3 - n_2) \right). \quad (9.4.7)$$

The first of fine-tuning conditions (9.4.5) substituted in Eq. (9.3.6) yields

$$S_{BH, non-BPS, Z \neq 0} \approx \frac{\pi}{8(2 + \sqrt{2})} \left(\vartheta^2 + 3 + 2\sqrt{2} \right) |N_3|^2 \approx 0.5066\pi |N_3|^2. \quad (9.4.8)$$

As one can see from Eq. (9.1.6)-(9.1.7), such a solution corresponds to a non-BPS, $Z \neq 0$ LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} \neq 0$). The real eigenvalues (9.3.4) for such a non-BPS, $Z \neq 0$ critical solution read

$$\begin{aligned} \lambda_{\pm, non-BPS, Z \neq 0} &\approx 8\psi\bar{\psi} |N_3|^2 \frac{C_2^2}{C_0^2} \left[(1 + 5(3 - 2\sqrt{2})\vartheta^2) \pm 3(3 - 2\sqrt{2})\vartheta^2 \left(1 + (3 + 2\sqrt{2}) \frac{C_0 C_6}{C_2 C_4} \right) \right] \approx \\ &\approx 8\psi\bar{\psi} |N_3|^2 \frac{C_2^2}{C_0^2} [7.9 \pm 5.5] > 0, \end{aligned} \quad (9.4.9)$$

where in the second line we replaced some constants with their numerical values by using Eq. (9.3.7).

Thus, the non-BPS, $Z \neq 0$ LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (9.4.5)-(9.4.7) is a (local) minimum of V_{BH} and consequently an attractor in a strict sense.

Let us now find the fine-tuning conditions for PF BH charges supporting the considered non-BPS, $Z \neq 0$ LG attractor for the mirror *octic* $\mathcal{M}'_8$. This amounts to solving Eqs. (9.4.5)-(9.4.7) by recalling the definitions (6.31) and (6.32). By doing so, one obtains the following three different sets of constraining relations on PF BH charges:

$$\text{II.1 : } \left\{ \begin{array}{l} n_{2,\pm} = (a_{1,\pm} + a_2)n_1 \approx \begin{cases} \text{"+"} : 2.4581n_1 \\ \text{"-"} : -1.32378n_1 \end{cases} , \\ n_{3,\pm} = (a_{1,\pm} - a_2)n_1 = -n_{2,\mp}, \\ n_4 = -n_1, \\ a_{1,\pm}(\vartheta) \equiv \pm \frac{\sqrt{2}}{\vartheta+1} \sqrt{\frac{3(2-\sqrt{2})\vartheta+\sqrt{2}}{(2-\sqrt{2})\vartheta-\sqrt{2}}} \approx \pm 1.89094, \\ a_2(\vartheta) \equiv \frac{\sqrt{2}+\vartheta(2-\sqrt{2})}{\sqrt{2}(1+\vartheta)} \approx 0.56716, \\ n_1 \neq 0; \end{array} \right. \quad (9.4.10)$$

$$\text{II.2 : } \left\{ \begin{array}{l} n_2 = \frac{\sqrt{2}-1+\vartheta}{-1+(\sqrt{2}-1)\vartheta} n_1 \approx 18.83713n_1, \\ n_3 = n_2, \\ n_4 = n_1, \\ n_1 \neq 0; \end{array} \right. \quad (9.4.11)$$

$$\text{II.3 : } \left\{ \begin{array}{l} \left\{ \begin{array}{l} n_{2,\pm} + n_{3,\pm} = a_{\pm}, \\ n_{1,\pm} + n_{4,\pm} = b_{\pm}, \\ n_{2,\pm} - n_{3,\pm} = c, \\ n_{1,\pm} - n_{4,\pm} = d, \end{array} \right\} \iff \left\{ \begin{array}{l} n_{1,\pm} = \frac{b_{\pm}+d}{2}, \\ n_{2,\pm} = \frac{a_{\pm}+c}{2}, \\ n_{3,\pm} = \frac{a_{\pm}-c}{2}, \\ n_{4,\pm} = \frac{b_{\pm}-d}{2}, \end{array} \right. \\ a_{\pm}(\vartheta; b_{\pm}, c, d) \equiv -\frac{c+d-(\sqrt{2}+1)[c-(\sqrt{2}-1)d]\vartheta}{c-d+[c-(\sqrt{2}-1)d]\vartheta} b_{\pm}(\vartheta; c, d) \approx \left(1.76316 + \frac{c}{-4.1598c+d}\right) b_{\pm}(\vartheta; c, d) \approx (9.4.12) \\ \approx \pm 2.29892 \sqrt{\frac{c-0.053086d}{c-0.55928d} \frac{(c-0.31647d)(c-0.56716d)}{c-0.24039d}}, \\ b_{\pm}(\vartheta; c, d) \equiv \pm (1+\vartheta) \sqrt{\frac{1+(\sqrt{2}+1)\vartheta}{2(-1+3(\sqrt{2}+1)\vartheta)}} \left[c - \frac{1+(\sqrt{2}-1)\vartheta}{1+\vartheta} d \right] \sqrt{\frac{c-(\sqrt{2}+1)\frac{-1+(\sqrt{2}-1)\vartheta}{1+(\sqrt{2}+1)\vartheta}d}{c-(\sqrt{2}+1)\frac{-1-3(\sqrt{2}-1)\vartheta}{1-3(\sqrt{2}+1)\vartheta}d}} \approx \\ \approx \pm 1.71651(c-0.56716d) \sqrt{\frac{c-0.053086d}{c-0.55928d}}. \end{array} \right.$$

By using the definitions (6.32) and (6.33) for the case $k = 8$, one obtains (recall the first of definitions (6.28))

$$\begin{aligned} (n_1, n_2, n_3, n_4) &= (p^0 - 2q_0, p^0 + 2q_1, -p^0 + 2p^1 + 6q_0 - 2q_1, -p^0 + 4q_0) \\ &\Updownarrow \\ \Gamma &= (-2n_1 - n_4, \frac{1}{2}(3n_1 - n_2 - n_3 + 3n_4), \frac{1}{2}(n_1 + n_4), \frac{1}{2}(-2n_1 + n_2 - n_4)). \end{aligned} \quad (9.4.13)$$

9.4.1 Analysis of non-BPS, $Z \neq 0$ supporting BH charge configurations

Let us analyze more in depth the obtained supporting BH charge configurations.

II.1. As for $k = 5$, this configuration is characterized by a physically meaningful “ $\pm$ ” split, affecting the charge-dependent expression of the BH entropy. By using Eqs. (9.4.7), (9.4.8) and (9.4.13), Eqs. (9.4.10) yield

$$\begin{aligned} N_{3,II.1,\pm} &= \frac{\sqrt{2}\sqrt{2-\sqrt{2}}}{\vartheta+1} \left[\pm \sqrt{\frac{3(\sqrt{2}-1)\vartheta+1}{(\sqrt{2}-1)\vartheta-1}} + i \right] n_1 \approx 0.2826 (\pm 5.12109 + i) n_1; \\ S_{BH,non-BPS,Z \neq 0,II.1,\pm} &\approx 0.5066\pi |N_{3,II.1,\pm}|^2 \approx \begin{cases} \text{“+”} : 1.10157\pi (p^0 - 2q_0)^2. \\ \text{“-”} : -1.02065\pi (p^0 - 2q_0)^2. \end{cases} \end{aligned} \quad (9.4.1.1)$$

In order to give an example of the range of the symplectic (electric/magnetic) charges in the considered BH charge configuration, by recalling that PF BH charges are integers as are the symplectic BH charges (see definitions (6.32)-(6.33)), one can fulfill the fine-tuning conditions (9.4.10) by taking *e.g.*

$$\begin{aligned} n_1 &= 10 \\ &\Downarrow \\ n_{2,\pm} &\approx \begin{cases} \text{“+”} : 25 \\ \text{“-”} : -13 \end{cases}, n_{3,\pm} \approx \begin{cases} \text{“+”} : 13 \\ \text{“-”} : -25 \end{cases}, n_4 = -10. \end{aligned} \quad (9.4.1.2)$$

Using Eq. (9.4.13), one finally gets

$$\Gamma_{\pm} \approx \begin{cases} \text{“+”} : (-10, -19, 0, 7) \\ \text{“-”} : (-10, 19, 0, -11) \end{cases}. \quad (9.4.1.3)$$

II.2. As for $k = 5$, this configuration is the only one not having any kind of split, and having a purely real N_3 . It is worth remarking that Eq. (9.4.11) is the analogue for the mirror *octic* $\mathcal{M}'_8$ of the fine-tuning condition (7.4.13) adopted for the mirror *quintic* $\mathcal{M}'_5$ in [9] (in particular, see Sect. 4 and App. C of such a Ref.). By using Eqs. (9.4.7), (9.4.8) and (9.4.13), Eqs. (9.4.11) yield

$$\begin{aligned} N_{3,II.2} &= \frac{2\sqrt{2}\sqrt{2-\sqrt{2}}}{-1+(\sqrt{2}-1)\vartheta} n_1 \approx 12.56955 (p^0 - 2q_0); \\ S_{BH,non-BPS,Z \neq 0,II.2} &\approx 0.5066\pi |N_{3,II.2}|^2 \approx 80.03965\pi (p^0 - 2q_0)^2. \end{aligned} \quad (9.4.1.4)$$

In order to give an example of the range of the symplectic (electric/magnetic) charges in the considered BH charge configuration, by recalling that PF BH charges are integers as are the symplectic BH charges (see definitions (6.32)-(6.33)), one can fulfill the fine-tuning conditions (9.4.11) by taking *e.g.*

$$\begin{aligned} n_1 &= 10 \\ \Downarrow \\ n_2 &\approx 188, \quad n_3 \approx 188, \quad n_4 = 10. \end{aligned} \tag{9.4.1.5}$$

Using Eq. (9.4.13), one finally gets

$$\Gamma = (-30, -158, 10, 79). \tag{9.4.1.6}$$

II.3. As for $k = 5$, this case, beside being split in “ $\pm$ ” (as the case II.1), is also parameterized by the couple $(c, d) \in \mathbb{Z}^2$ ($\mathbb{R}^2$ in the -semi- classical limit of large charges), further constrained by the request $a_{\pm}(b_{\pm}, c, d), b_{\pm}(c, d) \in \mathbb{Z}$ ($\mathbb{R}$ for large charges), implying that

$$c \neq 0.24039d, \quad \text{and} \quad \left\{ \begin{array}{l} \left\{ \begin{array}{l} c \geq 0.053086d, \\ c > 0.55928d; \end{array} \right. \\ \text{or} \\ \left\{ \begin{array}{l} c \leq 0.053086d, \\ c < 0.55928d. \end{array} \right. \end{array} \right. \tag{9.4.1.7}$$

By using Eq. (9.4.7), Eqs. (9.4.12) yield

$$\begin{aligned} N_{3,II.3,\pm}(c, d) &= \frac{\sqrt{2+\sqrt{2}}}{2} \left\{ [(1-\sqrt{2})a_{\pm}(c, d) + b_{\pm}(c, d)] - i[c + (1-\sqrt{2})d] \right\} = \\ &= \frac{\sqrt{2+\sqrt{2}}}{2} \left\{ \frac{\sqrt{2}c - \sqrt{2}(\sqrt{2}-1)d}{c-d+[c-(\sqrt{2}-1)d]^{\vartheta}} b_{\pm}(c, d) - i[c + (1-\sqrt{2})d] \right\} = \\ &= \frac{\sqrt{2+\sqrt{2}}}{2} \left\{ \pm \sqrt{\frac{1+(\sqrt{2}+1)^{\vartheta}}{-1+3(\sqrt{2}+1)^{\vartheta}}} \sqrt{\frac{c-(\sqrt{2}+1)^{\frac{-1+(\sqrt{2}-1)^{\vartheta}}{1+(\sqrt{2}+1)^{\vartheta}}d}{c-(\sqrt{2}+1)^{\frac{-1-3(\sqrt{2}-1)^{\vartheta}}{1-3(\sqrt{2}+1)^{\vartheta}}d}} - i} \right\} [c + (1-\sqrt{2})d] \approx \\ &\approx 0.92388 \left\{ \pm 0.63381 \sqrt{\frac{c-0.053086d}{c-0.55928d}} - i \right\} (c - 0.41421d); \end{aligned} \tag{9.4.1.8}$$

$$\begin{aligned} |N_{3,II.3,\pm}(c, d)|^2 &= \\ &= \frac{2+\sqrt{2}}{4} \left\{ \frac{[1+(\sqrt{2}+1)^{\vartheta}]c + (\sqrt{2}+1)[1-(\sqrt{2}-1)^{\vartheta}]d}{[-1+3(\sqrt{2}+1)^{\vartheta}]c + (\sqrt{2}+1)[-1-3(\sqrt{2}-1)^{\vartheta}]d} + 1 \right\} [c + (1-\sqrt{2})d]^2 \approx \\ &\approx 1.19644 \frac{(c-0.41421d)^3}{c-0.55928d}. \end{aligned} \tag{9.4.1.9}$$

By recalling Eqs. (9.4.8) and (9.4.13), one finally gets (notice that in this case the “ $\pm$ ” split is unphysical, because it does not affect the charge-dependent expression of the BH entropy) :

$$\begin{aligned}
S_{BH, non-BPS, Z \neq 0, II.3, \pm}(c, d) &\approx 0.5066\pi |N_{2, II.3, \pm}(c, d)|^2 \approx 0.60611\pi \frac{(c-0.41421d)^3}{c-0.55928d} \approx \\
&\approx 0.60611\pi \frac{(2p^0-2p^1-3.51474q_0+4q_1)^3}{0.88144p^0-2p^1-2.64432q_0+4q_1} = \\
&= S_{BH, non-BPS, Z \neq 0, II.3, \pm}(p^0, p^1, q_0, q_1).
\end{aligned} \tag{9.4.1.10}$$

Thus, by comparing Eqs. (9.4.1.1), (9.4.1.4) and (9.4.1.10), also for $k = 8$ one can conclude that *all* three distinct sets of fine-tuning conditions for PF BH charges (9.4.10)-(9.4.12) do support a non-BPS, $Z \neq 0$ LG attractor in a strict sense, but they yield different charge-dependent expressions for the classical BH entropy $S_{BH, non-BPS, Z \neq 0}$. As non-BPS, $Z \neq 0$ attractor, the LG point of the moduli space of the Fermat mirror *octic* $\mathcal{M}'_8$ turns out to originate a *threefold splitting* of the supporting BH charges, and correspondingly of the purely charge-dependent classical (Bekenstein-Hawking) BH entropy. Once again, this confirms the results of the recent investigation in [35].

Finally, by recalling the definition (2.2.11), one can compute the supersymmetry-breaking order parameter for the non-BPS, $Z \neq 0$ LG attractor in the mirror *octic* $\mathcal{M}'_8$; by using Eqs. (9.1.4), (9.1.6), (9.1.7) and (9.4.5), one gets

$$\mathcal{O}_{non-BPS, Z \neq 0} \equiv \left[\frac{g^{-1}|DW|^2}{|W|^2} \right]_{non-BPS, Z \neq 0} = \left(g^{-1}|D \ln \widetilde{W}|^2 \right)_{non-BPS, Z \neq 0} \approx \frac{3+2\sqrt{2}}{\vartheta^2} \approx 0.72. \tag{9.4.1.11}$$

10 $k = 10$: Mirror *Dectic* near LG Point

10.1 Geometric Setup

For the mirror *dectic* $\mathcal{M}'_{10}$ the computations (*but not the results!*) go the same way as for the mirror *octic* $\mathcal{M}'_8$.

For $k = 10$ the definitions (6.12) yield

$$C_{10, 2l-1} = 0 = C_{10, 5l-1}, \quad l \in \mathbb{N}; \tag{10.1.1}$$

moreover, since $F_{10, m+10, n} = F_{10, m, n+10} = -F_{10, m, n}$ (see the third of properties (6.26)), the only independent elements of the rank-2 tensor F_{10} belong to the 10×10 matrix

$$F_{10,m\,n} = \quad (10.1.2)$$

$$\begin{pmatrix} \sqrt{5(5+2\sqrt{5})} & -3-\sqrt{5} & 0 & 3+\sqrt{5} & -\sqrt{5(5+2\sqrt{5})} & 2+\sqrt{5} & 0 & -2 & 0 & 2+\sqrt{5} \\ -3-\sqrt{5} & \sqrt{5+2\sqrt{5}} & 3-\sqrt{5} & -\sqrt{2(5+\sqrt{5})} & 2+\sqrt{5} & -\sqrt{10-2\sqrt{5}} & -2+\sqrt{5} & 0 & 2 & -\sqrt{25-2\sqrt{5}} \\ 0 & 3-\sqrt{5} & -\sqrt{5(5-2\sqrt{5})} & 2 & -\sqrt{5(5-2\sqrt{5})} & 3-\sqrt{5} & 0 & 2-\sqrt{5} & 0 & -2+\sqrt{5} \\ 3+\sqrt{5} & -\sqrt{2(5+\sqrt{5})} & 2 & -\sqrt{5-2\sqrt{5}} & -2+\sqrt{5} & 0 & -3+\sqrt{5} & \sqrt{10-2\sqrt{5}} & -2-\sqrt{5} & \sqrt{25+2\sqrt{5}} \\ -\sqrt{5(5+2\sqrt{5})} & 2+\sqrt{5} & -\sqrt{5(5-2\sqrt{5})} & -2+\sqrt{5} & 0 & 2-\sqrt{5} & \sqrt{5(5-2\sqrt{5})} & -2-\sqrt{5} & \sqrt{5(5+2\sqrt{5})} & 8 \\ 2+\sqrt{5} & -\sqrt{10-2\sqrt{5}} & 3-\sqrt{5} & 0 & 2-\sqrt{5} & \sqrt{5-2\sqrt{5}} & -2 & \sqrt{2(5+\sqrt{5})} & -3-\sqrt{5} & \sqrt{25+2\sqrt{5}} \\ 0 & -2+\sqrt{5} & 0 & -3+\sqrt{5} & \sqrt{5(5-2\sqrt{5})} & -2 & \sqrt{5(5-2\sqrt{5})} & -3+\sqrt{5} & 0 & -2+\sqrt{5} \\ -2 & 0 & 2-\sqrt{5} & \sqrt{10-2\sqrt{5}} & -2-\sqrt{5} & \sqrt{2(5+\sqrt{5})} & -3+\sqrt{5} & -\sqrt{5+2\sqrt{5}} & 3+\sqrt{5} & -\sqrt{25-2\sqrt{5}} \\ 0 & 2 & 0 & 2-\sqrt{5} & \sqrt{5(5+2\sqrt{5})} & -3-\sqrt{5} & 0 & 3+\sqrt{5} & -\sqrt{5(5+2\sqrt{5})} & 2+\sqrt{5} \\ 2+\sqrt{5} & -\sqrt{25-2\sqrt{5}} & -2+\sqrt{5} & \sqrt{25+2\sqrt{5}} & 8 & \sqrt{25+2\sqrt{5}} & -2+\sqrt{5} & -\sqrt{25-2\sqrt{5}} & 2+\sqrt{5} & 0 \end{pmatrix}$$

Let us now write down all the relevant quantities up to the needed order (here and below, unless otherwise specified, we omit the Fermat parameter $k = 10$):

$$\tilde{K} \approx (5 - \sqrt{2}) \frac{C_2^2}{C_0^2} (\psi \bar{\psi})^2 \left[1 + \frac{(5-\sqrt{2})}{2} \frac{C_2^2}{C_0^2} (\psi \bar{\psi})^2 \right] + \mathcal{O}(\psi^9); \quad (10.1.3)$$

$$g \approx 4(5 - \sqrt{2}) \frac{C_2^2}{C_0^2} \psi \bar{\psi} \left[1 + 2(5 - \sqrt{2}) \frac{C_2^2}{C_0^2} (\psi \bar{\psi})^2 \right] + \mathcal{O}(\psi^7); \quad (10.1.4)$$

$$\widetilde{W} \approx N_1 + \frac{C_2}{C_0} N_3 \psi^2 - \frac{C_6}{C_0} \bar{N}_3 \psi^6 + \mathcal{O}(\psi^8). \quad (10.1.5)$$

Now, by using the formulæ of the general analysis exploited in Sect. 6, we can get the “effective BH potential” and the holomorphic superpotential, as well as their (covariant) derivatives, up to $\mathcal{O}(\psi^2)$:

$$\widetilde{W} = N_1 + \frac{C_2}{C_0} N_3 \psi^2; \quad (10.1.6)$$

$$D\widetilde{W} = 2\frac{C_2}{C_0} N_3 \psi + 2(\sqrt{5} - 2) \frac{C_2^2}{C_0^2} N_1 \psi \bar{\psi}^2; \quad (10.1.7)$$

$$D^2\widetilde{W} = -24\frac{C_6}{C_0} \bar{N}_3 \psi^4; \quad (10.1.8)$$

$$\tilde{V}_{BH} = |N_1|^2 + (\sqrt{5} + 2)|N_3|^2 + 2\frac{C_2}{C_0} N_3 \bar{N}_1 \psi^2 + 2\frac{C_2}{C_0} \bar{N}_3 N_1 \bar{\psi}^2; \quad (10.1.9)$$

$$\partial\tilde{V}_{BH} = 4\psi \left(\frac{C_2}{C_0} N_3 \bar{N}_1 - 3(\sqrt{5} + 2) \frac{C_6}{C_2} (\bar{N}_3)^2 \psi^2 + \frac{C_2^2}{C_0^2} (|N_1|^2(\sqrt{5} - 2) + |N_3|^2) \bar{\psi}^2 \right); \quad (10.1.10)$$

$$\partial^2 \tilde{V}_{BH} = 4 \left(\frac{C_2}{C_0} N_3 \bar{N}_1 - 9(\sqrt{5} + 2) \frac{C_6}{C_2} (\bar{N}_3)^2 \psi^2 + \frac{C_2^2}{C_0^2} (|N_1|^2(\sqrt{5} - 2) + |N_3|^2) \bar{\psi}^2 \right); \quad (10.1.11)$$

$$\partial\bar{\partial}\tilde{V}_{BH} = 8 \frac{C_2^2}{C_0^2} (|N_1|^2(\sqrt{5} - 2) + |N_3|^2) \psi \bar{\psi}. \quad (10.1.12)$$

Contrary to the cases of mirror *quintic* and *sextic* (see Sects. 7 and 8, respectively), and similarly to the case of mirror *octic* (see Sect. 9), for the case of mirror *dectic* it is evident from Eqs. (10.1.9)-(10.1.12) that we truncate the series expansion of the “effective BH potential” and of its second derivatives around the LG point up to $\mathcal{O}(\psi^2)$ included, and the series expansion of its first derivative around the LG point up to $\mathcal{O}(\psi^3)$ included. This is due to the absence of an $\mathcal{O}(\psi)$ term in expression of $\tilde{V}_{BH}$ given by Eq. (10.1.9). As mentioned in Sect. 9, such a fact can be traced back to the non-regularity of g at $\psi = 0$ (see Eq. (10.1.4)).

10.1.1 The Holomorphic Prepotentials F and $\mathcal{F}$

As done for $k = 5, 6, 8$ respectively in Subsubsects. 7.1.1, 8.1.1 and 9.1.1, in order to compute the holomorphic prepotential $F(X(\psi))$ and its “Kähler-invariant counterpart” $\mathcal{F}(\psi)$ defined by Eq. (4.1.9), we have to recall the general formulæ for (mirror) Fermat CY_3 s introduced in Sect. 6, and specialize them for $k = 10$. By recalling the definitions (6.38), (6.40), (6.41), (6.42), one can compute²⁶:

$$\xi_{10,m} = \begin{pmatrix} 2\cos\left(\frac{\pi m}{10}\right) + \cos\left(\frac{3\pi m}{10}\right) - i\sin\left(\frac{3\pi m}{10}\right) \\ -\cos\left(\frac{\pi m}{10}\right) + i\sin\left(\frac{\pi m}{10}\right) \\ 2i\sin\left(\frac{3\pi m}{10}\right) \\ 2i\sin\left(\frac{\pi m}{10}\right) \end{pmatrix}; \quad (10.1.1.1)$$

$$\begin{aligned} \Phi_{10,m,n} &= \\ &= \left[\cos\left(\frac{3\pi(n-m)}{10}\right) - \cos\left(\frac{\pi(n-m)}{10}\right) + \cos\left(\frac{\pi(n+m)}{10}\right) - \cos\left(\frac{3\pi(n+m)}{10}\right) + \right. \\ &\quad \left. + 5\cos\left(\frac{\pi(n+m)}{8}\right) + 3\cos\left(\frac{3\pi(n+m)}{8}\right) + 3\cos\left(\frac{\pi(3n+m)}{8}\right) + 3\cos\left(\frac{\pi(n+3m)}{8}\right) \right] + \\ &+ i \left[\sin\left(\frac{3\pi(n-m)}{10}\right) - \sin\left(\frac{\pi(n-m)}{10}\right) + 2\sin\left(\frac{\pi(3n-m)}{10}\right) - \sin\left(\frac{\pi(n+m)}{10}\right) + \right. \\ &\quad \left. + 2\sin\left(\frac{\pi(3n+m)}{10}\right) + \sin\left(\frac{3\pi(n+m)}{10}\right) \right] \neq \Phi_{10,(m,n)}; \end{aligned} \quad (10.1.1.2)$$

²⁶As implied by Eqs. (7.1.1.2), (8.1.1.2), (9.1.1.2) and (10.1.1.2), it holds that

$$\text{Re}(\Phi_{k,m,n}) = \text{Re}(\Phi_{k,(m,n)}), \quad \forall k = 5, 6, 8, 10,$$

while the same relation for the imaginary part holds only for $k = 5$. From the definitions (6.38) and (6.40), this fact can be traced back to value of k itself and to the explicit form of the matrices M_k s, given by Eqs. (6.16) and (6.17).

$$\begin{aligned}
F_{k=10,l=9}(X_{10}(\psi)) &= -\frac{2\pi^6}{5^4} \sum_{m=1}^9 \sum_{n=1}^9 (-1)^{m+n} C_{10,m-1} C_{10,n-1} \psi^{m+n-2} \Phi_{10,(m,n)} = \\
&= -\frac{2\pi^6}{5^4} \left[\begin{aligned} &\frac{(3+\sqrt{5})(2\sqrt{10}+i(5+\sqrt{5})^{3/2})\pi}{10\Gamma^2(\frac{4}{5})\Gamma^6(\frac{9}{10})2^{9/10}} + \\ &-\frac{5(5-3\sqrt{5})(1+i\sqrt{5-2\sqrt{5}})\pi}{\Gamma^2(\frac{2}{5})\Gamma^6(\frac{7}{10})2^{6/5}}\psi^4 + \\ &+\frac{25\sqrt{5(25+2\sqrt{5})}}{96\pi^3}\psi^8 \end{aligned} \right] + \mathcal{O}(\psi^9) = F_{k=10,l=9}(\psi^4); \tag{10.1.1.3}
\end{aligned}$$

$$\begin{aligned}
\mathcal{F}_{10}(\psi) &\equiv F_{10}\left(\frac{X_{10}^1(\psi)}{X_{10}^0(\psi)}\right) = \frac{F_{10}(X_{10}^0(\psi), X_{10}^1(\psi))}{(X_{10}^0(\psi))^2} = \frac{1}{2} \frac{\sum_{m,n=1}^{\infty} (-1)^{m+n} C_{10,m-1} C_{10,n-1} \Phi_{10,(m,n)} \psi^{m+n-2}}{[\sum_{r=1}^{\infty} (-1)^r C_{10,r-1} \xi_{10,r}^1 \psi^{r-1}]^2} \approx \\
&\approx \frac{\Phi_{10,1,1}}{2(\xi_{10,1}^1)^2} \left[1 + \frac{C_{10,2}(\Phi_{10,1,3} + \Phi_{10,3,1})}{C_{10,0}\Phi_{10,1,1}} \psi^2 + \mathcal{O}(\psi^3) \right] \left[1 - 2 \frac{C_{10,2}\xi_{10,3}^1}{C_{10,0}\xi_{10,1}^1} \psi^2 + \mathcal{O}(\psi^3) \right] \approx \\
&\approx \frac{\Phi_{10,1,1}}{2(\xi_{10,1}^1)^2} \left[1 - 2 \left(\frac{\xi_{10,3}^1}{\xi_{10,1}^1} - \frac{\Phi_{10,1,3} + \Phi_{10,3,1}}{2\Phi_{10,1,1}} \right) \frac{C_{10,2}}{C_{10,0}} \psi^2 \right] + \mathcal{O}(\psi^3), \tag{10.1.1.4}
\end{aligned}$$

where in the formula (10.1.1.4) for the Kähler-invariant holomorphic prepotential $\mathcal{F}_{10}$ only the leading terms in the *LG limit* $\psi \rightarrow 0$ have been retained. Eq. (10.1.1.3) is the exact expression (complete up to $\mathcal{O}(\psi^8)$ included) of the holomorphic prepotential²⁷ F_{10} in a suitable neighbourhood of the LG point of the moduli space of the (mirror) *dectic* $\mathcal{M}'_{10}$.

10.2 “Criticality Condition” Approach and Attractor Equation

Let us now find the solutions of the AE $\partial \tilde{V}_{BH}(\psi, \bar{\psi}; q, p) = 0$, and check their stability. Since we are working near the LG point, by using Eq. (10.1.9) we can rewrite the AE for $\mathcal{M}'_{10}$ as follows:

$$\frac{C_2}{C_0} N_3 \bar{N}_1 + \frac{C_2^2}{C_0^2} \left(|N_1|^2 (\sqrt{5} - 2) + |N_3|^2 \right) \bar{\psi}^2 \approx 3(\sqrt{5} + 2) \frac{C_6}{C_2} (\bar{N}_3)^2 \psi^2 \tag{10.2.1}$$

Solving Eq. (9.2.1), we will find one (or more) set(s) of BH charges supporting $\psi \approx 0$ to be a critical point of V_{BH} . Since we are working near the LG point, it is clear that the first term in the l.h.s. of Eq. (10.2.1) must be small enough. This implies the following fine-tuning condition:

$$N_3 \bar{N}_1 \approx 0. \tag{10.2.2}$$

²⁷It is worth noticing that F_8 and F_{10} (respectively given by Eqs. (9.1.1.3) and (10.1.1.3)) are in turn actually functions of ψ^2 (as explicitly checked up to $\mathcal{O}(\psi^{10})$ included) and ψ^4 (as explicitly checked up to $\mathcal{O}(\psi^{16})$ included). This fact distinguishes the cases $k = 8, 10$ from the cases $k = 5, 6$ (respectively given by Eqs. (7.1.1.3) and (8.1.1.3)), and it can be traced back to the explicit form of the matrices M_k s (given by Eqs. (6.16) and (6.17)) and of the constants $C_{k,m-1}$ (defined in the first of Eqs. (6.12)).

10.3 Critical Hessian of V_{BH}

By using (the $n_V = 1$ case of) Eq. (3.1.11) and Eqs. (10.1.11)-(10.1.12) evaluated along the criticality condition (10.2.1)-(10.2.2), it can be computed that the components of $H_{\text{real form}}^{\tilde{V}_{BH}}$ (given by Eq. (7.3.1)) constrained by the Eqs. (10.2.1)-(10.2.2) read as follows:

$$\mathcal{A} = -12(\sqrt{5} + 2) \frac{C_6}{C_2} ((\bar{N}_3)^2 \psi^2 + (N_3)^2 \bar{\psi}^2) + 8 \frac{C_2^2}{C_0^2} \psi \bar{\psi} \left(|N_1|^2 (\sqrt{5} - 2) + |N_3|^2 \right); \quad (10.3.1)$$

$$\mathcal{B} = 12(\sqrt{5} + 2) \frac{C_6}{C_2} ((\bar{N}_3)^2 \psi^2 + (N_3)^2 \bar{\psi}^2) + 8 \frac{C_2^2}{C_0^2} \psi \bar{\psi} \left(|N_1|^2 (\sqrt{5} - 2) + |N_3|^2 \right); \quad (10.3.2)$$

$$\mathcal{C} = -12i(\sqrt{5} + 2) \frac{C_6}{C_2} ((N_3)^2 \bar{\psi}^2 - (\bar{N}_3)^2 \psi^2). \quad (10.3.3)$$

The resulting real eigenvalues of $H_{\text{real form}}^{\tilde{V}_{BH}}$ constrained by Eqs. (10.2.1)-(10.2.2) read:

$$\lambda_{\pm} \approx 8\psi\bar{\psi} \left[\frac{C_2^2}{C_0^2} \left(|N_1|^2 (\sqrt{5} - 2) + |N_3|^2 \right) \pm 3(\sqrt{5} + 2) \frac{C_6}{C_2} |N_3|^2 \right]. \quad (10.3.4)$$

By recalling Eq. (6.34) and using Eq. (10.1.9) with $\psi \approx 0$ and constrained by Eqs. (10.2.1)-(10.2.2), one obtains that the purely charge-dependent LG critical values of the “effective BH potential” for the mirror *dectic* $\mathcal{M}'_{10}$ are

$$V_{BH, LG-critical, k=10} \approx \frac{1}{\sqrt{5(5 + 2\sqrt{5})}} \left[|N_1|^2 + (\sqrt{5} + 2)|N_3|^2 \right]; \quad (10.3.5)$$

by recalling formula (1.3), this directly yields the following purely charge-dependent values of the BH entropy at the LG critical points of $V_{BH,10}$ in the moduli space of $\mathcal{M}'_{10}$:

$$S_{BH, LG-critical, k=10} \approx \frac{\pi}{\sqrt{5(5 + 2\sqrt{5})}} \left[|N_1|^2 + (\sqrt{5} + 2)|N_3|^2 \right]. \quad (10.3.6)$$

From the definition (6.31), for $k = 10$ and $m = 1, 3$ one respectively gets that

$$N_1 = -\frac{1}{2} \sqrt{\frac{5 - \sqrt{5}}{2}} \left(n_1 + n_4 + \frac{(1 + \sqrt{5})}{2} (n_2 + n_3) \right) + i \frac{\sqrt{5} + 1}{4} \left(n_4 - n_1 + \frac{3 - \sqrt{5}}{2} (n_3 - n_2) \right); \quad (10.3.7)$$

$$N_3 = \frac{1}{2} \sqrt{\frac{(5 - \sqrt{5})}{2}} \left(n_1 + n_4 - \frac{\sqrt{5} - 1}{2} (n_2 + n_3) \right) + i \frac{\sqrt{5} - 1}{4} \left(n_4 - n_1 + \frac{3 + \sqrt{5}}{2} (n_3 - n_2) \right). \quad (10.3.8)$$

Let us write down here the numerical values of constants relevant to our treatment:

$$C_0 \approx 1.57, \quad C_2 \approx -0.66, \quad C_6 \approx 0.077; \quad 3(\sqrt{5} + 2) \frac{C_6}{C_2} \approx -1.48. \quad (10.3.9)$$

10.4 Solutions to Attractor Equations

Let us now analyze more in depth the species of LG attractor points arising from the AE (10.2.2). As it can be easily seen, the AE (10.2.2) has two *non-degenerate* solutions:

I. The first non-degenerate solution to AE (10.2.2) is

$$N_3 \approx 0. \quad (10.4.1)$$

This is nothing but the $k = 8$ solution (9.4.1). As one can see from Eq. (10.1.6)-(10.1.7), also for $k = 10$ such a solution corresponds to a $\frac{1}{2}$ -BPS LG critical point of V_{BH} ($\widetilde{W} \neq 0$, $D\widetilde{W} = 0$). From Eq. (10.3.8), in order to get the solution (10.4.1), one has to fine-tune two PF BH charges out of four in the following way:

$$n_3 \approx \frac{1}{2}(\sqrt{5} - 1)(n_1 + n_2), \quad n_4 \approx -\frac{1}{2}(\sqrt{5} - 1)n_1 + n_2. \quad (10.4.2)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition $N_1 \neq 0$. As it was for $k = 5, 6, 8$, also the real eigenvalues (10.3.4) for the $\frac{1}{2}$ -BPS critical solution coincide and, as it is well known [39, 15, 16], are strictly positive:

$$\lambda_{+, \frac{1}{2}-BPS} = \lambda_{-, \frac{1}{2}-BPS} \approx 8(\sqrt{5} - 2)\psi\bar{\psi}\frac{C_2^2}{C_0^2}|N_1|_{N_3 \approx 0}^2 > 0. \quad (10.4.3)$$

Consequently, the $\frac{1}{2}$ -BPS LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (10.4.2) is a stable extremum, since it is a (local) minimum of V_{BH} , and it is therefore an attractor in a strict sense. The classical (Bekenstein-Hawking) BH entropy at such a (class of) $\frac{1}{2}$ -BPS LG attractor(s) takes the value

$$S_{BH, \frac{1}{2}-BPS} = \pi V_{BH, \frac{1}{2}-BPS} \approx 0.166\pi|N_1|_{N_3 \approx 0}^2, \quad (10.4.4)$$

where $|N_1|_{N_3 \approx 0}$ is given by Eq. (10.3.7) constrained by Eq. (10.4.2):

$$N_1|_{N_3 \approx 0} \approx -\frac{\sqrt{5}}{2} \left(\sqrt{5 - 2\sqrt{5}}n_1 + \sqrt{\frac{5 + \sqrt{5}}{2}}n_2 + i \left(n_1 - \frac{\sqrt{5} - 1}{2}n_2 \right) \right). \quad (10.4.5)$$

II. The second non-degenerate solution to AE (10.2.2) is

$$N_1 \approx 0. \quad (10.4.6)$$

Interestingly, this is nothing but the $k = 6$ solution (8.4.6). As one can see from Eq. (10.1.6)-(10.1.7), also for $k = 10$ such a solution corresponds to a non-BPS, $Z = 0$ LG critical point of V_{BH} ($\widetilde{W} = 0$, $D\widetilde{W} \neq 0$). The real eigenvalues (10.3.4) for such a non-BPS, $Z = 0$ critical solution read

$$\lambda_{\pm, non-BPS, Z=0} \approx 8\psi\bar{\psi} \left[\frac{C_2^2}{C_0^2} \pm 3(\sqrt{5} + 2)\frac{C_6}{C_2} \right] |N_3|_{N_1 \approx 0}^2. \quad (10.4.7)$$

Substituting the numerical values (10.3.9) of the involved constants in Eq. (10.4.7), one reaches the conclusion that one eigenvalue is positive and the other one is negative:

$$\lambda_{\pm, non-BPS, Z=0} \approx 8\psi\bar{\psi} [0.18 \mp 1.48] |N_3|_{N_1 \approx 0}^2 \lesseqgtr 0. \quad (10.4.8)$$

Let us now find the fine-tuning conditions for PF BH charges supporting the considered non-BPS, $Z = 0$ LG attractor for the mirror *dectic* $\mathcal{M}'_{10}$. This amounts to solving Eq. (10.4.6) by recalling Eq. (10.3.7). By doing so, one gets the following unique set of constraining relations on PF BH charges:

$$n_3 \approx -\frac{1}{2}(1 + \sqrt{5})(n_1 + n_2), \quad n_4 \approx \frac{1}{2}(1 + \sqrt{5})n_1 + n_2. \quad (10.4.9)$$

The charges n_1, n_2 are not fixed; they only satisfy the non-degeneration condition $N_3 \neq 0$. Thus, the non-BPS, $Z = 0$ LG critical point $\psi \approx 0$ supported by the PF BH charge configuration (10.4.9) is a saddle point of V_{BH} and consequently it is *not* an attractor in a strict sense.

The classical (Bekenstein-Hawking) BH entropy at such a (class of) non-BPS, $Z = 0$ LG saddle point(s) takes the value

$$S_{BH, non-BPS, Z=0} = \pi V_{BH, non-BPS, Z=0} \approx 0.7\pi |N_3|_{N_1 \approx 0}^2, \quad (10.4.10)$$

where $N_3|_{N_1 \approx 0}$ is given by Eq. (10.3.8) constrained by Eq. (10.4.9):

$$N_3|_{N_1 \approx 0} \approx \frac{\sqrt{5}}{2} \left(\sqrt{5 + 2\sqrt{5}} n_1 + \sqrt{\frac{5 - \sqrt{5}}{2}} n_2 - i \left(n_1 + \frac{\sqrt{5} + 1}{2} n_2 \right) \right). \quad (10.4.11)$$

Similarly to the treatment of Sects. 7-9, one can also switch to the symplectic (electric/magnetic) basis for BH charges by using Eq. (6.33) for $k = 10$, and thus re-express the fine-tuning conditions (10.4.2) and (10.4.9) in terms of the (components of the) symplectic BH charge vector Γ defined by Eq. (6.28).

11 Special Kähler Geometrical Identities and Attractors

11.1 $n_V = 1$ Identities near the LG Point

Let us consider the real part of the $n_V = 1$ case of SK geometry identities (2.1.14) [55, 8, 13, 15, 16, 24]; by taking into account the change in the notation of the symplectic charge vectors pointed out in the Footnote before Eq. (6.28), one achieves:

$$\tilde{\Gamma}^T = 2e^K \text{Im} [W\bar{\Pi} + g^{-1} (\bar{D}\bar{W}) D\Pi], \quad (11.1.1)$$

where the 1×4 BH symplectic charge vector $\tilde{\Gamma}$ is defined in Eq. (6.28).

Next, let us switch to more convenient variables for the treatment of 1-modulus SK geometries endowing the moduli space of Fermat CY_3 s. By recalling the definition (6.32) of the 1×4 PF BH charge vector n , we can rewrite Eq. (11.1.1) as follows (here and below, unless otherwise specified, we omit the classifying Fermat parameter $k = 5, 6, 8, 10$):

$$n^T = \frac{2}{\chi F_{11}} e^{\tilde{K}} \text{Im} [\tilde{W}\tilde{\Upsilon} + g^{-1} (\bar{D}\bar{W}) D\tilde{\Upsilon}], \quad (11.1.2)$$

where the notations introduced in Sect. (6) have been used. Furthermore, we defined the 4×1 holomorphic vector

$$\tilde{\Upsilon}(\psi) \equiv \frac{1}{AC_0} M^T \Sigma^T \Pi(\psi) = \frac{1}{AC_0} m^T \varpi(\psi), \quad (11.1.3)$$

where Eqs. (6.15) and (6.23) have been used.

Let us investigate Eq.(11.1.2) in a certain neighbourhood of the LG point $\psi = 0$. The treatment given in Sects. 7-10 yields that, by its very definition (11.1.3), $\tilde{\Upsilon}$ has the following series expansion near the LG point²⁸:

$$k = 5 : \begin{cases} \tilde{\Upsilon} = \varphi_1 + \frac{C_1}{C_0} \varphi_2 \psi, \\ D\tilde{\Upsilon} = \frac{C_1}{C_0} \left[\varphi_2 + 2\frac{C_2}{C_1} \bar{\varphi}_2 \psi + \frac{C_1}{C_0} (\sqrt{5} - 2) \varphi_1 \bar{\psi} \right]; \end{cases} \quad (11.1.4)$$

$$k = 6 : \begin{cases} \tilde{\Upsilon} = \varphi_1 + \frac{C_1}{C_0} \varphi_2 \psi, \\ D\tilde{\Upsilon} = \frac{C_1}{C_0} \left[\varphi_2 + \frac{1}{3} \frac{C_1}{C_0} \varphi_1 \bar{\psi} \right]; \end{cases} \quad (11.1.5)$$

$$k = 8 : \begin{cases} \tilde{\Upsilon} = \varphi_1 + \frac{C_2}{C_0} \varphi_3 \psi^2, \\ D\tilde{\Upsilon} = 2\frac{C_2}{C_0} \psi \left[\varphi_3 - 2\frac{C_4}{C_2} \bar{\varphi}_3 \psi^2 + \frac{C_2}{C_0} (3 - 2\sqrt{2}) \varphi_1 \bar{\psi}^2 \right]; \end{cases} \quad (11.1.6)$$

$$k = 10 : \begin{cases} \tilde{\Upsilon} = \varphi_1 + \frac{C_2}{C_0} \varphi_3 \psi^2, \\ D\tilde{\Upsilon} = 2\frac{C_2}{C_0} \psi \left[\varphi_3 + \frac{C_2}{C_0} (\sqrt{5} - 2) \varphi_1 \bar{\psi}^2 \right]. \end{cases} \quad (11.1.7)$$

We defined the complex 4×1 vector²⁹

$$\varphi_{k,m} \equiv m_k \begin{pmatrix} \beta_k^{3m} \\ \beta_k^m \\ \beta_k^{-m} \\ \beta_k^{-3m} \end{pmatrix}, \quad (11.1.8)$$

²⁸Notice that, consistently with the approach to the truncation of series expansions near the LG point performed in Sects. 7-10, for $k = 5, 6$ we truncate up to $\mathcal{O}(\psi)$ included, whereas for $k = 8, 10$ we truncate up to $\mathcal{O}(\psi^2)$ included.

²⁹By recalling the definitions (6.23) and (6.38), it is worth noticing that

$$\varphi_{k,m} = M_k^T \Sigma \xi_{k,m}.$$

whose explicit forms (for $m = 1, 2, 3$, the only values relevant for the series expansion (11.1.4)-(11.1.7)) read as follows:

$$k = 5 : \begin{cases} \varphi_1 = \frac{1}{2} \sqrt{\frac{5+\sqrt{5}}{2}} \begin{pmatrix} -\sqrt{5+2\sqrt{5}} + i\sqrt{5} \\ -\sqrt{\frac{5+\sqrt{5}}{2}} + \frac{1}{2}i(5+3\sqrt{5}) \\ \sqrt{\frac{5+\sqrt{5}}{2}} + \frac{1}{2}i(5+3\sqrt{5}) \\ \sqrt{5+2\sqrt{5}} + i\sqrt{5} \end{pmatrix}, \\ \varphi_2 = \frac{1}{2} \sqrt{\frac{5-\sqrt{5}}{2}} \begin{pmatrix} \sqrt{5-2\sqrt{5}} - i\sqrt{5} \\ -\sqrt{\frac{5-\sqrt{5}}{2}} + \frac{1}{2}i(5-3\sqrt{5}) \\ \sqrt{\frac{5-\sqrt{5}}{2}} + \frac{1}{2}i(5-3\sqrt{5}) \\ -\sqrt{5-2\sqrt{5}} - i\sqrt{5} \end{pmatrix}; \end{cases} \quad (11.1.9)$$

$$k = 6 : \begin{cases} \varphi_1 = \frac{3}{2} \begin{pmatrix} -\sqrt{3} + i \\ 2i \\ 2i \\ \sqrt{3} + i \end{pmatrix}, \\ \varphi_2 = \frac{1}{2} \begin{pmatrix} 1 - i\sqrt{3} \\ -1 \\ 1 \\ -1 - i\sqrt{3} \end{pmatrix}; \end{cases} \quad (11.1.10)$$

$$k = 8 : \begin{cases} \varphi_1 = \frac{1}{2} \sqrt{2+\sqrt{2}} \begin{pmatrix} -2 - \sqrt{2} + i\sqrt{2} \\ -\sqrt{2} + i(2+\sqrt{2}) \\ \sqrt{2} + i(2+\sqrt{2}) \\ 2 + \sqrt{2} + i\sqrt{2} \end{pmatrix}, \\ \varphi_3 = \frac{1}{2} \sqrt{2-\sqrt{2}} \begin{pmatrix} -2 + \sqrt{2} + i\sqrt{2} \\ \sqrt{2} + i(-2+\sqrt{2}) \\ -\sqrt{2} + i(-2+\sqrt{2}) \\ 2 - \sqrt{2} + i\sqrt{2} \end{pmatrix}; \end{cases} \quad (11.1.11)$$

$$k = 10 : \begin{cases} \varphi_1 = \frac{1}{2} \sqrt{\frac{5+\sqrt{5}}{2}} \begin{pmatrix} -\frac{1}{2}\sqrt{3+\sqrt{5}} + i\sqrt{\frac{5+\sqrt{5}}{10}} \\ 1 + i\sqrt{1+\frac{2}{\sqrt{5}}} \\ -1 + i\sqrt{1+\frac{2}{\sqrt{5}}} \\ \frac{1}{2}\sqrt{3+\sqrt{5}} + i\sqrt{\frac{5+\sqrt{5}}{10}} \end{pmatrix}, \\ \varphi_3 = \frac{1}{2} \sqrt{\frac{5-\sqrt{5}}{2}} \begin{pmatrix} \frac{1}{2}\sqrt{-3+\sqrt{5}} + i\sqrt{\frac{5-\sqrt{5}}{10}} \\ 1 - i\sqrt{1-\frac{2}{\sqrt{5}}} \\ -1 - i\sqrt{1-\frac{2}{\sqrt{5}}} \\ \frac{1}{2}\sqrt{3-\sqrt{5}} + i\sqrt{\frac{5-\sqrt{5}}{10}} \end{pmatrix}. \end{cases} \quad (11.1.12)$$

Using Eqs. (11.1.3)-(11.1.12), one obtains that Eq. (11.1.2) near the LG point of the moduli space of Fermat CY_3 -compactifications (when consistently truncated up to the order in ψ considered above) reads

$$k = 5, 6 : \quad n^T = \frac{2}{\chi F_{11}} \text{Im} \left[N_1 \bar{\varphi}_1 - \frac{F_{11}}{F_{22}} \bar{N}_2 \varphi_2 \right]; \quad (11.1.13)$$

$$k = 8, 10 : \quad n^T = \frac{2}{\chi F_{11}} \text{Im} \left[N_1 \bar{\varphi}_1 - \frac{F_{11}}{F_{33}} \bar{N}_3 \varphi_3 \right]. \quad (11.1.14)$$

Thence, it is easy to check that, substituting the explicit forms of $N_{k,m}(q, p)$, $\varphi_{k,m}$ and $F_{k,mn}$ (see Eqs. (6.31), (11.1.8) and (6.25), respectively) in Eqs. (11.1.13)-(11.1.14), they become trivial identities, yielding nothing but

$$n_1 = n_1, \quad n_2 = n_2, \quad n_3 = n_3, \quad n_4 = n_4. \quad (11.1.15)$$

In other words, the 4 real Eqs. (11.1.2) are not equations, but rather they are identities. Therefore, they are satisfied at every point in the moduli space and for every BH charge configuration. Therefore, it is no surprise if, when evaluating them in a certain neighbourhood of the LG point as we did, one finds the identical relations (11.1.15). Thus, we found nothing new but another confirmation of a well known fact of SK geometry [55, 8, 13, 15, 16, 24].

11.2 “Special Kähler Geometrical Identities” Approach to LG Attractors in Fermat CY_3 s

However, the 4 real identities (11.1.2) can still be used to find extremal BH attractors, when properly evaluated along the constraints defining the various species of such attractors satisfying the criticality condition of the “effective BH potential” V_{BH} . Put another way, in the 1-modulus case with which we are concerned, when evaluated at the geometrical *loci* in the moduli space defining the various tipologies (*i.e.* $\frac{1}{2}$ -BPS, non-BPS $Z \neq 0$ and non-BPS $Z = 0$) of attractors, the 4 real identities (11.1.2) become 4 real equations. These are equivalent to the 2 real equations given by the real and imaginary part of the criticality condition $\partial V_{BH} = 0$. This approach has recently been used in [8] for the general n_V -moduli case, and then further investigated in [13]. The SK geometrical identities in the general n_V -moduli case had previously been formulated in [55] in terms of the decomposition of the third real cohomology $H^3(CY_3; \mathbb{R})$ of the CY_3 in the Dolbeaut cohomology basis (see [55], and [48] for further Refs.).

We will now focus on the 1-modulus case related to Fermat CY_3 -compactifications, and we will evaluate the 4 real identities (11.1.2) at the geometrical *loci* in the moduli space defining the various classes of extremal BH attractors. We will consequently show that solving the obtained 4 real equations is equivalent to solving the 2 real equations corresponding to the real and imaginary parts of the criticality condition (2.2.2). Thus, it follows that only 2 equations are independent out of the starting 4 ones. From a computational point of view, one can realize that, at least in the framework we are considering, the “criticality condition” approach is simpler than the “SK geometrical identities” approach, at least for the non-BPS, $Z \neq 0$ case.

Let us now evaluate the 4 real SK identities (11.1.2) along the 3 geometrical *loci* defining the 3 species of critical points of V_{BH} arising in SK geometry.

11.2.1 $\frac{1}{2}$ -BPS

The corresponding geometrical *locus* in the moduli space is given by the constraints $\widetilde{W} \neq 0$, $D\widetilde{W} = 0$, which directly solve the criticality condition (*i.e.* the 1-modulus AE) (2.2.2). By evaluating the 4 real SK identities (11.1.2) along such critical constraints, one gets

$$n^T = \frac{2}{\chi F_{11}} \left[e^{\tilde{K}} \text{Im} \left(\widetilde{W} \widetilde{\Upsilon} \right) \right]_{\frac{1}{2}\text{-BPS}}. \quad (11.2.1.1)$$

Such 4 real equations constrain the PF BH charge configurations along the *locus* $\widetilde{W} \neq 0$, $D\widetilde{W} = 0$ of $\frac{1}{2}$ -BPS critical points of V_{BH} in the moduli space ($\dim_{\mathbb{C}} = 1$) of Fermat CY_3 s. One can explicitly check that for all Fermat CY_3 s the solutions of the 4 real Eqs. (11.2.1.1) near the LG point give nothing but the $\frac{1}{2}$ -BPS-supporting PF BH charge configurations previously computed in Sects. 7-10 exploiting the so-called “criticality condition” approach.

11.2.2 Non-BPS, $Z \neq 0$

The corresponding geometrical *locus* in the moduli space is given by the conditions $\widetilde{W} \neq 0$, $D\widetilde{W} \neq 0$, further constrained by the criticality condition (2.2.2); by also recalling the definition (6.35), one obtains

$$\left(\overline{D\widetilde{W}} \right)_{non-BPS, Z \neq 0} = - \left[\frac{g^{-1} \left(\overline{D^2 \widetilde{W}} \right) D\widetilde{W}}{2\widetilde{W}} \right]_{non-BPS, Z \neq 0}. \quad (11.2.2.1)$$

By inserting Eq. (11.2.2.1) in the 4 real SK identities (11.1.2), one gets [8, 13, 16]

$$n^T = \frac{2}{\chi F_{11}} \left\{ e^{\tilde{K}} \text{Im} \left[\widetilde{W} \widetilde{\Upsilon} - \frac{g^{-2} \left(\overline{D^2 \widetilde{W}} \right) D\widetilde{W}}{2\widetilde{W}} D\widetilde{\Upsilon} \right] \right\}_{non-BPS, Z \neq 0} = \quad (11.2.2.2)$$

$$= \frac{2}{\chi F_{11}} \left\{ e^{\tilde{K}} \text{Im} \left[\widetilde{W} \widetilde{\Upsilon} + i \frac{g^{-3} C \left(\overline{D\widetilde{W}} \right)^2}{2\widetilde{W}} \overline{D\widetilde{\Upsilon}} \right] \right\}_{non-BPS, Z \neq 0}, \quad (11.2.2.3)$$

where, in the second line, we used the $n_V = 1$ case of the second SK differential relation of (2.1.8), yielding globally

$$D^2 \widetilde{W} = i C g^{-1} \overline{D\widetilde{W}}. \quad (11.2.2.4)$$

The 4 real Eqs. (11.2.2.2)-(11.2.2.3) constrain the PF BH charge configurations along the *locus* (11.2.2.1) of non-BPS $Z \neq 0$ critical points of V_{BH} in the moduli space of Fermat CY_3 s.

Let us for example consider the mirror *quintic* $\mathcal{M}'_5$. From the treatment given in Sect. 7 and above, the 4 real Eqs. (11.2.2.2) take the following form near the LG point:

$$n^T = \frac{2}{\chi F_{11}} \text{Im} \left[N_1 \bar{\varphi}_1 - (\sqrt{5} + 2) \xi \frac{N_2 N_2}{N_1} \varphi_2 \right], \quad (11.2.2.5)$$

where ξ is defined in Eq. (7.4.6). Substituting into Eqs. (11.2.2.5) the explicit expressions for the N s (see Sect. 7) and the φ s (see Eq. (11.1.9)) and performing long but straightforward computations, it can be shown that one generally recovers all the three distinct sets of BH charge configurations (7.4.12)-(7.4.14) supporting the considered non-BPS $Z \neq 0$ LG attractor. The same can be explicitly checked for the mirror *octic* $\mathcal{M}'_8$.

11.2.3 Non-BPS, $Z = 0$

The corresponding geometrical *locus* in the moduli space is given by the conditions $\widetilde{W} = 0$, $D\widetilde{W} \neq 0$, further constrained by the criticality condition (2.2.2); by also recalling and the definition (6.35), one obtains

$$D^2\widetilde{W}\Big|_{non-BPS, Z=0} = 0. \quad (11.2.3.1)$$

By recalling Eq. (2.2.5), the replacement of $\widetilde{W} = 0$ and of the condition (11.2.3.1) into the 4 real SK geometrical identities (11.1.2) yields the following 4 real equations:

$$n^T = -\frac{2}{\chi F_{11}} \left\{ g^{-1} e^{\tilde{K}} \text{Im} \left[\frac{\partial^2 \widetilde{W} + (\partial \tilde{K}) \partial \widetilde{W}}{\partial [\ln(g) - \tilde{K}]} \overline{D\tilde{Y}} \right] \right\}_{non-BPS, Z=0}. \quad (11.2.3.2)$$

Such 4 real equations constrain the PF BH charge configurations along the *locus* (11.2.3.1) of non-BPS $Z = 0$ critical points of V_{BH} in the moduli space of Fermat CY_3 s.

Let us for example consider the mirror *sextic* $\mathcal{M}'_6$. As one can easily check by using Eqs. (2.2.5) and (8.1.6)-(8.1.8), in this case $\widetilde{W} = 0$ directly satisfies the criticality condition (2.2.2). Consequently, rather than Eqs. (11.2.3.2), in order to exploit the so-called “*SK geometrical identities*” approach, one can consider the 4 real equations

$$n^T = \frac{2}{\chi F_{11}} \left\{ g^{-1} e^{\tilde{K}} \text{Im} \left[(\overline{\partial \widetilde{W}}) D\tilde{Y} \right] \right\}_{non-BPS, Z=0}, \quad (11.2.3.3)$$

obtained from identities (11.1.2) by simply putting $\widetilde{W} = 0$ and by replacing $\overline{D\widetilde{W}}$ with $\overline{\partial \widetilde{W}}$, as implied by $\widetilde{W} = 0$. From the treatment of Sect. 8 and above, the 4 real Eqs. (11.2.3.3) take the following form near the LG point:

$$n^T = -\frac{2}{\chi F_{22}} \text{Im} [\bar{N}_2 \varphi_2]. \quad (11.2.3.4)$$

Substituting the explicit expressions for N_2 (see Eq. (8.1.14)) and for φ_2 (see Eq. (11.1.10)) into Eqs. (11.2.3.4), it can be shown that one obtains nothing but the fine-tuning conditions (8.4.9) for PF BH charges supporting the considered non-BPS $Z = 0$ LG attractor. The same can be explicitly checked for the mirror *dectic* $\mathcal{M}'_{10}$.

12 The (LG Limit of) Yukawa Coupling Function of Fermat CY_3 s

12.1 Different Approaches to Computation and LG Limit

The *Yukawa coupling function* C is nothing but the case $n_V = 1$ of the fundamental tensor C_{ijk} of SK geometry, defined by Eqs. (2.1.9). Due to its covariant holomorphicity, C can be written as (see the first of Eqs. (2.1.9)):

$$C = e^K \mathcal{W}, \quad \bar{\partial} \mathcal{W} = 0. \quad (12.1.1)$$

Because of C has Kähler weights $(2, -2)$, the function $\mathcal{W}$, usually called the *holomorphic Yukawa coupling function*, has Kähler weights $(4, 0)$. The determination of C is far from being academic; as evident from the treatments exploited in

previous Sects., C and its (covariant) derivative are key ingredients in the study of extremal BH attractors in $n_V = 1$, $\mathcal{N} = 2$, $d = 4$ MESGTs.

Thus, let us now face the question of how to determine C in the framework of the 1-dim. SK geometry (endowing the moduli space of (mirror) Fermat CY_3 s).

There exist at least three independent (and equivalent, as we will discuss further below) ways to compute C :

I) Use Eqs. (4.2.5) and (12.1.1), obtaining

$$\begin{aligned}
C(\psi, \bar{\psi}) &= e^{K(\psi, \bar{\psi})} \mathcal{W}(\psi) = \\
&= e^K \left[\left[\partial X^0(\psi) \right]^3 \frac{\partial^3 F(X)}{(\partial X^0)^3} \Big|_{X=X(\psi)} + 3 \left[\partial X^0(\psi) \right]^2 \left[\partial X^1(\psi) \right] \frac{\partial^3 F(X)}{(\partial X^0)^2 \partial X^1} \Big|_{X=X(\psi)} + \right. \\
&\quad \left. + 3 \left[\partial X^1(\psi) \right]^2 \left[\partial X^0(\psi) \right] \frac{\partial^3 F(X)}{(\partial X^1)^2 \partial X^0} \Big|_{X=X(\psi)} + \left[\partial X^1(\psi) \right]^3 \frac{\partial^3 F(X)}{(\partial X^1)^3} \Big|_{X=X(\psi)} \right] = \\
&= e^K (X^0(\psi))^2 [e(\psi)]^3 \frac{\partial^3 \mathcal{F}(t)}{(\partial t)^3} \Big|_{t=t(\psi)} = \\
&= e^K (X^0(\psi)) \left[\begin{aligned} &3 [e(\psi)]^{-2} [\partial e(\psi)]^2 \partial \mathcal{F}(\psi) - [e(\psi)]^{-1} [\partial^2 e(\psi)] \partial \mathcal{F}(\psi) + \\ &-3 [e(\psi)]^{-1} [\partial e(\psi)] \partial^2 \mathcal{F}(\psi) + \partial^3 \mathcal{F}(\psi) \end{aligned} \right].
\end{aligned} \tag{12.1.2}$$

On the other hand, by recalling Eqs. (4.1.6), (4.1.7) (4.1.9) and (4.1.10), Eq. (6.18) can be further elaborated as follows:

$$K(z, \bar{z}) = -\ln \left\{ i |X^0(z)|^2 \left[2(\mathcal{F}(z) - \bar{\mathcal{F}}(\bar{z})) - \left(\frac{X^a(z)}{X^0(z)} - \frac{\bar{X}^a(\bar{z})}{\bar{X}^0(\bar{z})} \right) \left(e_a^i(z) \frac{\partial \mathcal{F}(z)}{\partial z^i} + \bar{e}_a^{\bar{i}}(\bar{z}) \frac{\partial \bar{\mathcal{F}}(\bar{z})}{\partial \bar{z}^{\bar{i}}} \right) \right] \right\}; \tag{12.1.3}$$

$\Downarrow$

$$e^{K(z, \bar{z})} = -i |X^0(z)|^{-2} \left[2(\mathcal{F}(z) - \bar{\mathcal{F}}(\bar{z})) - \left(\frac{X^a(z)}{X^0(z)} - \frac{\bar{X}^a(\bar{z})}{\bar{X}^0(\bar{z})} \right) \left(e_a^i(z) \frac{\partial \mathcal{F}(z)}{\partial z^i} + \bar{e}_a^{\bar{i}}(\bar{z}) \frac{\partial \bar{\mathcal{F}}(\bar{z})}{\partial \bar{z}^{\bar{i}}} \right) \right]^{-1}, \tag{12.1.4}$$

where

$$\begin{aligned}
\mathcal{F}(z) &\equiv \mathcal{F}\left(\frac{X(z)}{X^0(z)}\right) = (X^0(z))^{-2} F(X(z)); \\
e_a^i(z) &\equiv e_a^i\left(\frac{X(z)}{X^0(z)}\right), \quad \bar{e}_a^{\bar{i}}(\bar{z}) = \overline{e_a^i(z)}.
\end{aligned} \tag{12.1.5}$$

Thus, Eq. (12.1.2) can be rewritten as follows (for simplicity's sake, we omit notation of dependence on $(\psi, \bar{\psi})$):

$$C = e^K \mathcal{W} = -i \frac{[3e^{-2} (\partial e)^2 \partial \mathcal{F} - e^{-1} (\partial^2 e) \partial \mathcal{F} - 3e^{-1} (\partial e) \partial^2 \mathcal{F} + \partial^3 \mathcal{F}]}{\bar{X}^0 \left[2(\mathcal{F} - \bar{\mathcal{F}}) - \left(\frac{X^1}{\bar{X}^0} - \frac{\bar{X}^1}{\bar{X}^0} \right) (e^{-1} \partial \mathcal{F} + \bar{e}^{-1} \partial \bar{\mathcal{F}}) \right]}, \tag{12.1.6}$$

which can be also further elaborated by recalling Eq. (4.2.1).

Eq. (12.1.6) holds *globally* for a generic $n_V = 1$ SK geometry, with no Kähler gauge-fixing, with the only condition of $e \neq 0$ (and $X^0 \neq 0$). By specializing it for the 1-dim. SK geometry endowing the moduli space of (mirror) Fermat

CY_3 s, one has to replace the (k -indexed) quantities X^0 , X^1 , e and $\mathcal{F}$ with the first and second component of Eq. (6.22), with consequent Eq. (4.2.1), and with Eq. (6.42), respectively. Thence, one has to use the results obtained for $k = 5, 6, 8, 10$ in Sects. 7-10, which however converge only for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$, and have mainly considered in the *LG limit* $\psi \rightarrow 0$.

II) Use Eq. (12.1.1) and the relation $a_4 = \mathcal{W}^{-1}$ (given by the first of definitions (4.2.8)), where a_4 is the coefficient of the fourth derivative in the PF Eq. (4.2.7). It is important to notice that the relation $a_4 = \mathcal{W}^{-1}$ was obtained in [58], where the relation

$$D_\alpha U_\beta = C_{\alpha\beta\gamma} g^{\gamma\bar{\gamma}} \bar{U}_{\bar{\gamma}} \quad (12.1.7)$$

holds (see the second of Eqs. (8) of [58]), rather than the second of Eqs. (4.1.2), consistent with the conventions we use. This means that the C_{ijk} used in the present paper is nothing but $-i$ times the C_{ijk} used in [58]. In the case $n_V = 1$, by taking into account such a shift of phase and using (the $n_V = 1$ case of) Eq. (12.1.4), one finally gets the relation

$$C = -ie^K a_4^{-1} = - \left\{ a_4 |X^0|^2 \left[2(\mathcal{F} - \bar{\mathcal{F}}) - \left(\frac{X^1}{X^0} - \frac{\bar{X}^1}{\bar{X}^0} \right) (e^{-1} \partial \mathcal{F} + \bar{e}^{-1} \bar{\partial} \bar{\mathcal{F}}) \right] \right\}^{-1}. \quad (12.1.8)$$

As Eq. (12.1.6), Eq. (12.1.8) holds *globally* for a generic $n_V = 1$ SK geometry, with no Kähler gauge-fixing, with the only condition of $e \neq 0$ (and $X^0 \neq 0$). Let us specialize it for 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s. In order to do this, we need to introduce the Fermat classifying parameter k , and recall Eq. (6.24) (converging for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$), the first of Eqs. (6.35) and the second of Eqs. (5.2.8), thus obtaining (with the use of Tables 1 and 2):

$$C_k = -ie^{K_k} \frac{1}{\psi^{5-k} - \psi^5} = -\frac{i}{(2\pi)^6} \frac{(Ord(G_k))^2}{C_{k,0}^2 F_{k,11}} \frac{e^{\tilde{K}_k}}{1 - \psi^k} \psi^{k-5} = \quad (12.1.9)$$

$$= -\frac{i}{(2\pi)^6} \frac{(Ord(G_k))^2}{C_{k,0}^2 F_{k,11}} \frac{\psi^{k-5}}{1 - \psi^k} \left\{ 1 + \frac{1}{C_{k,0}^2 F_{k,11}} \left[2C_{k,0} \sum_{n=2}^{\infty} C_{k,n-1} F_{k,1n} Re(\psi^{n-1}) + \sum_{m,n=2}^{\infty} C_{k,m-1} C_{k,n-1} F_{k,mn} \psi^{m-1} \bar{\psi}^{n-1} \right] \right\}^{-1}; \quad (12.1.10)$$

notice that Eq. (12.1.9) holds globally, whereas Eq. (12.1.10) converges for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$ (as given by Eq. (6.24)). By recalling that $k = 5, 6, 8, 10$, in the *LG limit* $\psi \rightarrow 0$ such an expression can be approximated as follows:

$$\begin{aligned} \lim_{\psi \rightarrow 0} C_k &\approx -\frac{i}{(2\pi)^6} \frac{(Ord(G_k))^2}{C_{k,0}^2 F_{k,11}} \frac{\psi^{k-5}}{1 - \psi^k} \left\{ 1 - \frac{1}{C_{k,0}^2 F_{k,11}} \left[2C_{k,0} \sum_{n=2}^{\infty} C_{k,n-1} F_{k,1n} Re(\psi^{n-1}) + \sum_{m,n=2}^{\infty} C_{k,m-1} C_{k,n-1} F_{k,mn} \psi^{m-1} \bar{\psi}^{n-1} \right] \right\} \approx \\ &\approx -\frac{i}{(2\pi)^6} \frac{(Ord(G_k))^2}{C_{k,0}^2 F_{k,11}} \psi^{k-5} \longrightarrow -\frac{i}{(2\pi)^6} \frac{(Ord(G_k))^2}{C_{k,0}^2 F_{k,11}} \delta_{k,5}. \end{aligned} \quad (12.1.11)$$

Thus, in the 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s the *LG limit* of the Yukawa coupling function is holomorphic, with the leading term $\sim \mathcal{O}(\psi^{k-5})$, as given by Eq. (12.1.11).

III) Use Eq. (11.2.2.4). Clearly, it holds not only for the rescaled holomorphic central charge $\widetilde{W}$ (introduced in the framework of Fermat CY_3 - compactifications by the second of definitions (6.35)), but for a generic superpotential W of a 1-dim. SK geometry.

Apparently, Eq. (11.2.2.4) yields a way to compute C only along the *locus* $DW \neq 0$ of $n_V = 1$ SK manifolds, *i.e.* only away from $\frac{1}{2}$ -BPS critical points of V_{BH} . Instead, due to the fact that the fundamental relations (4.1.2) of SK geometry do hold globally, the approach based on Eq. (11.2.2.4) (which is nothing but the $n_V = 1$ case of the second of Eqs. (4.1.2)) has a global validity, yielding that

$$C = -ig \frac{D^2 W}{DW} = -i (\bar{\partial} \partial K) \frac{\left\{ \partial^2 + \partial^2 K + 2\partial K \partial + (\partial K)^2 - [\partial \ln (\partial \bar{\partial} K)] (\partial + \partial K) \right\} W}{[\bar{\partial} + (\bar{\partial} K)] \bar{W}}, \quad (12.1.12)$$

where Eqs. (2.2.3) and (2.2.5) have been used. Notice that the charge-dependence of DW and $D^2 W$ vanishes in the ratio $\frac{D^2 W}{DW}$ (C is charge-independent). As Eqs. (12.1.6) and (12.1.8), Eq. (12.1.12) holds *globally* for a generic $n_V = 1$ SK geometry, with *a priori* no Kähler gauge-fixing. In order to specialize it for the 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s, we need to introduce the Fermat classifying parameter k , and furthermore to recall the general treatment given in Sect. 6, and use the results obtained for $k = 5, 6, 8, 10$ in Sects. 7-10, which however converge only for $|\psi| < 1$, $0 \leq \arg(\psi) < \frac{2\pi}{k}$, and have mainly considered in the *LG limit* $\psi \rightarrow 0$. By doing this, one finally gets

$$\begin{aligned} k = 5 : \lim_{\psi \rightarrow 0} C_5 &\approx -2i (\sqrt{5} - 2) \frac{C_{5,1} C_{5,2}}{C_{5,0}^2}; \\ k = 6 : \lim_{\psi \rightarrow 0} C_6 &\approx 2i \frac{C_{6,1} C_{6,3}}{C_{5,0}^2} \psi; \\ k = 8 : \lim_{\psi \rightarrow 0} C_8 &\approx 16i (3 - 2\sqrt{2}) \frac{C_{8,2} C_{8,4}}{C_{8,0}^2} \psi^3; \\ k = 10 : \lim_{\psi \rightarrow 0} C_{10} &= 48i (\sqrt{5} - 2) \frac{C_{10,2} C_{10,6}}{C_{10,0}^2} \psi^5. \end{aligned} \quad (12.1.13)$$

In order for the LG limits (12.1.11) and (12.1.13) to coincide, it is easy to realize that $a_{4,k}$ must be normalized as follows

$$\begin{aligned} a_{4,k} &\longrightarrow \frac{(Ord(G_k))^2}{(2\pi)^6 F_{k,11} \mathcal{A}_k} a_{4,k}; \\ \mathcal{A}_5 &\equiv 2C_{5,1} C_{5,2} (\sqrt{5} - 2); \\ \mathcal{A}_6 &\equiv -2C_{6,1} C_{6,3}; \\ \mathcal{A}_8 &\equiv -16C_{8,2} C_{8,4} (3 - 2\sqrt{2}); \\ \mathcal{A}_{10} &\equiv -48C_{10,2} C_{10,6} (5 - \sqrt{2}). \end{aligned} \quad (12.1.14)$$

In order to preserve the set of solutions of the PF Eq. (5.2.8) and the properties of the coefficients $a_{n,k}$ (as given by Eqs. (4.2.8)-(4.2.11)), it is clear that the normalization (12.1.14) yields the multiplication of PF Eq. (5.2.8) by the overall constant $\frac{(Ord(G_k))^2}{(2\pi)^6 F_{k,11} \mathcal{A}_k}$. Thus, one can state that the compatibility between the approaches II and III to

the computation of the Yukawa coupling function C_k in the LG limit $\psi \rightarrow 0$ of the (k -parameterized) 1-dim. SK geometry of the moduli space of (mirror) Fermat CY_3 s yields the following (*irrelevant*, because overall and constant) normalization of the PF Eq. (5.2.8)

$$\sum_{n=0}^4 a_{n,k}(\psi) \partial^n V_h(\psi) = 0, \quad (12.1.15)$$

$$a_{n,k}(\psi) \equiv \frac{(Ord(G_k))^2}{(2\pi)^6 F_{k,11} \mathcal{A}_k} \left[-\sigma_n \psi^{n+1} + (-1)^n \tau_{n,k} \psi^{n+1-k} \right],$$

where the k -dependent constants $Ord(G_k)$, $F_{k,11}$, $\mathcal{A}_k$, σ_n and $\tau_{n,k}$ are given by Table 3, Eqs. (7.1.2), (8.1.2), (9.1.2), (10.1.2), (12.1.14), and Tables 1 and 2, respectively. Finally, it can be checked that the use of approach I does not give raise to any other problem of consistency/compatibility with the approaches II and III in the framework (of the LG limit) of 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s.

12.2 Check of Stability Conditions

By recalling Eqs. (12.1.8) and (12.1.9), and the first of definitions (6.35), from the correct normalization (12.1.14) of $a_{n,k}$ we can achieve the exact, k -parametrized global formula of the Yukawa coupling function C_k for the (1-dim. SK geometry endowing the moduli space of mirror) Fermat CY_3 s:

$$C_k = -ie^k (a_{4,k})^{-1} = -i \frac{\mathcal{A}_k}{C_{k,0}^2} \frac{\psi^{k-5}}{1-\psi^k} e^{\tilde{K}_k}. \quad (12.2.1)$$

Let us now reconsider Eq. (3.2.4); by using Eq. (12.2.1), we can compute the exact, k -parametrized global formula of the covariant derivative of the Yukawa coupling function of 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s:

$$DC_k = \partial C_k + \left[\left(\partial \tilde{K}_k \right) - 3 \partial \ln \left(\bar{\partial} \partial \tilde{K}_k \right) \right] C_k = -i \frac{\mathcal{A}_k}{C_{k,0}^2} \left[\frac{5\psi^4 + (k-5)\psi^{4-k}}{\psi^{5-k} - \psi^5} + 2 \left(\partial \tilde{K}_k \right) - 3 \left(\frac{\bar{\partial} \partial^2 \tilde{K}_k}{\bar{\partial} \partial \tilde{K}_k} \right) \right] \frac{\psi^{k-5}}{1-\psi^k} e^{\tilde{K}_k}. \quad (12.2.2)$$

We now have all the ingredients to check the stability conditions for non-BPS $Z \neq 0$ and non-BPS $Z = 0$ critical points of V_{BH} in the *LG limit* of the 1-dim. SK geometry endowing the moduli space of (mirror) Fermat CY_3 s. By recalling Eqs. (3.2.2.8) and (3.2.3.7), using Eqs. (12.2.1) and (12.2.2), the results obtained for the LG limit of (the moduli space of mirror) Fermat CY_3 s for $k = 5, 6, 8, 10$ in Sects. 7, 8, 9 and 10 respectively, yield that:

1) for $k = 5, 8$ the condition (3.2.2.8) for $\psi \approx 0$ to be a *stable* non-BPS $Z \neq 0$ critical point of $V_{BH,k=5,8}$ (and thus to be an *attractor* in strict sense) is satisfied, consistently with the results obtained in Subsects. 7.3-7.4 and 9.3-9.4, respectively;

2) for $k = 6, 10$ the condition (3.2.3.7) for $\psi \approx 0$ to be a *stable* non-BPS $Z = 0$ critical point of $V_{BH,k=6,10}$ (and thus to be an *attractor* in strict sense) is *not* satisfied, consistently with the results obtained in Subsects. 8.3-8.4 and 10.3-10.4, respectively.

12.3 The Exact Holomorphic Yukawa Coupling Function of Fermat CY_3 s

Having performed a self-consistent treatment of Yukawa coupling function C in the LG limit, we can now reconsider the relation $a_4 = \mathcal{W}^{-1}$ (given by the first of definitions (4.2.8)); indeed, from the correct normalization (12.1.14) of $a_{n,k}$, we can achieve the exact, k -parametrized global formula of the holomorphic Yukawa coupling function $\mathcal{W}_k$ for the (1-dim. SK geometry endowing the moduli space of mirror) Fermat CY_3 s:

$$\mathcal{W}_k(\psi) = [a_{4,k}(\psi)]^{-1} = \frac{(2\pi)^6 F_{k,11} \mathcal{A}_k}{(\text{Ord}(G_k))^2} \frac{\psi^{k-5}}{1 - \psi^k}. \quad (12.3.1)$$

By using such an exact global formula, the evaluation of $\mathcal{W}_k$ near the three species of regular singular points of PF Eq. (12.1.15) of Fermat CY_3 s yields:

$$LG \text{ limit} : \lim_{\psi \rightarrow 0} \mathcal{W}_k(\psi) = \frac{(2\pi)^6 F_{k,11} \mathcal{A}_k}{(\text{Ord}(G_k))^2} \lim_{\psi \rightarrow 0} \frac{\psi^{k-5}}{1 - \psi^k} \approx \frac{(2\pi)^6 F_{k,11} \mathcal{A}_k}{(\text{Ord}(G_k))^2} \psi^{k-5} \longrightarrow \frac{(2\pi)^6 F_{k,11} \mathcal{A}_k}{(\text{Ord}(G_k))^2} \delta_{k,5}; \quad (12.3.2)$$

$$Conifold \text{ limit} : \left| \lim_{\psi^k \rightarrow 1} \mathcal{W}_k(\psi) \right| = \infty; \quad (12.3.3)$$

$$Large \text{ complex structure modulus limit} : \lim_{\psi \rightarrow \infty} \mathcal{W}_k(\psi) \approx -\frac{(2\pi)^6 F_{k,11} \mathcal{A}_k}{(\text{Ord}(G_k))^2} \psi^{-5} \longrightarrow 0. \quad (12.3.4)$$

Remark

A final comment concerns the vanishing of C_8 (and $\mathcal{W}_8$) in the LG limit, as yielded by Eqs. (12.1.11) (and (12.3.2)). This might seem in contrast with the result, obtained in Sect. 9, that for $k = 8$ only $\frac{1}{2}$ -BPS and non-BPS $Z \neq 0$ LG critical points of V_{BH} (both stable) exist, depending on the supporting BH charge configuration(s). For non-BPS $Z \neq 0$ attractors, this implies that $C_{8,non-BPS,Z \neq 0} \neq 0$, as given for a generic 1-dim. SK geometry by Eq. (2.2.10). The contrast with the result (12.1.11) can be simply explained by pointing out that for the (mirror) octic $\mathcal{M}'_8$ the exact LG point $\psi = 0$ cannot be a non-BPS, $Z \neq 0$ critical point of V_{BH} . Rather, such a kind of critical point turns out to be realized for $\psi \approx 0$, *i.e.* in a suitable neighbourhood of the LG point. Indeed, as remarked below Eq. (7.2.1) and as understood in all the treatment of the present paper, the actual BH charges satisfying the $\mathcal{N} = 2$, $d = 4$ AE in the LG limit are very close to the found one, and also the critical value of ψ may not be zero (*i.e.* the *exact* LG point), rather it may belong to a suitable neighbourhood of the LG point.

13 Conclusions and Outlook

In the present work we investigated non-degenerate extremal BH attractors near the so-called LG point $\psi = 0$ (herein named *LG attractors*) of the moduli space (with $\dim_{\mathbb{C}} = 1$) of the class of (mirror) Fermat Calabi-Yau threefolds. We found the BH charge configurations supporting $\psi \approx 0$ to be a critical point of the real, positive-definite “effective BH potential” V_{BH} defined in Eq. (2.2.1).

In order to do this, we exploited two different approaches:

i) “criticality condition” approach: we solved at $\psi \approx 0$ the 2 real *criticality conditions* of V_{BH} , corresponding in the 1-modulus case to the real and imaginary part of the *Attractor Eq.* (2.2.2) (see Sects. 7-10);

ii) “SK geometrical identities” approach: we evaluated at $\psi \approx 0$ the 4 real fundamental identities (11.1.1) of 1-modulus SK geometry at the geometrical *loci* corresponding to the various species of critical points of V_{BH} (see Sect. 11).

We found that the results of two such solving approaches do coincide, in spite of the different number of real Eqs. involved in approaches 1 and 2. The equivalence of the above-mentioned approaches to find the critical points of V_{BH} (and the BH charge configurations supporting them) is explicit proof of the fact that the relations (11.1.1) actually are *identities* and not equations, *i.e.* that, for any point of the moduli space at which we evaluate them, they do *not* give any constraint on the charge configuration.

It is worth pointing out that the “criticality condition” approach had been previously exploited in literature only for the following cases:

i.a) *mirror quintic* ($k = 5$) in [9], where however peculiar *Ansätze* (on the BH charge configuration and on ψ in the neighbourhood of the LG point) were used, implying a certain loss of generality;

i.b) *mirror sextic* ($k = 6$) in [10].

On the other hand, the “SK geometrical identities” approach (and its equivalence with the “criticality condition” one) had been hitherto exploited only in [13]; in such a Ref., the mirror *quintic* was considered within the same simplifying *Ansätze* formulated in [9], obtaining a complete agreement with the results of [9].

As a by-product of our computations, we extended the results of [9] and [13] to full generality (see Sect. 7). Moreover, we found that the analysis of the stability of $\psi \approx 0$ as a non-BPS, $Z = 0$ critical point of V_{BH} in the mirror *sextic*, performed in Sect. 7 of [10], suffers from some problems of inconsistency. Indeed, in [10] it was found that the LG point (supported by a certain BH charge configuration characterizing it as a non-BPS, $Z = 0$ critical point of V_{BH}) is *stable* (minimum of V_{BH}). Instead, our computations (see Sect. 8), which carefully took into account the relevant orders in ψ and $\bar{\psi}$ in the truncation of the series expansion around $\psi \approx 0$, allow us to conclude that, *for the same supporting BH charge configuration*, the LG point is *unstable* (namely, a saddle point of V_{BH}).

We also checked the stability of $\psi \approx 0$ in two ways:

1) by inspecting the Hessian matrix of V_{BH} in correspondence to the various BH charge configurations supporting the LG point to be a critical point of V_{BH} (see Subsects. 7.3-7.4, 8.3-8.4, 9.3-9.4 and 10.3-10.4 for $k = 5, 6, 8, 10$, respectively);

2) by using the stability conditions for non-BPS $Z \neq 0$ and non-BPS $Z = 0$ critical points of V_{BH} in 1-modulus SK geometry, respectively obtained in Subsubsects. 3.2.2 and 3.2.3.

A sketchy summary of our results is given by the following Table:

The stability of $\psi \approx 0$ as a $\frac{1}{2}$ -BPS attractor agrees with the known results from general analysis of SK geometry of scalar manifolds in $\mathcal{N} = 2$, $d = 4$ supergravity coupled to n_V Abelian vector multiplets [39, 15, 16].

For non-BPS LG critical points of V_{BH} the central charge Z turns out to be a crucial quantity for stability. Indeed,

$k \longrightarrow$	5	6	8	10
$\frac{1}{2}$ -BPS	<i>stable</i> , 1 charge config.	<i>stable</i> , 1 charge config.	<i>stable</i> , 1 charge config.	<i>stable</i> , 1 charge config.
non-BPS, $Z \neq 0$	<i>stable</i> , 3 charge configs.	—	<i>stable</i> , 3 charge configs.	—
non-BPS, $Z = 0$	—	<i>unstable</i> , 1 charge config.	—	<i>unstable</i> , 1 charge config.

Table 5: **Species and stability of the LG critical point $\psi \approx 0$ of V_{BH} in the moduli space of (mirror) Fermat CY_3 s**

regardless of the kind of BH charge configuration supporting them, the non-BPS, $Z \neq 0$ LG attractors, when they exist, are found to be stable (minima of V_{BH}). On the other hand, the non-BPS, $Z = 0$ LG attractors, when they exist, are found to be unstable (saddle points of V_{BH}).

It is interesting to compare such a result to what happens in the large volume limit of CY_3 -compactifications (of Type IIA superstrings). Indeed, in such a framework (with a generic number n_V of complex structure moduli) in [9] it was shown that the stability of non-BPS, $Z \neq 0$ critical points of V_{BH} (and therefore their actual attractor behaviour) within a certain supporting BH charge configuration, crucially depends on the possible vanishing of p^0 , *i.e.* of the asymptotical magnetic flux of the graviphoton field strength, whose microscopical interpretation corresponds to a $D6$ -brane wrapping p^0 times a 3-cycle of the considered CY_3 . Nevertheless, also in such a context in the 1-modulus case the non-BPS, $Z \neq 0$ critical points of V_{BH} are always stable, and therefore they are attractors in a strict sense.

Furthermore, one can also observe that all Fermat CY_3 s admit *only one* kind of non-BPS LG attractors, either with $Z \neq 0$ or with $Z = 0$; for the allowed values of the classifying Fermat parameter $k = 5, 6, 8, 10$, one gets the “pattern” shown in Table 5 above.

Once again, such a feature is exhibited also by the large volume limit of CY_3 -compactifications (of Type IIA superstrings), whose related SK geometry is characterized by cubic holomorphic prepotentials; indeed, it can be explicitly computed that the 1-modulus prepotential (3.2.3.11) corresponding to the homogeneous symmetric SK manifold $\frac{SU(1,1)}{U(1)}$ (see the remark at the end of Subsubsect. 3.2.3, and [22] and Refs. therein), admits $\frac{1}{2}$ -BPS and non-BPS, $Z \neq 0$ critical points of V_{BH} (both stable) only.

It is worth mentioning that the fourth order linear ordinary differential Picard-Fuchs equations of Fermat CY_3 s (5.2.8) (or equivalently (12.1.15)) (specified by Tables 1 and 2) exhibit other two species of regular singular points, namely the k -th roots of unity ($\psi^k = 1$, the so-called *conifold points*) and the *point at infinity* $\psi \longrightarrow \infty$ in the moduli space, corresponding to the so-called *large complex structure modulus limit*. It would be interesting to solve criticality conditions for V_{BH} near such regular singular points, *i.e.* to investigate *extremal BH conifold attractors* and *extremal BH large complex structure attractors* in the moduli space of 1-modulus (Fermat) CY_3 s, also in view of recent investigations of extremal BH attractors in specific examples of 2-moduli CY_3 -compactifications [25].

When CY_3 -compactifications with more than one complex structure deformation modulus are considered, it is clear

that interesting situations might arise other than the ones present at 1-modulus level. Indeed, differently from what has been studied so far [25], in such frameworks all three species of extremal BH (LG) attractors (namely $\frac{1}{2}$ -BPS, non-BPS $Z \neq 0$ and non-BPS $Z = 0$) should exist, each typology being supported by distinct, zero-overlapping BH charge configurations. *Ça va sans dire* that such an issue deserves more investigation and analyzing efforts.

Finally, it is worth spending a few words concerning the instability of non-BPS, $Z = 0$ (LG) attractors in the 1-modulus case. It would be intriguing to extend to such a framework the same conjecture formulated in [24]. In Sect. 5 of such a Ref., in the framework of (the large volume limit of CY_3 -compactifications leading to) the peculiarly symmetric case of cubic *stu* model, it was argued that the instability of the considered non-BPS attractors might be only apparent, since such attractors might correspond to multi-centre stable attractor solutions, whose stable nature should be “resolved” only at sufficiently small distances. As mentioned, it would be interesting to extend such a conjecture to the non-BPS, $Z = 0$ (LG) attractors, also in relation to the possible existence of *non-BPS lines of marginal stability* [65, 66].

Acknowledgments

It is a pleasure to acknowledge proofreading by Mrs. Suzy Vascotto and Mrs. Helen Webster.

A. Y. would like to thank the INFN Frascati National Laboratories for the kind hospitality extended to him during the work for the present paper.

The work of S.B. has been supported in part by the European Community Human Potential Program under contract MRTN-CT-2004-005104 “*Constituents, fundamental forces and symmetries of the Universe*”.

The work of S.F. has been supported in part by the European Community Human Potential Program under contract MRTN-CT-2004-005104 “*Constituents, fundamental forces and symmetries of the Universe*”, in association with INFN Frascati National Laboratories and by D.O.E. grant DE-FG03-91ER40662, Task C.

The work of A.M. has been supported by a Junior Grant of the “*Enrico Fermi*” Centre, Rome, in association with INFN Frascati National Laboratories.

The work of A.Y. was supported in part by the grants NFSAT-CRDF ARPI-3328-YE-04 and INTAS-05-7928.

References

- [1] S. Ferrara, R. Kallosh and A. Strominger, *$\mathcal{N}=2$ Extremal Black Holes*, Phys. Rev. **D52**, 5412 (1995), [hep-th/9508072](#).
- [2] A. Strominger, *Macroscopic Entropy of $\mathcal{N}=2$ Extremal Black Holes*, Phys. Lett. **B383**, 39 (1996), [hep-th/9602111](#).
- [3] S. Ferrara and R. Kallosh, *Supersymmetry and Attractors*, Phys. Rev. **D54**, 1514 (1996), [hep-th/9602136](#).
- [4] S. Ferrara and R. Kallosh, *Universality of Supersymmetric Attractors*, Phys. Rev. **D54**, 1525 (1996), [hep-th/9603090](#).
- [5] A. Sen, *Black Hole Entropy Function and the Attractor Mechanism in Higher Derivative Gravity*, JHEP **09**, 038 (2005), [hep-th/0506177](#).
- [6] K. Goldstein, N. Iizuka, R. P. Jena and S. P. Trivedi, *Non-Supersymmetric Attractors*, Phys. Rev. **D72**, 124021 (2005), [hep-th/0507096](#).
- [7] A. Sen, *Entropy Function for Heterotic Black Holes*, JHEP **03**, 008 (2006), [hep-th/0508042](#).
- [8] R. Kallosh, *New Attractors*, JHEP **0512**, 022 (2005), [hep-th/0510024](#).
- [9] P. K. Tripathy and S. P. Trivedi, *Non-Supersymmetric Attractors in String Theory*, JHEP **0603**, 022 (2006), [hep-th/0511117](#).
- [10] A. Giryavets, *New Attractors and Area Codes*, JHEP **0603**, 020 (2006), [hep-th/0511215](#).
- [11] K. Goldstein, R. P. Jena, G. Mandal and S. P. Trivedi, *A C-Function for Non-Supersymmetric Attractors*, JHEP **0602**, 053 (2006), [hep-th/0512138](#).
- [12] M. Alishahiha and H. Ebrahim, *Non-supersymmetric attractors and entropy function*, JHEP **0603**, 003 (2006), [hep-th/0601016](#).
- [13] R. Kallosh, N. Sivanandam and M. Soroush, *The Non-BPS Black Hole Attractor Equation*, JHEP **0603**, 060 (2006), [hep-th/0602005](#).
- [14] B. Chandrasekhar, S. Parvizi, A. Tavanfar and H. Yavartanoo, *Non-supersymmetric attractors in R^2 gravities*, [hep-th/0602022](#).
- [15] S. Bellucci, S. Ferrara and A. Marrani, *On some properties of the Attractor Equations*, Phys. Lett. **B635**, 172 (2006), [hep-th/0602161](#).

- [16] S. Bellucci, S. Ferrara and A. Marrani, *Supersymmetric Mechanics. Vol.2: The Attractor Mechanism and Space-Time Singularities* (LNP **701**, Springer-Verlag, Heidelberg, 2006).
- [17] G. L. Cardoso, D. Lüst and J. Perz, *Entropy Maximization in the presence of Higher-Curvature Interactions*, JHEP **05**, 028 (2006), [hep-th/0603211](#).
- [18] B. Sahoo and A. Sen, *Higher-derivative corrections to non-supersymmetric extremal black holes*, [hep-th/0603149](#).
- [19] S. Ferrara and R. Kallosh, *On $\mathcal{N}=8$ attractors*, Phys. Rev. D **73**, 125005 (2006), [hep-th/0603247](#).
- [20] M. Alishahiha and H. Ebrahim, *New attractor, Entropy Function and Black Hole Partition Function*, [hep-th/0605279](#).
- [21] S. Ferrara and M. Günaydin, *Orbits and attractors for $\mathcal{N}=2$ Maxwell-Einstein supergravity theories in five dimensions*, [hep-th/0606108](#).
- [22] S. Bellucci, S. Ferrara, M. Günaydin and A. Marrani, *Charge Orbits of Symmetric Special Geometries and Attractors*, Int. J. Mod. Phys. **A21**, 5043 (2006), [hep-th/0606209](#).
- [23] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen and S. P. Trivedi, *Rotating Attractors*, [hep-th/0606244](#).
- [24] R. Kallosh, N. Sivanandam and M. Soroush, *Exact Attractive non-BPS STU Black Holes*, [hep-th/0606263](#).
- [25] P. Kaura and A. Misra, *On the Existence of Non-Supersymmetric Black Hole Attractors for Two-Parameter Calabi-Yau's and Attractor Equations*, [hep-th/0607132](#).
- [26] G. L. Cardoso, V. Grass, D. Lüst and J. Perz, *Extremal non-BPS Black Holes and Entropy Extremization*, [hep-th/0607202](#).
- [27] J. F. Morales and H. Samtleben, *Entropy Function and Attractors for AdS Black Holes*, [hep-th/0608044](#).
- [28] B. Chandrasekhar, H. Yavartanoo and S. Yun, *Non-Supersymmetric Attractors in BI black holes*, [hep-th/0611240](#).
- [29] L. Andrianopoli, R. D'Auria, S. Ferrara and M. Trigiante, *Extremal Black Holes in Supergravity*, in : *"String Theory and Fundamental Interactions"*, M. Gasperini and J. Maharana eds. (LNP, Springer, Berlin-Heidelberg, 2007), [hep-th/0611345](#).
- [30] A.E. Mosaffa, S. Randjbar-Daemi and M.M. Sheikh-Jabbari, *Non-Abelian Magnetized Blackholes and Unstable Attractors*, [hep-th/0612181](#).
- [31] G.L. Cardoso, B. de Wit and S. Mahapatra, *Black hole entropy functions and attractor equations*, [hep-th/0612225](#).

- [32] R. D'Auria, S. Ferrara and M. Trigiante, *Critical points of the Black-Hole potential for homogeneous special geometries*, **hep-th/0701090**.
- [33] S. Bellucci, S. Ferrara and A. Marrani, *Attractor Horizon Geometries of Extremal Black Holes*, Contribution to the Proceedings of the XVII SIGRAV Conference, 4–7 September 2006, Turin, Italy, **hep-th/0702019**.
- [34] L. Andrianopoli, R. D'Auria, S. Ferrara and M. Trigiante, *Black Hole Attractors in $\mathcal{N}=1$ Supergravity*, **hep-th/0703178**.
- [35] K. Saraikin and C. Vafa, *Non-Supersymmetric Black Holes and Topological Strings*, **hep-th/0703214**.
- [36] J. D. Bekenstein, Phys. Rev. **D7**, 2333 (1973) $\diamond$ S. W. Hawking, Phys. Rev. Lett. **26**, 1344 (1971); in: “*Black Holes*” (*Les Houches 1972*), C. DeWitt and B. S. DeWitt eds. (Gordon and Breach, New York, 1973) $\diamond$ S. W. Hawking, Nature **248**, 30 (1974) $\diamond$ S. W. Hawking, Comm. Math. Phys. **43**, 199 (1975).
- [37] G. W. Gibbons and C. M. Hull, *A Bogomol'ny Bound for General Relativity and Solitons in $\mathcal{N}=2$ Supergravity*, Phys. Lett. **B109**, 190 (1982).
- [38] R. Arnowitt, S. Deser and C. W. Misner, *The Dynamics of General Relativity*, in : “*Gravitation: an Introduction to Current Research*”, L. Witten ed. (Wiley, New York, 1962).
- [39] S. Ferrara, G. W. Gibbons and R. Kallosh, *Black Holes and Critical Points in Moduli Space*, Nucl. Phys. **B500**, 75 (1997), **hep-th/9702103**.
- [40] S. Ferrara and M. Günaydin, *Orbits of Exceptional Groups, Duality and BPS States in String Theory*, Int. J. Mod. Phys. **A13**, 2075 (1998), **hep-th/9708025**.
- [41] A. Forsyth, *Theory of Differential Equations, Vol. 4* (Dover Publications, New York, 1959).
- [42] A. Strominger and E. Witten, *New Manifolds for Superstring Compactification*, Commun. Math. Phys. **101**, 341 (1985).
- [43] P. Candelas, X. C. De La Ossa, P. S. Green and L. Parkes, *A Pair of Calabi-Yau Manifolds as an Exactly Soluble Superconformal Theory*, Nucl. Phys. **B359**, 21 (1991).
- [44] P. Candelas, X. C. De La Ossa, P. S. Green and L. Parkes, *An Exactly Soluble Superconformal Theory from a Mirror Pair of Calabi-Yau Manifolds*, Phys. Lett. **B258**, 118 (1991).
- [45] A. C. Cadavid and S. Ferrara, *Picard-Fuchs Equations and the Moduli Space of Superconformal Field Theories*, Phys. Lett. **B267**, 193 (1991).
- [46] A. Klemm and S. Theisen, *Considerations of One Modulus Calabi-Yau Compactifications: Picard-Fuchs Equations, Kähler Potentials and Mirror Maps*, Nucl. Phys. **B389**, 153 (1993), **hep-th/9205041**.

- [47] A. Font, *Periods and Duality Symmetries in Calabi-Yau Compactifications*, [hep-th/9203084](#).
- [48] A. Ceresole, R. D'Auria and S. Ferrara, *The Symplectic Structure of $\mathcal{N}=2$ SUGRA and Its Central Extension*, Talk given at ICTP Trieste Conference on Physical and Mathematical Implications of Mirror Symmetry in String Theory, Trieste, Italy, 5-9 June 1995, Nucl. Phys. Proc. Suppl. **46** (1996), [hep-th/9509160](#).
- [49] A. Ceresole, R. D'Auria, S. Ferrara and A. Van Proeyen, *Duality Transformations in Supersymmetric Yang-Mills Theories Coupled to Supergravity*, Nucl. Phys. **B444**, 92 (1995), [hep-th/9502072](#).
- [50] J. Bagger and E. Witten, *The Gauge Invariant Supersymmetric Nonlinear Sigma Model*, Phys. Lett. **B118**, 103 (1982).
- [51] B. Craps, F. Roose, W. Troost and A. Van Proeyen, *The Definitions of Special Geometry*, [hep-th/9606073](#).
- [52] B. Craps, F. Roose, W. Troost and A. Van Proeyen, *What is Special Kähler Geometry?*, Nucl. Phys. **B503**, 565 (1997), [hep-th/9703082](#).
- [53] L. Castellani, R. D'Auria and S. Ferrara, *Special Geometry without Special Coordinates*, Class. Quant. Grav. **7**, 1767 (1990) $\diamond$ L. Castellani, R. D'Auria and S. Ferrara, *Special Kähler Geometry: an Intrinsic Formulation from $\mathcal{N}=2$ Space-Time Supersymmetry*, Phys. Lett. **B241**, 57 (1990).
- [54] R. D'Auria, S. Ferrara and P. Fré, *Special and Quaternionic Isometries: General Couplings in $\mathcal{N}=2$ Supergravity and the Scalar Potential*, Nucl. Phys. **B359**, 705 (1991) $\diamond$ L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara, P. Fré and T. Magri, *$\mathcal{N}=2$ Supergravity and $\mathcal{N}=2$ Super Yang-Mills Theory on General Scalar Manifolds : Symplectic Covariance, Gaugings and the Momentum Map*, J. Geom. Phys. **23**, 111 (1997), [hep-th/9605032](#) $\diamond$ L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara and P. Fré, *General Matter Coupled $\mathcal{N}=2$ Supergravity*, Nucl. Phys. **B476**, 397 (1996), [hep-th/9603004](#).
- [55] S. Ferrara, M. Bodner and A. C. Cadavid, *Calabi-Yau Supermoduli Space, Field Strength Duality and Mirror Manifolds*, Phys. Lett. **B247**, 25 (1990).
- [56] E. Cremmer, C. Kounnas, A. Van Proeyen, J.P. Derendinger, S. Ferrara, B. de Wit and L. Girardello, *Vector Multiplets Coupled To $\mathcal{N}=2$ Supergravity: Superhiggs Effect, Flat Potentials And Geometric Structure*, Nucl. Phys. **B250**, 385 (1985).
- [57] E. Cremmer and A. Van Proeyen, *Classification of Kähler Manifolds in $\mathcal{N}=2$ Vector Multiplet Supergravity Couplings*, Class. Quant. Grav. **2**, 445 (1985).
- [58] S. Ferrara and J. Louis, *Flat Holomorphic Connections and Picard-Fuchs Identities from $\mathcal{N}=2$ Supergravity*, Phys. Lett. **B278**, 240 (1992), [hep-th/9112049](#).

- [59] A. Ceresole, R. D'Auria, S. Ferrara, W. Lerche and J. Louis, *Picard-Fuchs Equations and Special Geometry*, Int. J. Mod. Phys. **A8**, 79 (1993), [hep-th/9204035](#) $\diamond$ A. Ceresole, R. D'Auria, S. Ferrara, W. Lerche, J. Louis and T. Regge, *Picard-Fuchs Equations, Special Geometry and Target Space Duality*, in: “*Mirror Symmetry II*”, B. R. Greene and S.-T. Yau eds. (American Mathematical Society - International Press, 1997).
- [60] A. Strominger, *Special Geometry*, Commun. Math. Phys. **133**, 163 (1990).
- [61] P. Candelas, M. Lynker and R. Schimmrigk, *Calabi-Yau Manifolds in Weighted $P(4)$* , Nucl. Phys. **B341**, 383 (1990).
- [62] B. R. Greene and M. R. Plesser, *$(2, 2)$ and $(2, 0)$ Superconformal Orbifolds*, Harvard Univ. Rept. HUTP-89/A043 $\diamond$ B. R. Greene and M. R. Plesser, *Duality in Calabi-Yau Moduli Space*, Harvard Univ. Rept. HUTP-89-A043A, Nucl. Phys. **B338**, 15 (1990).
- [63] P. Aspinwall, A. Lütken and G. G. Ross, *Construction and Couplings of Mirror Manifolds*, Phys. Lett. **B241**, 373 (1990).
- [64] A. Giryavets, S. Kachru, P. K. Tripathy, S. P. Trivedi, *Flux Compactifications on Calabi-Yau Threefolds*, JHEP **0404**, 003 (2004), [hep-th/0312104](#).
- [65] F. Denef, *Supergravity Flows and D-Brane Stability*, JHEP **0008**, 050 (2000), [hep-th/0005049](#).
- [66] F. Denef, *On the Correspondence between D-Branes and Stationary Supergravity Solutions of Type II Calabi-Yau Compactifications*, [hep-th/0010222](#).